

Supplementary Material

Supplementary Table 1 showed the levels of cytokines and chemokines produced after PHA-stimulation whole blood from VL patients in the active moment, during the treatment and the end of treatment.

Supplementary table 1. Median concentration (pg/ml) (IQR) of cytokine/chemokine after PHA stimulation of 13 VL patients from Gondar, Ethiopia.

Cytokine/ Chemokine	D0	W1	EOT
IFN- γ	0.1 (0-5.54)	2.52 (0.39-3.71)	3.39 (1.56-15-27) ^{b*}
TNF- α	8.03 (5.19-19-74)	62.00 (8.81-214.3) ^{a**}	206.50 (114.9-726.7) ^{b****, c**}
IP-10	5.19 (0-73.10)	34.26 (1.53-112.90)	122.10 (39.70-842.10) ^{b**}


^a D0 vs W1

^b D0 vs EOT

^c W1 vs EOT

D0: active moment of the disease; W1: one week during the treatment; EOT: end of treatment. * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$.

In the supplementary Figure 1, we showed the fold change in each cytokine/chemokine concentration after one week of treatment (W1) and at the end of the treatment (EOT), in relation to their concentration at the time of active disease (D0) after SLA-stimulation from *L. infantum*.



Supplementary figure 1. Fold changes in cytokine/chemokine concentrations during treatment of 13 VL patients, measured in soluble *L. infantum* antigen stimulated plasma. Fold increase during W1 and EOT compared to D0 was calculated by dividing the value of (W1, EOT) by D0 values. P values are represented for comparison with time of diagnosis (D0). * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.