

Research Article

High-Density Lipoprotein Binds to *Mycobacterium avium* and Affects the Infection of THP-1 Macrophages

Naoya Ichimura,^{1,2} Megumi Sato,¹ Akira Yoshimoto,¹ Kouji Yano,¹ Ryunosuke Ohkawa,¹ Takeshi Kasama,³ and Minoru Tozuka¹

¹Analytical Laboratory Chemistry, Field of Applied Laboratory Science, Graduate School of Health Care Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-Ku, Tokyo 113-8519, Japan

²Department of Clinical Laboratory, Medical Hospital of Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-Ku, Tokyo 113-8519, Japan

³Instrumental Analysis Research Division, Research Center for Medical and Dental Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-Ku, Tokyo 113-8519, Japan

Correspondence should be addressed to Minoru Tozuka; mtozuka.alc@tmd.ac.jp

Received 25 February 2016; Accepted 13 June 2016

Academic Editor: Shinichi Oikawa

Copyright © 2016 Naoya Ichimura et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

High-density lipoprotein (HDL) is involved in innate immunity toward various infectious diseases. Concerning bacteria, HDL is known to bind to lipopolysaccharide (LPS) and to neutralize its physiological activity. On the other hand, cholesterol is known to play an important role in mycobacterial entry into host cells and in survival in the intracellular environment. However, the pathogenicity of *Mycobacterium avium* (*M. avium*) infection, which tends to increase worldwide, remains poorly studied. Here we report that HDL indicated a stronger interaction with *M. avium* than that with other Gram-negative bacteria containing abundant LPS. A binding of apolipoprotein (apo) A-I, the main protein component of HDL, with a specific lipid of *M. avium* might participate in this interaction. HDL did not have a direct bactericidal activity toward *M. avium* but attenuated the engulfment of *M. avium* by THP-1 macrophages. HDL also did not affect bacterial killing after ingestion of live *M. avium* by THP-1 macrophage. Furthermore, HDL strongly promoted the formation of lipid droplets in *M. avium*-infected THP-1 macrophages. These observations provide new insights into the relationship between *M. avium* infection and host lipoproteins, especially HDL. Thus, HDL may help *M. avium* to escape from host innate immunity.

1. Introduction

High-density lipoprotein (HDL), known as antiatherosclerotic lipoprotein, is also involved in innate immunity [1]. For instance, HDL binds to lipopolysaccharides (LPS) and lipoteichoic acid (LTA) derived from microorganisms and neutralizes the physiological activity of these molecules [2, 3]. Additionally, HDL is known to have a bactericidal activity toward *Yersinia enterocolitica* serotype O:3; this activity is mediated by the complement system [4].

Mycobacterium avium (*M. avium*) is a species of nontuberculous mycobacteria and causes opportunistic infections in immunocompromised hosts [5]. Although the incidence of *M. avium* infection is increasing worldwide [6], its

pathogenicity remains poorly understood unlike that of *M. tuberculosis*. Mycobacterial infection is known to be affected by host lipid metabolism. In particular, cholesterol performs an important function in the invasion and survival of mycobacteria inside macrophages [7–11]. Cholesterol accumulation in the host cell membrane is observed at the entry point of *M. tuberculosis* and of other mycobacteria, and cholesterol depletion inhibits the invasion of cells by mycobacteria. After phagocytosis of mycobacteria, they are enveloped by the cell membrane containing cholesterol-rich domains; tryptophan aspartate-containing coat protein (TACO) is recruited to the phagosomes and prevents the fusion of these organelles with lysosomes [12]. Consequently, the engulfed mycobacteria evade degradation by lysosomes

and survive inside the host cell. The invading mycobacteria modulate lipid metabolism in the host cell and promote formation of lipid droplets (LDs), which are mainly composed of neutral lipids such as triacylglycerol and cholesteryl ester [13]. LDs physiologically contribute to lipid storage and lipid metabolism and are available to mycobacteria as a carbon source [10, 14].

Scavenger receptor type B1 (SR-B1), a receptor of apolipoprotein A-I (apoA-I) present in HDL as a main component, is generally known to perform the function of transferring esterified cholesterol from matured HDL into the cytosol in accordance with the cholesterol gradient [15]. SR-B1 was also identified as a nonopsonic phagocytic receptor for mycobacteria because suppression of SR-B1 expression attenuates phagocytosis of mycobacteria [16, 17].

We screened various bacteria for binding with HDL and found that *M. avium* exhibits a stronger interaction with HDL than do other Gram-negative bacilli. The fact that HDL binds to *M. avium* has never been reported and the effect for innate immunity is unclear. The aim of the present study was to elucidate the molecular mechanism of HDL's binding to *M. avium* and the physiological meaning of this interaction.

2. Materials and Methods

2.1. Bacterial Strains and Growth Conditions. *M. avium* (ATCC 700737) was cultured for 3 weeks in Middlebrook 7H9 broth (Difco) supplemented with the oleic acid-albumin-dextrose complex (OADC; Becton Dickinson). Gram-negative bacteria (isolated at the Clinical Laboratory, Medical Hospital of Tokyo Medical and Dental University) were cultured on trypticase-soy-agar with 5% sheep blood (Becton Dickinson) before use.

2.2. Cell Culture, Differentiation, and Infection. The THP-1 cell line was obtained from ATCC (Manassas, VA) and maintained at $2\text{--}10 \times 10^5$ cells/mL in the RPMI 1640 medium (Sigma-Aldrich) supplemented with 10% heat-inactivated fetal bovine serum (Invitrogen), with a penicillin-streptomycin-L-glutamine solution (Wako), and 1x nonessential amino acids (GIBCO). The THP-1 cells were induced to differentiate into THP-1 macrophages by phorbol myristate acetate (PMA, Sigma-Aldrich) for 72 h. After washing with phosphate-buffered saline (PBS), the THP-1 macrophages were incubated in the serum-free RPMI 1640 supplemented with Nutridoma-SP (Roche) and then infected with *M. avium* (multiplicity of infection [MOI] 20:1) with or without HDL (50 μg protein/mL) for 24 h at 35°C in a humidified atmosphere containing 5% CO₂.

2.3. Isolation of HDL and Purification of apoA-I. HDL (1.063–1.210 g/mL) was isolated by means of ultracentrifugation from pooled serum samples obtained from healthy subjects. ApoA-I was purified from HDL according to the previously described method [18]. HDL and apoA-I were dialyzed against PBS and stored at 4°C and –20°C, respectively, until use.

2.4. Western Blot Analysis. Bacteria (3×10^8) were washed with PBS and then incubated with 40 μL of normal human serum (NHS), heat-inactivated serum (HIS), HDL, apoA-I, or PBS for 10 min at 37°C. After washing with PBS, the bacteria were mixed with lysis buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 0.1% SDS, and 0.5% sodium deoxycholate). The lysates were then analyzed by electrophoresis on 12.5% SDS-polyacrylamide gels, and the separated proteins were transferred onto polyvinylidene fluoride (PVDF) membranes (Millipore). ApoA-I was detected with a goat anti-apoA-I polyclonal antibody (Academy Bio-Medical Company) followed by a peroxidase- (POD-) conjugated rabbit anti-goat IgG antibody (Medical & Biological Laboratories). Finally, apoA-I was visualized using hydroperoxide and 3,3'-diaminobenzidine as the substrate.

2.5. Dot Blot Analysis. This analysis was carried out using a previously described method [19] with a slight modification. Lipids were extracted from the outer cell wall of *M. avium* by means of Folch's extraction procedure [20] and were then spotted onto a nitrocellulose membrane. The membrane was sequentially incubated with 5% (w/v) skim milk and 50 $\mu\text{g}/\text{mL}$ HDL in 10 mM Tris-HCl (pH 8.0) containing 140 mM NaCl and 0.1% Tween 20 (TBS-T) for 1 h and 5 h, respectively, at room temperature. The membrane was washed with TBS-T and then incubated with a POD-conjugated anti-apoA-I polyclonal antibody (Cosmo Bio). ApoA-I was visualized as described above.

2.6. Thin Layer Chromatography (TLC) and TLC Blot Analysis. The lipids obtained from *M. avium* were spotted onto three Silica Gel 60 plates (Merck Millipore). The TLC plates were simultaneously developed with CHCl₃/CH₃OH (95:5, v/v). After the run was completed, one of the plates was treated with 20% sulfuric acid solution to visualize the separated lipids, and the other 2 were subjected to TLC blot analysis according to a previously described method [21]. Briefly, the TLC plates were dipped into isopropyl alcohol containing 0.2% aqueous CaCl₂. The separated lipids were thermally transferred onto a PVDF membrane. One membrane was incubated with 1% bovine serum albumin (BSA) in TBS-T and subsequently incubated with 83 $\mu\text{g}/\text{mL}$ of apoA-I in TBS-T at 4°C overnight. The membrane was washed with TBS-T and incubated with the POD-conjugated anti-apoA-I polyclonal antibody (Binding Site). ApoA-I was visualized as described above. To analyze by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS), the lipid bound to apoA-I was extracted from the other membrane by means of CHCl₃/CH₃OH (2:1, v/v) according to comparison with the position of the apoA-I spot on the membrane visualized by the TLC blotting described above.

2.7. Growth Curve Assay. Live *M. avium* cells ($3 \times 10^8/\text{mL}$) were briefly sonicated to disperse clumps and mixed with the Middlebrook 7H9 broth supplemented with OADC with or without HDL (50 μg protein/mL). Absorbance of the medium at 530 nm was monitored for 144 h.




FIGURE 1: Interaction between bacteria and HDL. (a) The bacterial samples (lane 1: *Pseudomonas aeruginosa*, lane 2: *Aeromonas* species, lane 3: *Klebsiella oxytoca*, lane 4: *Klebsiella pneumoniae*, lane 5: *Serratia marcescens*, lane 6: *Citrobacter koseri*, lane 7: *Pseudomonas aeruginosa*, lane 8: *Enterobacter cloacae*, lane 9: *Acinetobacter baumannii*, lane 10: *Stenotrophomonas maltophilia*, lane 11: *Mycobacterium avium*, and lane 12: *Escherichia coli*) were mixed with normal human serum (NHS). The NHS conjugated bacteria were washed and solubilized by means of lysis buffer. The lysates were separated by 12.5% SDS-PAGE and immunoblotted with anti-apoA-I antibody. The molecular weight markers are also shown (M). (b) *M. avium* was incubated with NHS, heat-inactivated serum (HIS), HDL, and apoA-I for 10 min at 37°C. After washing with PBS, *M. avium* was solubilized and subjected to immunoblotting for apoA-I as described above.

2.8. Phagocytosis Assay Using Flow Cytometry. This assay was carried out according to the previously described method [22, 23]. Briefly, *M. avium* was autoclaved and stained with 0.5 mg/mL fluorescein isothiocyanate (FITC, DOJIN laboratories) in PBS for 30 min. After exhaustive washing, we resuspended the stained *M. avium* at 3×10^8 /mL in PBS and then stored the suspension at -80°C until use. THP-1 macrophages (10^6 /well) were cultured with the FITC-conjugated *M. avium* in the presence of HDL (50 μg protein/mL), apoA-I (50 μg /mL), or BSA (50 μg /mL). After quenching extracellular fluorescence using trypan blue (1.2 mg/dL), we fixed the cells with CellFIX (Becton Dickinson) and subjected them to flow cytometric analysis on a Navios flow cytometer (Beckman Coulter). The data were analyzed in the Kaluza software (Beckman Coulter). The FITC-conjugated *E. coli*, which was prepared by the similar method described above, was also analyzed by the phagocytosis assay as the reference.

2.9. Assay of Colony-Forming Units (CFUs). THP-1 macrophages (10^6 /well) were infected with live *M. avium* (MOI 20:1) with or without HDL (50 μg protein/mL) for 24 h. The cells engulfing *M. avium* were washed three times with PBS and further cultured in the absence of *M. avium* and HDL for 0, 24, and 48 h. Then, THP-1 macrophages were lysed with PBS containing 0.1% SDS to recover the engulfed *M. avium*. The lysates diluted 100-fold with PBS were cultured on Middlebrook 7H11 agar plates (Difco) supplemented with OADC. CFUs were counted after 3-week incubation at 37°C.

2.10. Staining and Quantification of LDs. THP-1 macrophages (5×10^5 /well) were infected with live *M. avium* (MOI 20:1) with or without HDL (50 μg protein/mL) for 24 h. After being fixed with a 10% formaldehyde solution for 15 min, they are stained with Oil red O (ORO) for 5 min. The cells were counterstained with Mayer's hematoxylin for 2 min in samples to be examined under a light microscope. Lipid bodies were counted for 50 cells. For quantification of LDs amount, infected THP-1 macrophages (5×10^5 /well) were stained with only ORO as described above. ORO was then extracted in 500 μL of 100% isopropyl alcohol, and absorbance was measured at 540 nm as described previously [24].

2.11. Statistical Analysis. All of the values were shown by the mean \pm standard error of the mean (SEM). Statistical analysis was performed by R software. CFUs, the number of LDs, and the others were statistically analyzed by paired Student's *t*-test, Kruskal-Wallis test, and multifactorial analysis of variance (ANOVA) followed by Tukey post hoc tests, respectively.

2.12. Ethics Statement. Blood samples were obtained from the healthy donors who had provided written informed consent. The protocol of this study was approved by the Research Ethics Committee of Tokyo Medical and Dental University (decision number 1491).

3. Results

3.1. Interaction of *M. avium* with HDL through apoA-I. To screen the bacteria for interaction with HDL, some species of bacteria that were incubated with normal human serum (NHS) were analyzed by western blotting with an anti-apoA-I antibody (Figure 1(a)). The apoA-I signal was well visible in the lysate of *M. avium* incubated with NHS but only barely visible in samples from other Gram-negative bacteria. To confirm the direct interaction between apoA-I and *M. avium*, we carried out western blotting of the lysate of *M. avium* incubated with heat-inactivated serum (HIS), NHS, isolated HDL, or purified apoA-I (Figure 1(b)). The apoA-I signal was well visible after incubation not only with NHS but also with the HIS, HDL, and apoA-I, indicating that HDL bound directly to *M. avium* via apoA-I.

3.2. ApoA-I Binds to the Outer Cell Wall Lipid of *M. avium*. We hypothesized that apoA-I interacts with the abundant lipids on the *M. avium* cell wall because of amphipathic α -helices present in the apoA-I molecule. To test this hypothesis, we performed a western blot assay of the lipids extracted from the outer cell wall of *M. avium* (Figure 2(a)). ApoA-I was detected in the spots of the extracted lipids incubated with HDL and the intensity of signals increased with the amount of spotted lipids (at a constant HDL concentration). The TLC blot assay was carried out to determine whether apoA-I binds to a specific lipid molecule of *M. avium* (Figure 2(b)). Several lipid spots on the TLC plate were nonspecifically visualized by a charring reagent (Figure 2(b), left). One




FIGURE 2: Interaction between lipids of *M. avium* and HDL. (a) The lipids extracted from *M. avium* were spotted in the indicated amount (12.5–100.0 μg) on a nitrocellulose membrane. The membrane was then incubated with HDL (50 μg protein/mL), and HDL bound to the lipids was visualized using an anti-apoA-I antibody. (b) TLC plates with the spotted lipids (150 μg each sample) extracted from *M. avium* were developed using $\text{CHCl}_3/\text{CH}_3\text{OH}$ (95:5, v/v). One plate was treated with a 20% H_2SO_4 solution to visualize the fractionated lipids (TLC). The fractionated lipids on the other plate were thermally transferred onto a PVDF membrane, which was then incubated with HDL (83 μg protein/mL). The lipid that bound to HDL was visualized using an anti-apoA-I antibody (TLCB).

of those spots visibly interacted with apoA-I by TLC blot analysis (Figure 2(b), right), indicating that apoA-I formed a complex with this specific lipid of *M. avium*.

The lipids extracted from the apoA-I-positive spot of the TLC blot were analyzed by MALDI-TOF MS (Supplemental Figure 1 in Supplementary Material available online at <http://dx.doi.org/10.1155/2016/4353620>). Two prominent peaks at m/z 1243.8 and m/z 1271.8 were observed along with the several peaks with an interval of 28 Da, suggesting that the lipid that bound to apoA-I included a variety of fatty acid moieties.

3.3. Analysis of Bactericidal Activity of HDL. The growth curve analysis of *M. avium* in the presence or absence of HDL was performed to determine whether HDL directly kills *M. avium* (Figure 3). No significant difference was observed in the absorbance of the medium with or without HDL for 6 days. This finding indicated that HDL itself had no bactericidal activity toward *M. avium*.

3.4. Effects of HDL on Engulfment of *M. avium* by THP-1 Macrophages. To find out whether HDL participates in the nonopsonic phagocytosis of *M. avium* by THP-1 macrophages, we used flow cytometry to analyze the effect of HDL on the number of THP-1 macrophages engulfing FITC-conjugated *M. avium* (Figure 4(a)). The percentage of FITC-positive cells decreased with the addition of HDL and apoA-I but did not change significantly after addition of BSA. The percentage of FITC-positive cells which is obtained using FITC-conjugated *E. coli* indicated a similar result to that of FITC-conjugated *M. avium*; however the addition of HDL did not decrease the percentage of FITC-positive cells. When the number of FITC-positive cells of control samples was set




FIGURE 3: Bactericidal activity of HDL. *M. avium* was cultured with (+HDL; 50 μg protein/mL) or without (control) HDL and absorbance (at 530 nm) of the medium was measured at 0, 24, 48, 72, and 144 h. The values were indicated by mean \pm SEM ($n = 3$).

to 100%, HDL and apoA-I decreased the number of FITC-positive cells to $68.7\% \pm 1.5\%$ and $83.3\% \pm 0.4\%$, respectively (Figure 4(b)). BSA did not affect the engulfment of *M. avium* by THP-1 macrophages ($101.8\% \pm 2.0\%$). In contrast, FITC-positive cells did not decrease, if anything increased, in the case of FITC-conjugated *E. coli*.




FIGURE 4: Effects of HDL on the engulfment of *M. avium* by THP-1 macrophages. (a) THP-1 macrophages (10^6 /well) were cultured with FITC-conjugated *M. avium* or *E. coli* in the presence or absence (control) of HDL ($50 \mu\text{g}$ protein/mL), apoA-I ($50 \mu\text{g}$ /mL), or BSA ($50 \mu\text{g}$ /mL) for 24 h. After extensive washing and quenching of extracellular bacteria, the cells were subjected to flow cytometric analysis. Representative scatter profiles are presented. (b) The percentages of FITC-positive cells compared to the control using FITC-*M. avium* or FITC-*E. coli* were quantitatively compared. The values were indicated by mean \pm SEM ($n = 3$, * $P < 0.05$, ** $P < 0.01$). NS: not significant.

3.5. Effects of HDL on Bactericidal Activity of THP-1 Macrophages. Assay of CFUs was performed to assess the number of live *M. avium* inside THP-1 macrophages by means of *M. avium* recovered from the lysed THP-1 macrophages (Figure 5(a)). The results that were expressed in CFU/well were $(8.5 \pm 1.2) \times 10^4$ and $(11.7 \pm 0.6) \times 10^4$ (mean \pm SE) in the presence and the absence of HDL, respectively. This observation is consistent with the result obtained by the flow cytometric analysis (Figure 4), namely, that HDL attenuated the engulfment of *M. avium* by THP-1 macrophages. To assess the survival rate of *M. avium* inside THP-1 macrophages, the infected THP-1 macrophages were further cultured in the absence of *M. avium* and HDL for 24 and 48 h. The number of CFUs per well indicated a tendency to decrease in a time-dependent manner (Figure 5(b)). However, no significant difference was observed in the decreasing profiles between the presence and the absence of HDL, suggesting that the coexistence of HDL could not affect bacterial killing after ingestion by THP-1 macrophage.

3.6. Effects of HDL on LDs Formation. After the infection of THP-1 macrophages with live *M. avium*, LDs inside the

cells were stained with Oil red O (ORO) (Figure 6(a)). The numbers of LDs per cell were significantly increased by the coexistence of live *M. avium* and HDL (Figure 6(b)). LDs amounts were also estimated by the absorbance at 540 nm of the extracted ORO (Figure 6(c)). The existence of HDL or *M. avium* alone indicated no significant effect in the quantity of ORO compared with the control. However, the coexistence of HDL and *M. avium* caused an increase in the LDs amount.

4. Discussion

In this study, we found that HDL binds to *M. avium* via the interaction between apoA-I (the main structural component of HDL) and a specific lipid molecule of *M. avium*. The specific lipid could be extremely nonpolar considering its behavior during TLC analysis. Although a structure of the specific lipid molecule was analyzed by MALDI-TOF-MS/MS (data not shown), we failed to identify it among the known lipid molecules of *M. avium*.

It should be noted that *M. avium* causes inflammation and increases permeability of blood vessels, increasing the chance of interaction with leaked HDL. If this specific lipid of *M.*




FIGURE 5: Effects of HDL on the viability of *M. avium* engulfed by THP-1 macrophages. (a) THP-1 macrophages (10^6 /well) were infected with live *M. avium* in the presence or absence (control) of HDL ($50 \mu\text{g}$ protein/mL) for 24 h. *M. avium* was recovered from lysed THP-1 macrophages for an assay of colony-forming units (CFUs; 3 weeks culture). (b) A part of infected THP-1 macrophages was further cultured for 24 and 48 h in a fresh medium. CFUs were also determined as described above. The data were indicated as the percentages against CFUs obtained without further cultivation. The values were indicated by mean \pm SEM ($n = 3$, * $P < 0.05$).

avium plays a role in host pathogenesis, HDL may neutralize the activity of this lipid, just as in the case of LPS and LTA. It could be important to identify the lipid molecule in order to elucidate a participation of HDL in *M. avium* infection.

We explored the influence of HDL's binding to *M. avium* on an innate immunity in experiments with THP-1 macrophages. HDL does not directly kill *M. avium*, but the presence of HDL significantly decreases the number of THP-1 macrophages engulfing *M. avium*, and HDL indicated no effect on bacterial killing after ingestion. It suggests that HDL allows *M. avium* to escape engulfment by THP-1 macrophages. SR-B1, known as an apoA-I receptor, also plays a role in nonopsonic phagocytosis of mycobacteria [16, 17]. According to these observations, we can hypothesize that HDL competitively inhibits the binding of *M. avium* to SR-B1. In this study, we did not analyze SR-B1, and further research is needed to assess the interaction between *M. avium* and HDL, including the participation of SR-B1.

When mycobacteria are recognized by Toll-like receptors 2 or 6 (TLR2/6), LDs formation is enhanced by TLR signaling [25, 26]. Moreover, some researchers proposed a mechanism behind the enhancement of LDs formation during mycobacterial infection: upregulation of a series of scavenger receptors including SR-B1 [27]. In our study, *M. avium* infection did not apparently upregulate LDs formation under serum-free condition; however, the coexistence of

HDL increased the number and the amount of LDs in THP-1 macrophages, suggesting that HDL-cholesterol is utilized as a raw material for LDs formation. Two mechanisms may explain this observation. According to one mechanism, HDL-cholesterol is internalized together with *M. avium* via phagocytosis. According to the other, although approximately 30% of the engulfment of *M. avium* by THP-1 macrophages was attenuated by the coexistence of HDL (Figure 4), the HDL-cholesterol influx is enhanced by upregulation of SR-B1 in THP-1 macrophages engulfing *M. avium*. Consequently, TLR signaling and HDL-cholesterol influx via SR-B1 may promote LDs formation in the presence of both *M. avium* and HDL. Further research is needed to determine the actual mechanism.

On the basis of the present study, we believe that HDL plays a crucial role in the infection of THP-1 macrophages by *M. avium*. Identification of the lipid molecule of *M. avium* that binds to HDL via apoA-I and the analysis of the expression of the scavenger receptor, SR-B1, in an upcoming project, should shed some light on the pathogenesis of *M. avium* infections and on the relevant immune response.

5. Conclusion

HDL affects *M. avium* infection through the binding between apoA-I, the main component of HDL, and a specific lipid of




FIGURE 6: High-density lipoprotein (HDL) promotes a formation of THP-1 macrophage-derived foam cells during *M. avium* infection. THP-1 macrophages were infected by *M. avium* with (+) or without (-) HDL. (a) Representative light microscopic images of the cells stained with Oil red O (ORO) and hematoxylin are shown. The scale bar is 25 nm. (b) The numbers of lipid droplets (LDs) per cell were calculated by counting of those number for 50 cells. (c) The amount of LDs was also quantitatively analyzed by measuring absorbance (at 540 nm) of ORO extracted from the cells. The values were indicated by mean \pm SEM ($n = 3$, * $P < 0.05$, ** $P < 0.01$). NS: not significant.

M. avium. This interaction may enable *M. avium* to escape from innate immunity THP-1 macrophages.

Competing Interests

The authors declare that there are no competing interests regarding the publication of this paper.

Acknowledgments

The authors thank Dr. Naoko Tojo (Department of Clinical Laboratory, Medical Hospital of Tokyo Medical and Dental University) for generously helping their collaborators and coauthors. They also thank Dr. Etsuko Sawabe (Department of Clinical Laboratory, Medical Hospital of Tokyo Medical

and Dental University) for generously providing assistance with handling of the microorganisms.

References

- [1] K. R. Feingold and C. Grunfeld, "The role of HDL in innate immunity," *Journal of Lipid Research*, vol. 52, no. 1, pp. 1–3, 2011.
- [2] L. Cai, A. Ji, F. C. de Beer, L. R. Tannock, and D. R. van der Westhuyzen, "SR-BI protects against endotoxemia in mice through its roles in glucocorticoid production and hepatic clearance," *The Journal of Clinical Investigation*, vol. 118, no. 1, pp. 364–375, 2008.
- [3] J. H. M. Levels, P. R. Abraham, E. P. Van Barneveld, J. C. M. Meijers, and S. J. H. Van Deventer, "Distribution and kinetics of lipoprotein-bound lipoteichoic acid," *Infection and Immunity*, vol. 71, no. 6, pp. 3280–3284, 2003.
- [4] M. Biedzka-Sarek, J. Metso, A. Katefides et al., "Apolipoprotein A-I exerts bactericidal activity against *Yersinia enterocolitica* serotype O:3," *The Journal of Biological Chemistry*, vol. 286, no. 44, pp. 38211–38219, 2011.
- [5] I. M. Orme and D. J. Ordway, "Host response to nontuberculous mycobacterial infections of current clinical importance," *Infection and Immunity*, vol. 82, no. 9, pp. 3516–3522, 2014.
- [6] C. H. Weiss and J. Glassroth, "Pulmonary disease caused by nontuberculous mycobacteria," *Expert Review of Respiratory Medicine*, vol. 6, no. 6, pp. 597–613, 2012.
- [7] D. A. Keown, D. A. Collings, and J. I. Keenana, "Uptake and persistence of *Mycobacterium avium* subsp. paratuberculosis in human monocytes," *Infection and Immunity*, vol. 80, no. 11, pp. 3768–3775, 2012.
- [8] P. Peyron, C. Bordier, E.-N. N'Diaye, and I. Maridonneau-Parini, "Nonopsonic phagocytosis of *Mycobacterium kansasii* by human neutrophils depends on cholesterol and is mediated by CR3 associated with glycosylphosphatidylinositol-anchored proteins," *Journal of Immunology*, vol. 165, no. 9, pp. 5186–5191, 2000.
- [9] J. Gatfield and J. Pieters, "Essential role for cholesterol in entry of mycobacteria into macrophages," *Science*, vol. 288, no. 5471, pp. 1647–1651, 2000.
- [10] K. A. Mattos, V. C. G. Oliveira, M. Berrêdo-Pinho et al., "Mycobacterium leprae intracellular survival relies on cholesterol accumulation in infected macrophages: a potential target for new drugs for leprosy treatment," *Cellular Microbiology*, vol. 16, no. 6, pp. 797–815, 2014.
- [11] C. de Chastellier and L. Thilo, "Cholesterol depletion in *Mycobacterium avium*-infected macrophages overcomes the block in phagosome maturation and leads to the reversible sequestration of viable mycobacteria in phagolysosome-derived autophagic vacuoles," *Cellular Microbiology*, vol. 8, no. 2, pp. 242–256, 2006.
- [12] K. Tanigawa, K. Suzuki, H. Kimura et al., "Tryptophan aspartate-containing coat protein (CORO1A) suppresses Toll-like receptor signalling in *Mycobacterium leprae* infection," *Clinical and Experimental Immunology*, vol. 156, no. 3, pp. 495–501, 2009.
- [13] K. A. Mattos, F. A. Lara, V. G. C. Oliveira et al., "Modulation of lipid droplets by *Mycobacterium leprae* in Schwann cells: a putative mechanism for host lipid acquisition and bacterial survival in phagosomes," *Cellular Microbiology*, vol. 13, no. 2, pp. 259–273, 2011.
- [14] A. Brzostek, J. Pawelczyk, A. Rumijowska-Galewicz, B. Dziadek, and J. Dziadek, "Mycobacterium tuberculosis is able to accumulate and utilize cholesterol," *Journal of Bacteriology*, vol. 191, no. 21, pp. 6584–6591, 2009.
- [15] S. Xu, M. Laccotripe, X. Huang, A. Rigotti, V. I. Zannis, and M. Krieger, "Apolipoproteins of HDL can directly mediate binding to the scavenger receptor SR-BI, an HDL receptor that mediates selective lipid uptake," *Journal of Lipid Research*, vol. 38, no. 7, pp. 1289–1298, 1997.
- [16] J. A. Philips, E. J. Rubin, and N. Perrimon, "Drosophila RNAi screen reveals CD36 family member required for mycobacterial infection," *Science*, vol. 309, no. 5738, pp. 1251–1253, 2005.
- [17] G. Schäfer, R. Guler, G. Murray, F. Brombacher, and G. D. Brown, "The role of scavenger receptor BI in infection with mycobacterium tuberculosis in a murine model," *PLoS ONE*, vol. 4, no. 12, Article ID e8448, 2009.
- [18] Y. Usami, K. Matsuda, M. Sugano et al., "Detection of chymase-digested C-terminally truncated apolipoprotein A-I in normal human serum," *Journal of Immunological Methods*, vol. 369, no. 1–2, pp. 51–58, 2011.
- [19] J. M. Stevenson, I. Y. Perera, and W. F. Boss, "A phosphatidylinositol 4-kinase pleckstrin homology domain that binds phosphatidylinositol 4-monophosphate," *Journal of Biological Chemistry*, vol. 273, no. 35, pp. 22761–22767, 1998.
- [20] J. Folch, M. Lees, and G. H. Sloane Stanley, "A simple method for the isolation and purification of total lipides from animal tissues," *The Journal of Biological Chemistry*, vol. 226, no. 1, pp. 497–509, 1957.
- [21] T. Taki and D. Ishikawa, "TLC blotting: application to microscale analysis of lipids and as a new approach to lipid-protein interaction," *Analytical Biochemistry*, vol. 251, no. 2, pp. 135–143, 1997.
- [22] M. J. Hewish, A. M. Meikle, S. D. Hunter, and S. M. Crowe, "Quantifying phagocytosis of *Mycobacterium avium* complex by human monocytes in whole blood," *Immunology & Cell Biology*, vol. 74, no. 4, pp. 306–312, 1996.
- [23] J. Nuutila and E.-M. Lilius, "Flow cytometric quantitative determination of ingestion by phagocytes needs the distinguishing of overlapping populations of binding and ingesting cells," *Cytometry Part A*, vol. 65, no. 2, pp. 93–102, 2005.
- [24] Q. Lin, Y.-J. Lee, and Z. Yun, "Differentiation arrest by hypoxia," *Journal of Biological Chemistry*, vol. 281, no. 41, pp. 30678–30683, 2006.
- [25] K. A. Mattos, V. G. C. Oliveira, H. D'Avila et al., "TLR6-driven lipid droplets in *Mycobacterium leprae*-infected Schwann cells: immunoinflammatory platforms associated with bacterial persistence," *Journal of Immunology*, vol. 187, no. 5, pp. 2548–2558, 2011.
- [26] H. D'Avila, R. C. N. Melo, G. G. Parreira, E. Werneck-Barroso, H. C. Castro-Faria-Neto, and P. T. Bozza, "Mycobacterium bovis bacillus Calmette-Guérin induces TLR2-mediated formation of lipid bodies: Intracellular domains for eicosanoid synthesis in vivo," *Journal of Immunology*, vol. 176, no. 5, pp. 3087–3097, 2006.
- [27] A. A. Elamin, M. Stehr, and M. Singh, "Lipid droplets and *Mycobacterium leprae* infection," *Journal of Pathogens*, vol. 2012, Article ID 361374, 10 pages, 2012.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

