

## Research Article

# Assessment of Sodium and Potassium Intakes in Children Aged 6 to 18 Years by 24 h Urinary Excretion in City of Rabat, Morocco

Naima Saeid <sup>1</sup>, Mohammed Elmzibri <sup>1</sup>, Abdeslam Hamrani <sup>1</sup>, Qandoussi Latifa,<sup>1</sup> Hakim Belghiti,<sup>2</sup> Hicham El Berri,<sup>3</sup> Kaoutar Benjeddou,<sup>1</sup> Amina Bouziani <sup>1</sup>, Hasnae Benkirane <sup>1</sup>, Youness Taboz <sup>1</sup>, Asmae Elhamdouchi <sup>1</sup>, Khalid El Kari <sup>1</sup>, and Hassan Aguentaou<sup>1</sup>

<sup>1</sup>Joint Research Unit in Nutrition and Food, URAC 39, (Ibn Tofail University-CNESTEN), Regional Designated Center of Nutrition (AFRA/IAEA), Rabat-Kénitra, Morocco

<sup>2</sup>Military Hospital Mohammed V, Rabat, Morocco

<sup>3</sup>Ministry of Health, Rabat, Morocco

Correspondence should be addressed to Naima Saeid; saeid\_na@yahoo.com

Received 10 February 2018; Accepted 21 May 2018; Published 1 August 2018

Academic Editor: Michael B. Zemel

Copyright © 2018 Naima Saeid et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** The incidence of noncommunicable diseases (NCDs) has greatly increased, mainly due to high level of dietary sodium. Thus, reduction of sodium intake in population has been recognized as one of the most cost-effective strategies to reduce NCDs. The aim of this study was to estimate sodium and potassium consumption in a sample of Moroccan children as a baseline study to implement national strategy for salt intake reduction. **Methods.** The study was conducted on 131 children aged 6–18 years recruited from Rabat and its region. Sodium excretion and potassium excretion were measured on 24 h urinary collection, and the creatinine excretion was used to validate completeness of urine collections. **Results.** The average of urinary sodium was  $2235.3 \pm 823.2$  mg/day, and 50% of children consume more than 2 g/d of sodium (equivalent to 5 g/day of salt), recommended by the WHO. However, daily urinary excretion of potassium was  $1431 \pm 636.5$  mg/day, and 75% of children consume less than adequate intake. Sodium consumption increased significantly with age. Of particular interest, 46.7% of children aged 6–8 years and 49.3% of children aged 9–13 years consume more than the corresponding upper limits. **Conclusions.** Children have high sodium and low potassium status. There is evidence of the urgent need to implement a strategy for reduction of dietary sodium intake in Morocco.

## 1. Background

Worldwide, noncommunicable diseases (NCDs) are the leading cause of mortality and morbidity [1] accounting globally for 60% of all deaths and 43% of disease burden [2]. Evidence that diet is an important risk factor for NCDs has been reported by various researchers. Indeed, Western dietary patterns involving an over consumption of fatty, sugary, and salty foods proliferate in many emerging and developing countries [3].

Nowadays, it is widely accepted that an excessive intake of salt is associated with a broad range of NCDs, such as

hypertension, cardiovascular diseases (CVDs), cancer, and osteoporosis [4, 5].

Most individuals suffering from higher blood pressure have been shown to have high sodium and low potassium consumptions [6]. In fact, several studies have shown a stronger relation between sodium/potassium ratio and blood pressure than sodium or potassium alone. Moreover, a high consumption of potassium, commonly from fruits and vegetables, can counteract the negative effects of high sodium intake on blood pressure [7, 8]. On the other hand, numerous studies have shown that blood pressure levels in infancy and childhood are moderately predictive for blood pressure later in life [9, 10].

Blood pressure in childhood or early adult life rises progressively through to middle age and predicts both the blood pressure level and presence of hypertension in later life [11]. So tracking of dietary habits from early childhood into adulthood has shown that children with extremely high levels of sodium and lower potassium intakes tend to maintain those levels over time [12, 13].

Therefore, diet in childhood can be a significant determinant of adult dietary habits even after several decades [14].

The World Health Organization (WHO) recommendation outlines the importance of prioritizing sodium intake reduction, as a main approach to reduce blood pressure and to decrease the risk of CVDs and strokes. Accordingly, the WHO recommends for adults the consumption of less than 2 g of sodium/day (5 g/day of salt) and more than 3.5 g of potassium/day [15]. For children, the reduction of sodium intake prevents and decreases risk of hypertension during childhood and adulthood. In this case, the maximum recommended level of sodium intake should be adjusted downward based on the energy requirements of children relative to those of adults [16]. Following these recommendations, the adequate intake for sodium for young people is set at 1.5 g/day (equivalent to 3.8 g of salt) without reducing the intake of other nutrients. The Institute of Medicine (IOM) sets the tolerable upper limit (UL) intake level at 2300 mg per day. The UL is the highest daily nutrient intake level that is likely to pose no risk of adverse health effects (e.g., for sodium, increased blood pressure). The UL depends on the age; for children aged from 6 to 8 years, the UL is set at 1900 mg/day, and for children aged from 9 to 13 years and 14 to 18 years, the ULs are set at 2200 mg/day and 2300 mg/day, respectively. For instance, the IOM has reported that the adequate intake of potassium is 3.8 g/day for children aged 6–8 years and reaches 4.7 g/day for adolescents and adults [17].

Sodium intake can be estimated indirectly either from a questionnaire or food consumption data or directly by the measurement of urinary excretion. Because of the problems of underestimation of sodium intakes based on dietary surveys in most studies, 24-hour urinary sodium excretion has become the “gold standard” method of obtaining data on sodium and potassium intakes in population surveys [18]. However, the assessment of daily intakes by using 24 h dietary recall (three times) provides useful additional information on the nature of food consumed and helps to implement specific strategies based on studied population consumption.

However, fewer data are available on urinary sodium excretion in children and young people than that in adults, and these are mainly limited to the developed countries such as Europe and North America [19, 20].

This is mainly due to methodological difficulties in obtaining complete and valid urinary excretion for children [21]. The majority of studies on sodium and potassium consumption in children are based on food consumption, which is still difficult in this case.

In Morocco, high blood pressure is a public health problem; its prevalence was estimated at 33% according to the national survey carried out in 2000, and this prevalence is expected to continue to rise [22]. Thus, reduction of salt

consumption would be a suitable strategy to fight against NCDs in Morocco. On the other hand, data on the daily intake of salt in Morocco through the diet of Moroccans remain limited. The most recent estimate was carried out by the Ministry of Health on 2008 before the addition of iodine to salt and revealed that the salt intake for adults reached 7–12 g/person/day [23].

Therefore, we have planned in this study to assess the consumption of sodium and potassium among a sample of Moroccan children as a part of a baseline survey (pilot study) to adapt and implement the WHO strategy to reduce the salt intake to the Moroccan context by using the 24 h urinary sodium/potassium excretion.

## 2. Materials and Methods

**2.1. Subjects.** This transversal study was conducted between September 2015 and June 2016. It was carried out among children aged 6–18 years, enrolling two primary schools and one middle school in Rabat and its nearest region.

The city of Rabat is the political and administrative capital of Morocco, and it is located on the Atlantic coast. The region of Rabat accounts about 1.6 million inhabitants, thus becoming the second largest agglomeration of the country after Casablanca [24]. The unemployment rate is 22.5% and the illiteracy is about 20.7% [25], whereas 16% of the population of Rabat are affected by poverty [26].

The study protocol was approved by the Ethics Committee for Biomedical Research, Faculty of Medicine and Pharmacy of Rabat, Morocco, and written informed consent was obtained from each parent of the recruited child.

**2.2. Clinical Survey.** A face-to-face interview was conducted with the participant's parents to collect information on health problems (exclusion criteria), date of birth, and consumption of medications and/or supplements.

**2.3. Anthropometric Measurements.** The anthropometric measurements were taken at schools according to the WHO recommendations [27]. Weight was measured using an electronic scale to the nearest 0.1/0.2 kg (Seca GmbH, Germany). Height was measured in the standing position, barefoot, using Shorr Board portable at the nearest of 0.1 cm (formerly Shorr Productions, LLC, USA). Waist circumference (WC) was measured at 0.1 cm threshold using an inelastic tape measure. The nutritional status was evaluated by different Z-scores (BMI Z-score, height-for-age Z-score, weight-for-age Z-score, and weight-for-height Z-score) calculated by the Anthroplus software [28].

The weight-for-age curve enables countries that routinely measure only weight to monitor growth throughout childhood. BMI-for-age is the recommended indicator for assessing thinness, overweight, and obesity in children aged 10–19 years.

**2.4. Blood Pressure.** Blood pressure was measured using an automatic vital signs monitor (OMRON M6 Comfort-HEM-7321 E). Measurements were completed in the sitting

position after 10 minutes of rest with the child's arm positioned at the level of the heart on a rested table. The cuff was positioned on the child's arm by aligning the marked cuff artery indicator with the brachial artery. Three blood pressure readings were taken on the right arm at 1-minute intervals. The average of the three measurements will be used for analysis.

**2.5. Urine Sampling.** A single-timed 24-hour urine collection was obtained for estimation of electrolyte excretion. Instructions on urine collection were provided for parents, and simplified pictorial instructions were given to children. As requested, all participants were invited to discard the overnight urine to start the urine collection period with an empty bladder. During the following 24 hours, all urine voided was collected in 5-liter wide-mouth, rimmed polypropylene bottles. Start and finish urine collection times and any missed collections were also recorded on the 24-hour urine bottle. If the duration of the collection was not exactly 24 hours (but within 20–28 hours), urinary sodium, potassium, creatinine, and total volume were normalized to a 24-hour period [29].

Moreover, urine collections were being considered incomplete and then excluded if (1) the volume is less than 300 ml; (2) more than few drops of urine were lost during urine collection; (3) time outside of <20 hours or >28 hours; and (4) urinary creatinine was <280 mg/l or >2590 mg/l. In these cases, participants were excluded or invited to repeat the urine collection.

**2.6. Urine Analysis.** The total volume of urine collected was measured using a specially devised linear measuring scale, and an aliquot (20 ml) from the complete samples was stored at  $-20^{\circ}\text{C}$  until analysis. Urinary sodium and potassium concentrations were assessed using inductively coupled plasma mass spectrometry (ICP-MS; Thermo Scientific X-SERIES 2); coefficient of variation was 1.5% for sodium and 2.5% for potassium. An international reference material (Seronorm TM Trace Elements Urine) was used to control and validate the results. Total sodium and potassium excretions during 24 hours are calculated by multiplying the obtained concentrations by the volume of the collected urine and reported to 24-hour urine excretion [30]. Daily sodium and potassium intakes were expressed in grams, and sodium concentrations were converted to the salt equivalent (g) using the conversion factor 2.54 [21].  $\text{Na}^+/\text{K}^+$  ratio was also calculated.

Urinary creatinine excretion was analyzed by the kinetic approach according to Jaffé method [31], using the Cobas C311 (Roche Diagnostic, Meylan, France).

**2.7. Statistical Analysis.** All statistical analyses were carried out using SPSS for Windows, version 20 (SPSS Inc, Chicago). Variables were expressed as mean  $\pm$  standard deviation (SD) or as interquartile range. The normality of data was checked, and the association between sodium intake and the studied variables was assessed by the Kolmogorov–Smirnov test. The

significance level was established as 5% ( $<0.05$ ). Independent samples *t*-test was used to compare boys with girls within each age group for continuous data. The percentages of children above the maximum salt intake recommendations are reported.

The distributions of children by age groups were made according to the Institute of Medicine (IOM) children stratification [17].

### 3. Results

**3.1. Participants.** The flow diagram of the study is reported in Figure 1. A total of 205 children were recruited from the 3 schools. Among them, 10 were excluded for (1) non-attendance on days when 24 h urinary collection was collected; (2) having a disease that might affect the results (diabetes, renal disease, etc.); (3) being under special diet; and (4) taking antidiuretic drugs the week previous to the study. Complete information regarding clinical survey, anthropometric data, and 24 h urine samples was obtained from 195 eligible children. Among them, 64 children were withdrawn for nonvalid urine sampling: 5 children have reported that the collection time was less than 20 hours or more than 28 hours; 36 children have abnormal creatinine values; 6 children have collected less than 300 ml of urine; and 5 children have reported that they missed more than 2 times of collection. Therefore, only 131 participants have been retained in the study.

**3.2. Characteristics of Study Participants.** In our study, boys represent 51.9% (68/131) and girls 48.1% (63/131), with a sex ratio of 1.08. The mean age was  $9.8 \pm 2.4$  years. Among the 131 children, 19.1% aged 6–8 years (25/131), 68.7% aged 9–13 years (90/131), and 12.2% aged 14–18 years (16/131) (Table 1).

The characteristics of growth parameters for the studied children are presented in Table 1. For overall recruited children, the mean weight was  $32.7 \pm 11.3$  kg and the average height was  $137 \pm 14.4$  cm. For nutritional status, the BMI-Z-score indicates that 9.9% children are underweight, 12.2% present overweight status, and 8.4% are obese. Statistical analysis showed no significant difference between boys and girls for all parameters.

**3.3. Assessment of Sodium and Potassium Status.** Excretion of urinary sodium and potassium is summarized in Table 2. The average of urine creatinine excretion was  $852.8 \pm 352.7$  mg/day. The mean of daily urinary excretion of sodium in the whole group was  $2235.3 \pm 823.2$  mg/day (equivalent to  $5667.9 \pm 2077.7$  mg/day of salt). Overall, 50% of children consume more than 2 g/day of sodium (equivalent to 5 g/day of salt). However, daily urinary excretion of potassium was  $1431 \pm 636.5$  mg/day, and 75% of children consume less than adequate intake. The mean sodium-to-potassium ratio was  $1.7 \pm 0.7$ , so children consume double the amount of the adequate intake. A comparison of daily sodium and potassium urinary excretion, sodium/potassium ratio, and creatinine level between boys




FIGURE 1: Flowchart of the study.

TABLE 1: General characteristics of the studied population.

	Total (n = 131)		Boys (n = 68)		Girls (n = 63)		p values
	Mean ± SD	Median (interquartile range)	Mean ± SD	Median (interquartile range)	Mean ± SD	Median (interquartile range)	
Age (years)	9.8 ± 2.4	9.7 (8.4, 11.2)	9.8 ± 2.4	9.4 (8.1, 11.0)	10.5 ± 2.6	9.9 (8.9, 11.4)	0.1
Weight (kg)	32.7 ± 11.3	30.4 (25.0, 38.4)	32.7 ± 11.3	29.5 (24.8, 39.4)	34.1 ± 12.4	31.6 (25.9, 37.9)	0.4
Height (cm)	137 ± 14.4	137.0 (130, 144.9)	137 ± 14.4	137.2 (126.7, 144.0)	138.5 ± 13.8	136 (130, 148)	0.5
BMI	17.1 ± 3.6	16.5 (14.6, 18.6)	17.1 ± 3.8	16.08 (14.5, 18.2)	17.3 ± 3.5	16.8 (14.8, 18.8)	0.7
BAZ	-0.1 ± 1.6	0.05 (-1.1, 0.7)	-0.1 ± 1.8	0.05 (-1.1, 0.75)	-0.15 ± 1.3	0.03 (-1.9, 0.9)	0.3
Thinness	9.9	—	10.3	—	9.5	—	
Normal	69.5	—	72.1	—	66.7	—	
Overweight	12.2	—	8.8	—	15.9	—	0.8
Obesity	8.4	—	8.8	—	7.9	—	
Blood pressure							
Systolic (mmHg)	98.5 ± 11.7	98 (90–107)	98.1 ± 12	98.3 (88.4–108.4)	98.9 ± 10.1	99 (91.3–107)	0.7
Diastolic (mmHg)	69.3 ± 9.9	70 (61–77.5)	68.8 ± 10.7	70 (60–77.8)	69.8 ± 9.01	70 (62–77.3)	0.5

p values by one-way ANOVA for means or Mann-Whitney U test for medians. Results are presented as mean ± standard deviation or proportion (%); BMI (body mass index), BAZ (BMI Z-score of body mass index for age), and Z-scores were determined according to [27].

and girls are also reported in Table 2 and showed no significant statistical difference.

Table 3 provides the distribution of sodium intake according to age, gender, and nutritional status. Data clearly showed that for all age groups, sodium intake exceeds the upper limits (ULs) according to ages' energetic needs. Sodium intake increases with age, children aged 6–8 years consume an average of 1800.0 (1450.1, 2145.3) mg/day, and 46.7% of children consume over the UL. Children aged 9–13

years consume 2193.4 (1843.6, 2793.8) mg/day, and more than 49.3% have over UL. Finally, 26.7% of children aged 14–18 years have over ULs and consume an average of 2138.0 (1876.7, 2392.5) mg/day. Differences between age groups are statistically significant ( $p < 0.001$ ). Distribution of sodium intake and the corresponding proportion of children exhibiting sodium status over the UL level, according to gender and nutritional status, showed no statistically significant difference ( $p < 0.05$ ).

TABLE 2: Urinary sodium and potassium excretion according to the sex.

	Total ( <i>n</i> = 131)		Boys ( <i>n</i> = 68)		Girls ( <i>n</i> = 63)		<i>p</i> value
	Mean ± SD	Median (interquartile range)	Mean ± SD	Median (interquartile range)	Mean ± SD	Median (interquartile range)	
Sodium (mg/d)*	2235.3 ± 823.2	2094.2 (1640.3, 2498.0)	2184.3 ± 783.3	2057.0 (1600.0, 2462.5)	2290.3 ± 867.2	2138.0 (1765.3, 2669.1)	0.4
Potassium (mg/d)	1431 ± 636.5	1251.6 (994.3, 1813.0)	1469.4 ± 721.3	1217.7 (902.9, 1940.6)	1389.5 ± 532.8	1288.3(1022.6, 1288.3)	0.47
Sodium-to-potassium ratio	1.7 ± 0.7	1.7 (1.2, 2.1)	1.7 ± 0.7	1.6 (1.2, 2.03)	1.8 ± 0.7	1.7 (1.2, 2.2)	0.4
Salt (mg/d)	5667.9 ± 2077.7	5316.3 (4166.4, 6345.0)	5548.3 ± 1989.6	5224.7 (4063.9, 6254.7)	5797.0 ± 2177.5	5430.6 (4483.8, 6779.4)	0.5
Creatinine (mg/d)*	852.8 ± 352.7	842 (620, 1020)	853.4 ± 324.4	866 (609.5, 1010.3)	852.1 ± 329.7	789 (621, 1058)	0.9
Volume (ml/d)*	0.8 ± 0.4	0.8 (0.7, 1.1)	0.9 ± 0.4	0.8 (0.6, 1.01)	0.9 ± 0.3	0.8 (0.7, 1.0)	0.3

\*Variables are not normally distributed (Kolmogorov–Smirnov test). *p* values by one-way ANOVA for means or Mann–Whitney *U* test for medians.

TABLE 3: Distribution of sodium excretion according to age groups, sex, and nutritional status.

	Sample, <i>n</i> (%)	Sodium intake (mg/day) Median (interquartile range)	UL (mg/d)	Proportion over UL level, <i>n</i> (%)	<i>p</i> value
Age group					
6–8 y	45 (22.8)	1800.0 (1450.1, 2145.3)	1900	21 (46.7)	
9–13 y	71 (63.6)	2193.4 (1843.6, 2793.8)	2200	35 (49.3)	<0.001
14–18 y	15 (13.6)	2138.0 (1876.7, 2392.5)	2300	4 (26.7)	
Gender					
Boys	68 (41.2)	2057.0 (1600.0, 2462.5)	2300	40 (58.8)	
Girls	63 (58.8)	2138.0 (1765.3, 2669.1)	2300	35 (55.6)	0.487
Anthropometric status					
Thinness	13 (9.9)	1988.5 (1545.8, 2851.9)	2300	4 (26.5)	
Normal	91 (69.5)	2107.4 (1733.9, 2478.2)	2300	48 (52.5)	
Overweight	16 (12.2)	2044.3 (1787.5, 3153.6)	2300	10 (62.5)	0.678
Obese	11 (8.4)	2076.3 (1624.0, 2373.1)	2300	3 (31.5)	

UL: the upper limit refers to the highest daily level of sodium that is likely to pose no risk of adverse health effects to almost all individuals in the general population. The UL is not a recommended intake, and there is no apparent benefit to consuming levels of sodium above the adequate intake (AI). Source: Institute of Medicine [17]. *p* values are determined using the Kruskal–Wallis test for medians.

## 4. Discussion

Worldwide, there are limited data on sodium and potassium intakes in children and young people and are limited to high-income countries such as Europe and North America [32]. In Morocco, and to our best knowledge, this study is the first investigation to assess sodium and potassium status in children.

In this study, the assessment of sodium status in children aged 6–18 years old, with 24-hour urinary sodium excretion, clearly showed that the average daily sodium consumption in Morocco exceeds the recommended values for both boys and girls. With 24-hour urinary sodium excretion, the mean sodium concentration was  $2235.3 \pm 823.2$  mg/day (equivalent to  $5667.9 \pm 2077.7$  mg/day of salt).

Our results are in agreement with previously reported data worldwide using the 24 h urinary excretion approach, indicating a highly average consumption of sodium, particularly in industrialized countries. In Europe, the mean of sodium intake among children was included between 2400 mg/day and 3000 mg/day [33–36]. In USA and China, the sodium intake is much higher, and reported data showed

the mean sodium intake was 3100 mg/day and 3400 mg/day, respectively [37, 38]. Up to now, the highest mean of dietary sodium intake was reported in Chinese children aged 12–16 years from rural Shanxi, with a mean intake of 4 g/day [37].

In 2004, the Canadian Community Health Survey (CCHS) has evaluated sodium intake in a population of children aged 9–18 years and have clearly showed that sodium intakes are too high and exceed the ULs for 99% of boys and 91% of girls, increasing the risk to develop adverse health effects immediately or at adult age [36].

On the other hand, it is widely accepted that potassium has a critical role in many physiological processes, and potassium needs are widely discussed. In 2010, the American Dietary Guidelines identified potassium as a nutrient to be increased in the diet [39]. For instance, the Institute of Medicine (IOM) has reported that the adequate intake of potassium is 3800 mg/day for children aged 6–8 years and reaches 4700 mg/day for adolescents and adults [17].

In our study, potassium consumption is low for almost children and does not reach the recommended intake levels for all age groups. Overall, the average urinary potassium concentration was  $1431 \pm 636.5$  mg/day, and more than 97%

of children consume less than adequate intake. The average sodium-to-potassium ratio is  $1.7 \pm 0.7$ , so sodium is over-consumed compared to low potassium consumption.

Recently, Campanozzi et al. have reported similar results in Italy with an estimated daily potassium intake of 1530 mg/day for boys and 1400 mg/day for girls, much lower than the age-specific adequate intakes. Moreover, over 96% of the boys and 98% of the girls have a potassium intake lower than the recommended adequate intake [33]. Equivalent situation was reported in the USA. Indeed, the average of dietary potassium intake of the U.S. children aged 6–11 years and 12–19 years was 2225 mg/day and 2100 mg/day, respectively [40].

This study highlights that our recruited children have high level of sodium and low level of potassium intakes, as compared to recommended daily intakes. During last decades, Morocco has registered a nutritional transition, leading to changes in dietary habits and lifestyle modifications, especially at the biggest cities such as the Rabat region. Accordingly, the processed food and restaurant/fast food have been dramatically increased. In Morocco, food was usually prepared at home, but growing interest is currently given to a commercially processed food, especially by young people, under 25 years, representing 55% of the whole population [41].

A considerable part of the salt consumed in Morocco comes from the important bread consumption by adults. Indeed, most bakeries use between 15 and 20 g of salt/kg in bread preparation, and in Moroccan diet, bread is excessively consumed during meals, with an average of 500 g/day/person, which leads to a daily intake of 8 to 9 g of salt through bread alone [42]. For children, salt coming from bread would be slightly lower due to less consumption of bread. Worldwide, dietary sodium comes mainly from pre-prepared meals, snacks, and fast food. According to the UK National Food Survey data collected in 2000, cereal products (including bread, other baked goods, and breakfast cereals) accounted for the greatest proportion (38%) of household sodium intake, whereas 10% are added at home, during cooking or at table, and only 10% are naturally brought by food [43]. In the UK, children and adolescents, aged 4–18 years, surveyed for the UK Diet and Nutrition Investigation in 1992, showed that the major sources of sodium reflect those for adults (cereals contribute 38–40% and meat products contribute 20–24%) [44].

Fruits and vegetables are considered as the main sources of potassium. In Morocco, consumption of fruits and vegetables is very low and is more pronounced in children, consuming less than 374 g/person/day, giving rise to the lower potassium intake in Moroccan children [45].

This inadequate status of sodium and potassium in Morocco reflects the worldwide situation with an excess of sodium and deficiency of potassium intakes. According to this worrying situation, the WHO has developed and is implementing a global strategy and effective policies for salt consumption reduction, targeting the main sources of dietary sodium for all age groups. In this strategy, modification of dietary habits is also pointed out through nutritional education of the whole population, especially children,

including enhancing legume and fruit consumption, limiting fast food and processed products, avoiding the use of table salt, and so on [46].

Excessive sodium and deficient potassium intakes could lead to metabolic dysfunctions in children. Tracking of dietary habits from early childhood to adulthood has shown that children with extremely high levels of sodium intake tend to maintain those levels over time [12]. Thus, close monitoring of these children is needed for better management of their health.

Moreover, nutritional campaigns, aiming to increase the awareness about correct sodium and potassium intakes, should focus on children and adolescents as a major target in the framework of the population strategies for prevention of noncommunicable diseases.

The present study is very informative and has much strength mainly related to the technical approach used for assessment of sodium and potassium. In fact, evaluation of sodium and potassium was done by an objective indicator, 24 h urinary excretion, using a validated protocol for 24 h urine collection. However, the main limitation of the study is the assessment of sodium and potassium status in only one 24 h urine excretion per child. Moreover, the study was done only on children from Rabat and its region, and further studies on representative children from the whole country are necessary to draw consistent conclusions regarding the status of sodium and potassium in Moroccan children and to elaborate more adapted strategies to fight against associated NCDs.

## 5. Conclusion

In conclusion, the present study gives evidence that children from Rabat and its nearest region have inadequate sodium/potassium status, with higher sodium and lower potassium intakes compared to recommended values. These results give an idea about the sodium and potassium status in Moroccan children and strongly support the urgent need to implement a systematic strategy for the reduction of dietary sodium intake in Morocco. This study must be completed by the assessment of sodium and potassium intakes in Moroccan adults to have a global picture on these micronutrients' status and a valid baseline to follow up the population and evaluate the WHO strategy for reduction of salt consumption in Morocco.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Ethical Approval

The study protocol was approved by the Ethics Committee for Biomedical Research, Faculty of Medicine and Pharmacy of Rabat, Morocco.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

The authors would like to thank all investigators and all participants who have agreed to participate in the study, parents and children. They also thank Dr. Hicham El Berri, Head of Division of Non-Communicable Diseases, Direction of Epidemiology and Disease Control/Epidemiology & Control Diseases Direction, for taking part in this pilot study.

## References

- [1] WHO, *Global Status Report on Non-Communicable Diseases*, World Health Organization, Geneva, Switzerland, 2010, [http://whqlibdoc.who.int/publications/2011/9789240686458\\_eng.pdf](http://whqlibdoc.who.int/publications/2011/9789240686458_eng.pdf).
- [2] WHO, *Effect of Reduced Sodium Intake on Blood Pressure, Renal Function, Blood Lipids and Other Potential Adverse Effects*, World Health Organization, Geneva, Switzerland, 2012, ISBN 978 92 4 150491 1.
- [3] T. Meier, K. Senfleben, P. Deumelandt, O. Christen, K. Riedel, and M. Langer, "Healthcare costs associated with an adequate intake of sugars, salt and saturated fat in Germany: a health econometrical analysis," *PLoS One*, vol. 10, no. 9, Article ID e0135990, 2015.
- [4] D. Mozaffarian, S. Fahimi, G. M. Singh et al., "Global sodium consumption and death from cardiovascular causes," *New England Journal of Medicine*, vol. 371, no. 7, pp. 624–634, 2014.
- [5] R. Caudarella, F. Vescini, E. Rizzoli, and C. M. Francucci, "Salt intake, hypertension, and osteoporosis," *Journal of Endocrinological Investigation*, vol. 32, no. 4S, pp. 15–20, 2009.
- [6] WHO, *Prevention of Recurrent Heart Attacks and Strokes in Low and Middle Income Populations: Evidence-Based Recommendations for Policy Makers and Health Professionals*, World Health Organization, Geneva, Switzerland, 2003.
- [7] K. T. Khaw and E. Barrett-Connor, "The association between blood pressure, age, and dietary sodium and potassium: a population study," *Circulation*, vol. 77, no. 1, pp. 53–61, 1988.
- [8] J. M. Geleijnse, D. E. Grobbee, and A. Hofman, "Sodium and potassium intake and blood pressure change in childhood," *BMJ*, vol. 300, no. 6729, pp. 899–902, 1990.
- [9] D. S. Celermajer and J. G. J. Ayer, "Childhood risk factors for adult cardiovascular disease and primary prevention in childhood," *Heart*, vol. 92, no. 11, pp. 1701–1706, 2006.
- [10] C. Berkey, N. M. Laird, I. Valadian, and J. Gardner, "Modelling adolescent blood pressure patterns and their prediction of adult pressures," *Biometrics*, vol. 47, no. 3, pp. 1005–1018, 1991.
- [11] R. M. Lauer and W. R. Clarke, "Childhood risk factors for high adult blood pressure: the Muscatine study," *Pediatrics*, vol. 84, pp. 633–641, 1989.
- [12] E. Patterson, J. Wärnberg, J. Kearney, and M. Sjöström, "The tracking of dietary intakes of children and adolescents in Sweden over six years: the European Youth Heart Study," *International Journal of Behavioral Nutrition and Physical Activity*, vol. 6, p. 91, 2009.
- [13] M. R. Singer, L. L. Moore, E. J. Garrahe, and R. C. Ellison, "The tracking of nutrient intake in young children: the Framingham Children's Study," *American Journal of Public Health*, vol. 85, no. 12, pp. 1673–1677, 1995.
- [14] V. Mikkilä, L. Räsänen, O. T. Raitakari, P. Pietinen, and J. Viikari, "Longitudinal changes in diet from childhood into adulthood with respect to risk of cardiovascular diseases: the cardiovascular risk in young Finns study," *European Journal of Clinical Nutrition*, vol. 58, no. 7, pp. 1038–1045, 2004.
- [15] World Health Organization, *Guideline Potassium Intake for Adults and Children*, WHO, Geneva, Switzerland, 2012.
- [16] World Health Organization, *Guideline: Sodium Intake for Adults and Children*, WHO, Geneva, Switzerland, 2012.
- [17] Institute of Medicine (IOM), *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate, Panel on Dietary Reference Intakes for Electrolytes and Water, Standing Committee on the Scientific Evaluation of Dietary Reference Intakes*, Institute of Medicine, Kirtipur, Nepal, 2004, ISBN: 0-309-53049-0.
- [18] S. A. Bingham, C. Gill, A. Welch et al., "Comparison of dietary assessment methods in nutritional epidemiology," *British Journal of Nutrition*, vol. 72, no. 4, pp. 619–642, 1994.
- [19] D. G. Simons-Morton and E. Obarzanek, "Diet and blood pressure in children and adolescents," *Pediatric Nephrology*, vol. 11, no. 2, pp. 244–249, 1997.
- [20] J. Lambert, C. Agostoni, I. Elmadfa et al., "Dietary intake and nutritional status of children and adolescents in Europe," *British Journal of Nutrition*, vol. 92, no. S2, pp. S147–S211, 2004.
- [21] M. B. Livingstone, P. J. Robson, and J. M. Wallace, "Issues in dietary intake assessment of children and adolescents," *British Journal of Nutrition*, vol. 92, no. S2, pp. S213–S222, 2004.
- [22] Ministère de la Santé (MS), *National Strategy of Nutrition 2011-2019 Morocco*, Ministère de la Santé, Rabat, Morocco, 2010.
- [23] Ministry of Health, *National Survey of Iron Deficiency, the Use of Iodized Salt and Vitamin A Supplementation 2000 and the Inquiry. National Multi-Indicators and Youth Health 2006-2007*, Moroccan Ministry of Health, Rabat, Morocco, 2008.
- [24] Royaume du Maroc, "Rapport du Haut-Commissariat au Plan (HCP). Recensement général de la population et de l'habitat," 2004, <http://www.hcp.ma/file/111366>.
- [25] Mairie de Rabat, September 2013, <http://www.mairiederabat.com/economie.html>.
- [26] Wilaya Rabat Salé Zemmour Zaër, *Plan Communal de Développement de la Ville de Rabat*, Wilaya, Rabat, Morocco, 2012.
- [27] World Health Organization, "Physical status: the use and interpretation of anthropometry. Report of a WHO expert committee technical," Report Series No. 854, WHO, Geneva, Switzerland, 1995.
- [28] WHO, *AnthroPlus for Personal Computers Manual: Software for Assessing Growth of the World's Children and Adolescents*, WHO, Geneva, Switzerland, 2009, <http://www.who.int/growthref/tools/en/>.
- [29] C. A. Grimes, J. R. Baxter, K. J. Campbell et al., "Cross-sectional study of 24-hour urinary electrolyte excretion and associated health outcomes in a convenience sample of Australian primary schoolchildren: the salt and other nutrients in children (SONIC) study protocol," *JMIR Research Protocols*, vol. 4, no. 1, p. e7, 2015.
- [30] D. Hunter, "Biochemical indicators of dietary intake," in *Nutritional Epidemiology*, W. Willett, Ed., pp. 174–243, University Press, Oxford, UK, 1998.
- [31] J. R. Delanghe and M. M. Speckaert, "Creatinine determination according to Jaffe—what does it stand for?," *Clinical Kidney Journal*, vol. 4, no. 2, pp. 83–86, 2011.

- [32] WHO, *Sodium Intakes around the World/Paul Elliott and Ian Brown; Background Document Prepared for the Forum and Technical Meeting on Reducing Salt Intake in Populations, Paris*, WHO, Geneva, Switzerland, October 2006, ISBN 978 92 4 159593 5.
- [33] A. Campanozzi, S. Avallone, A. Barbato et al., “High sodium and low potassium intake among Italian children: relationship with age, body mass and blood pressure,” *PLoS One*, vol. 10, no. 4, Article ID e0121183, 2015.
- [34] N. M. Marrero, F. J. He, P. Whincup, and G. A. MacGregor, “Salt intake of children and adolescents in South London: consumption levels and dietary sources,” *Hypertension*, vol. 63, no. 5, pp. 1026–1032, 2014.
- [35] L. Shi, D. Krupp, and T. Remer, “Salt, fruit and vegetable consumption and blood pressure development: a longitudinal investigation in healthy children,” *British Journal of Nutrition*, vol. 111, no. 4, pp. 662–671, 2014.
- [36] A. Aparicio, E. Rodríguez, E. Cuadrado-Soto et al., “Estimation of salt intake assessed by urinary excretion of sodium over 24 h in Spanish subjects aged 7–11 years,” *European Journal of Nutrition*, vol. 56, p. 171, 2017.
- [37] F. J. He, Y. Wu, X. X. Feng et al., “School based education programme to reduce salt intake in children and their families (School-EduSalt): cluster randomised controlled trial,” *BMJ*, vol. 350, p. h770, 2015.
- [38] NHANES, *U.S. Department of Agriculture, Agricultural Research Service and U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, What We Eat In America, NHANES, 2005–2006*, NCHS, Hyattsville, MD, USA, 2006.
- [39] USDA and USHSS, *Dietary Guidelines for Americans, 2010*, U.S. Government Printing Office, Washington, DC, USA, 7th edition, 2010.
- [40] M. K. Hoy and J. D. Goldman, *Potassium Intake of the U.S. Population: What We Eat In America, NHANES 2009-2010*, Food Surveys Research Group Dietary Data Brief No. 10, Food Surveys Research Group, Beltsville, MD, USA, 2012.
- [41] Euromonitor, *Consumer Lifestyles in Morocco*, Euromonitor International Limited, London, UK, 2012.
- [42] N. Mokhtar, H. Belhadj, D. Kress, A. Zerrari, and N. Chaouki, “Food-105 fortification program in Morocco,” *Food and Nutrition Bulletin*, vol. 22, no. 4, pp. 427–430, 2001.
- [43] HMSO, *Scientific Advisory Committee on Nutrition, Salt and Health*, London, UK, 2003, <https://www.gov.uk/government/publications/sacn-salt-and-health-report>.
- [44] I. J. Brown, I. Tzoulaki, V. Candeias, and P. Elliott, “Salt intakes around the world: implications for public health,” *International Journal of Epidemiology*, vol. 38, no. 3, pp. 791–813, 2009.
- [45] HCP, *Le Maroc Face à la Transition Démographique. Les Cahiers du Plan No 1 Février-Mars*, Haut-Commissariat au Plan, Casablanca, Morocco, 2005.
- [46] Forum, “OMS sur la réduction des apports en sel au niveau des populations,” in *Réduire les Apports en sel au Niveau des Populations: Rapport du Forum et de la Réunion Technique OMS* Paris, France, October 2006, ISBN 978 92 4 259537 6.



**Hindawi**

Submit your manuscripts at  
[www.hindawi.com](http://www.hindawi.com)

