

Table S1: Primer sequences used in this study**a) RT-qPCR primers**

Genes	Forward primer (5'-3')	Reverse primer (5'-3')
PD-L1	TGGCATTGCTGAACGCATTT	TGCAGCCAGGTCTAATTGTTTT
TET-1	CATCAGTCAAGACTTTAAGCCCT	CGGGTGGTTTAGGTTCTGTTT
TET-2	GATAGAACCAACCATGTTGAGGG	TGGAGCTTTGTAGCCAGAGGT
TET-3	GCCGGTCAATGGTGCTAGAG	CGGTTGAAGGTTTCATAGAGCC
DNMT3a	CCGATGCTGGGGACAAGAAT	CCCGTCATCCACCAAGACAC
DNMT3b	AGGGAAGACTCGATCCTCGTC	GTGTGTAGCTTAGCAGACTGG
β -ACTIN	AGAGCTACGAGCTGCCTGAC	AGCACTGTGTTGGCGTACAG

b) PCR primers for CpG methylation analysis

Genes	Forward primer (5'-3')	Reverse primer (5'-3')
PD-L1 promoter	TTGTTTTGGGTAGAGGTGGG	AAAAACCAAATACATTACCTATTCTTA

c) Sequencing Primer

Genes	Forward primer (5'-3')	Reverse primer (5'-3')
M-13	TGTAAAACGACGGCCAGT	CAGGAAACAGCTATGACC

d) ChIP-qPCR primers

Genes	Forward primer (5'-3')	Reverse primer (5'-3')
PD-L1 promoter	TATTTATAAGGTGGAAGTTTTG AGG	TGCCCAAGGCAGCAAATCCAG