Research Article

Prognostic Values for the mRNA Expression of the ADAMTS Family of Genes in Gastric Cancer

Liang Liang,¹ Jin-hui Zhu,¹ Gang Chen¹,² Xin-gan Qin¹, and Jun-qiang Chen¹

¹Department of Gastrointestinal Surgery, The First Affiliated Hospital of Guangxi Medical University, Nanning, Guangxi Zhuang Autonomous Region 530021, China

²Department of Pathology, The First Affiliated Hospital of Guangxi Medical University, Nanning, Guangxi Zhuang Autonomous Region 530021, China

Correspondence should be addressed to Xin-gan Qin; qinxingan_gxmu@163.com and Jun-qiang Chen; gxhans@163.com

Received 19 December 2019; Accepted 28 February 2020; Published 20 August 2020

Academic Editor: Akira Hara

Copyright © 2020 Liang Liang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The “A Disintegrin and Metalloproteinase with Thrombospondin Motif” (ADAMTS) family of genes is involved in the occurrence and development of different cancers. However, the prognostic value of these genes in gastric cancer (GC) has not been revealed. The present study was thus conducted to determine the prognostic value for the ADAMTS family of genes in GC. First, we evaluated the mRNA expression levels of the ADAMTS family in GC patients using a GEPIA dataset. Thereafter, we determined the prognostic value of these genes by analyzing their mRNA level using the Kaplan–Meier Plotter database. The mRNA expression level of ADAMTS12 was randomly validated by qRT-PCR and meta-analysis while its coexpression genes were derived using Coexpedia. Finally, we performed Gene Ontology (GO) annotation and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analyses using the OmicShare Tools. Compared to normal tissues, expression of ADAMTS2 and 12 was significantly higher while that of ADAMTS1, 13, and 15 was significantly lower in GC tissues. According to the RNA-seq and gene chip data, the ADAMTS family (6, 7, 12, 15, and 18) of genes was closely related to the prognosis of GC, and their high expression levels were associated with poor prognosis and survival time. In addition, ADAMTS12 was highly expressed in 20 pairs of GC tissues based on RT-PCR ($P = 0.016$) and meta-analysis (SMD: 0.73, 95% CI: 0.32–1.14, $P < 0.001$). GO and KEGG pathway analyses indicated that the ADAMTS12 coexpressed genes were enriched in the pathways of extracellular matrix organization, extracellular matrix structural constituent, extracellular matrix, and protein digestion and absorption. Herein, we discovered the prognostic values and biological roles of the ADAMTS genes in GC.

1. Introduction

Gastric cancer (GC) is the most common lethal cancer worldwide. Although it is ranked fifth based on the number of patients diagnosed each year, it has the third highest mortality rate of all cancers [1]. Although the techniques used for early screening have improved, and there have been considerable improvements in comprehensive prevention and treatment measures owing to surgical treatment [2, 3], the prognosis of GC remains unsatisfactory [2]. Therefore, establishing reliable biomarkers to predict the prognosis of

GC could contribute to the development of individualized clinical treatment.

Carcinogenesis is associated with abnormalities in different cellular and intercellular mechanisms such as extracellular matrix (ECM) remodeling. Among all ECM proteases, “A Disintegrin and Metalloproteinase with Thrombospondins” (ADAMTSs) are a relatively new group and are considered to be associated with carcinogenesis and the local/distant spread of cancer [4].

ADAMTS is a complex extracellular protease related to carcinogenesis and tumor protection. ADAMTS can cleave

or interact with different ECM components or regulatory factors, ultimately affecting cell adhesion, migration, proliferation, and angiogenesis. Differences in the over-expression, mutation, or epigenetic silencing of the ADAMTS genes were identified in different sources of tumors, indicating the direct effect of these metalloproteinases on the development of cancer [5].

Presently, only few studies have been carried out on the ADAMTS family in GC. Jiang found that *ADAMTS2* may be a potential biomarker for determining the prognosis of GC. This researcher also demonstrated that *ADAMTS1*, 8, and 18 were highly expressed in GC and its lymph node metastasis [6]. Conversely, however, other studies have found that the hypermethylation of *ADAMTS8* and 18 is related to a decrease in the expression of GC and may play an important role in the invasion and metastasis of this malignancy [7]. *ADAMTS12* may also play a role in the tumor process owing to its proteolytic activity or serve as a potential molecule involved in the regulation of cell adhesion [8].

Because the role of the ADAMTS family of genes in tumor diseases remains unclear, especially in the study of prognosis, we aimed to determine the prognostic value of their expression in GC.

2. Materials and Methods

2.1. Detection of the Gene Expression of the ADAMTS Family. The expression level of the ADAMTS family of genes was detected using the GEPIA tool (<http://gepia.cancer-pku.cn>) [9]. Normal samples were selected from TCGA and GTEx databases and compared to those from the TCGA tumor group, using a standard processing pipeline. To explore the expression of the ADAMTS family of genes in the different stages of tumor development, major pathological stages or subpathological stages were selected for plotting. All expression data were first $\log_2(\text{TPM} + 1)$ transformed for plotting. One-way ANOVA was employed for differential analysis, and disease state (tumor or normal) was used as a variable to calculate the differential expression. The RNA-seq data for stomach cancer, which were used in the heatmap, were downloaded from UCSC Xena project (<http://xena.ucsc.edu>). Genomic alterations of the ADAMTS family were analyzed by cBioPortal, an integrative analytic platform of TCGA [10].

2.2. Prognostic Analysis of the ADAMTS Gene Family in GC Using the Kaplan–Meier (K–M) Plotter. The K–M plotter database was established using gene expression data and survival information from Gene Expression Omnibus (GEO) and TCGA (<http://kmplot.com/analysis>) [11]. Briefly, the gene chip data from the GEO database and mRNA-seq data from the TCGA database were selected. The chip data contains GSE14210, GSE22377, GSE51105, GSE15459, GSE29272, GSE62254, and other chips. The clinical parameters of tumor subtypes (tumor stage and grade) were recorded for patients, including gender, treatment, and HER2 status. The prognosis of ADAMTS in these

subtypes was then compared. Thereafter, 19 genes were separately searched in the K–M plotter and the resulting images were downloaded from the website.

2.3. Verification of ADAMTS12 Expression Using the Chip Data and RT-PCR. We searched the chip data to identify the GC-related chip datasets, including cancer and its control group, and extracted the expression of *ADAMTS12*. The STATA v12.0 statistical software was employed in the meta-analysis to derive the continuous variables. Standardized mean difference (SMD) and 95% confidence interval (CI) were calculated using the Random (I–V heterogeneity) test. A two-sided *P* value <0.05 was considered to indicate statistical significance.

Following the retrieval of written informed consent, a total of 20 paired GC and corresponding adjacent non-cancerous tissues were collected from patients who underwent gastric resection at the Department of Gastrointestinal Surgery between January 2019 and May 2019 at the First Affiliated Hospital of Guangxi Medical University. All experiments were approved by the Ethics Committee of the Hospital. Total RNA extraction was conducted with NucleoZOL (Macherey–Nagel, Germany), following the protocol of the manufacturer. Reverse transcription of the extracted RNA and real-time quantitative PCR (qRT-PCR) were successively performed according to the instructions provided in the RT-PCR kit (TaKaRa Biotechnology) and the fluorescence quantitative PCR kit (TaKaRa Biotechnology). The primer sequences for *ADAMTS12* and β -actin were produced by Sangon Biotech Company (Shanghai, China). The following primer sequences were used: *ADAMTS12*, 5'-AACGCTATCG-CTTGTGCAAC-3' and 5'-CTCACAAGGATGTGCTGG-GT-3'; and β -actin, 5'-TGGCAGCCAGCACAATGAA-3' and 5'-CTAAGTCATAGTCCGCCTAGAAGCA-3'. The relative expression levels of *ADAMTS12* were quantified using the $2^{-\Delta\Delta CT}$ method.

2.4. Gene Function Enrichment Analysis. Coexpedia is a database that contains context-associated coexpression networks inferred from individual series of microarray samples for human and mouse from chip data. Hence, this database (<http://www.coexpedia.org>) was employed to analyze the *ADAMTS12*-related genes [12]. In addition, we analyzed the *ADAMTS12* coexpressed genes using RNA-seq data to determine the possible mechanisms of action of *ADAMTS12*. Kyoto Encyclopedia of Genes and Genomes (KEGG) and Gene Ontology (GO) analyses were performed using the OmicShare tools, a free online platform for data analysis (<http://www.omicshare.com/tools>).

3. Results

3.1. mRNA Expression Level of the ADAMTS Family in GC. Based on our search of the RNA-seq data, the expression level of *ADAMTS2*, *ADAMTS9*, *ADAMTS12*, *ADAMTS14*, and other genes in the GC group was higher than that found in the normal group (Figure 1) ($P < 0.05$). However, the

FIGURE 1: The expression of the ADAMTS family based on sequencing data. The mRNA expression of the ADAMTS family between tumor and normal; red: tumor; gray: normal; * $P < 0.01$. (a) ADAMTS1, (b) ADAMTS2, (c) ADAMTS3, (d) ADAMTS4, (e) ADAMTS5, (f) ADAMTS6, (g) ADAMTS7, (h) ADAMTS8, (i) ADAMTS9, (j) ADAMTS10, (k) ADAMTS12, (l) ADAMTS13, (m) ADAMTS14, (n) ADAMTS15, (o) ADAMTS16, (p) ADAMTS17, (q) ADAMTS18, (r) ADAMTS19, and (s) ADAMTS20.

FIGURE 2: Heatmap depicting the expression of the ADAMTS family in GC patients. Red represents high expression; blue represents low expression; black represents not detected.

FIGURE 3: Expression of the ADAMTS family at different clinical stages. One-way ANOVA was performed to calculate the expression of the ADAMTS family in the differential pathological stage. (a) ADAMTS1, (b) ADAMTS2, (c) ADAMTS3, (d) ADAMTS4, (e) ADAMTS5, (f) ADAMTS6, (g) ADAMTS7, (h) ADAMTS8, (i) ADAMTS9, (j) ADAMTS10, (k) ADAMTS12, (l) ADAMTS13, (m) ADAMTS14, (n) ADAMTS15, (o) ADAMTS16, (p) ADAMTS17, (q) ADAMTS18, (r) ADAMTS19, and (s) ADAMTS20.

FIGURE 4: The genomic alterations of the ADAMTS family in RNA-seq data. *ADAMTS12* was found to have the highest alteration (12%), with amplification detected to be the main cause of alterations.

expression level of *ADAMTS1* was significantly lower in the GC tissues than in normal tissues ($P < 0.05$) (Figure 1). The heatmap presented in Figure 2 shows the expression of each ADAMTS family of molecules in tumor tissues and the adjacent tissues. Figure 3 depicts the relationship between the ADAMTS molecules and tumor progression. Genes such as *ADAMTS1*, *ADAMTS10*, and *ADAMTS12* were found to increase with tumor progression ($P < 0.05$). By analyzing the ADAMTS family of genes, we found that *ADAMTS12* and *ADAMTS16* had the highest mutation rate (i.e., 12%), which was mainly due to amplification and missense mutation. The lowest rate of genetic change occurred in *ADAMTS6* (rate = 1.8%; Figure 4).

3.2. Prognostic Values of the ADAMTS Family in Patients with GC Based on RNA-Seq and Chip Data. Using the RNA-seq data, we determined the prognostic value for the mRNA expression of the ADAMTS family in GC. *ADAMTS4* (HR = 1.39, $P < 0.001$), *ADAMTS6* (HR = 1.75, $P < 0.001$), *ADAMTS7* (HR = 1.43, $P < 0.033$), *ADAMTS10* (HR = 1.45, $P < 0.026$), *ADAMTS12* (HR = 1.41, $P < 0.040$), *ADAMTS15* (HR = 1.49, $P < 0.016$), *ADAMTS16* (HR = 1.42, $P < 0.034$), *ADAMTS18* (HR = 1.59, $P = 0.005$), and other molecules were found to be associated with the overall survival (OS) of

patients with GC. In fact, the survival time of patients with a high expression of these molecules was significantly shorter than that of patients with a low expression of these molecules (Figure 5).

The expression levels of *ADAMTS10* (HR = 2.22, $P = 0.020$), *ADAMTS18* (HR = 2.02, $P = 0.036$), and *ADAMTS20* (HR = 2.47, $P = 0.006$) were recognized to be associated with the recurrence of GC. The survival time of patients with a high expression level of these genes was significantly lower than that of patients with a low expression level of these genes (Figure 6).

Finally, we verified the prognostic value of 19 genes of the ADAMTS family in the chip data. As a result, *ADAMTS1* (HR = 1.68, $P < 0.001$), *ADAMTS2* (HR = 1.49, $P < 0.001$), *ADAMTS3* (HR = 1.33, $P < 0.001$), *ADAMTS5* (HR = 1.4, $P < 0.002$), *ADAMTS6* (HR = 1.7, $P < 0.001$), *ADAMTS7* (HR = 1.63, $P < 0.001$), *ADAMTS8* (HR = 1.72, $P < 0.001$), *ADAMTS9* (HR = 1.46, $P < 0.001$), *ADAMTS12* (HR = 1.39, $P < 0.001$), *ADAMTS14* (HR = 1.58, $P < 0.001$), *ADAMTS15* (HR = 1.29, $P < 0.001$), *ADAMTS18* (HR = 1.53, $P < 0.001$), and *ADAMTS20* (HR = 1.67) were found to be closely related to OS in patients with GC ($P < 0.001$). The total survival time of patients with a high expression level of these molecules was significantly shorter than that of patients with low expression (Figure 7).

FIGURE 5: Continued.

FIGURE 5: The overall survival value for the expression of the ADAMTS family based on RNA-seq data. (a) ADAMTS1, (b) ADAMTS2, (c) ADAMTS3, (d) ADAMTS4, (e) ADAMTS5, (f) ADAMTS6, (g) ADAMTS7, (h) ADAMTS8, (i) ADAMTS9, (j) ADAMTS10, (k) ADAMTS12, (l) ADAMTS13, (m) ADAMTS14, (n) ADAMTS15, (o) ADAMTS16, (p) ADAMTS17, (q) ADAMTS18, (r) ADAMTS19, and (s) ADAMTS20.

FIGURE 6: Continued.

FIGURE 6: The relapse-free survival value for the expression of the ADAMTS family based on RNA-seq data. (a) ADAMTS1, (b) ADAMTS2, (c) ADAMTS3, (d) ADAMTS4, (e) ADAMTS5, (f) ADAMTS6, (g) ADAMTS7, (h) ADAMTS8, (i) ADAMTS9, (j) ADAMTS10, (k) ADAMTS12, (l) ADAMTS13, (m) ADAMTS14, (n) ADAMTS15, (o) ADAMTS16, (p) ADAMTS17, (q) ADAMTS18, (r) ADAMTS19, and (s) ADAMTS20.

FIGURE 7: Continued.

FIGURE 7: The overall survival value for the expression of the ADAMTS family based on chip data. (a) 222486_s_at ADAMTS1, (b) 226311_at ADAMTS2, (c) 214913_at ADAMTS3, (d) 235368_at ADAMTS5, (e) 237411_at ADAMTS6, (f) 228911_at ADAMTS7, (g) 235649_at ADAMTS8, (h) 1554697_at ADAMTS9, (i) 232133_at ADAMTS10, (j) 221421_s_at ADAMTS12, (k) 223844_at ADAMTS13, (l) 230167_at ADAMTS14, (m) 1553427_at ADAMTS15, (n) 238125_at ADAMTS16, (o) 1552726_at ADAMTS17, (p) 230040_at ADAMTS18, (q) 1553179_at ADAMTS19, and (r) 220717 ADAMTS20.

For postprogression survival (PPS) (Figure 8), we found that 14 genes, namely, *ADAMTS1* (HR = 2.14, $P < 0.001$), *ADAMTS2* (HR = 1.66, $P < 0.001$), *ADAMTS3* (HR = 1.27, $P < 0.001$), *ADAMTS5* (HR = 1.89, $P < 0.001$), *ADAMTS6* (HR = 1.97, $P < 0.001$), *ADAMTS7* (HR = 1.98, $P < 0.001$), *ADAMTS8* (HR = 1.7, $P < 0.001$), *ADAMTS9* (HR = 1.42, $P < 0.001$), *ADAMTS12* (HR = 1.59, $P < 0.001$), *ADAMTS14* (HR = 1.94, $P < 0.001$), *ADAMTS15* (HR = 1.43, $P < 0.001$), *ADAMTS17* (HR = 1.44, $P < 0.001$), *ADAMTS18* (HR = 1.67, $P < 0.001$), and *ADAMTS20* (HR = 1.97, $P < 0.001$), were closely related to PPS. The total survival time of patients with a high expression level

of these genes was significantly shorter than that of patients with a low expression level.

Through subgroup analysis, genes such as *ADAMTS15*, 16, and 18 were identified to be associated with unfavorable OS in stage II; *ADAMTS2*, 5, 6, 7, and 20 were associated with unfavorable OS in stage III; and *ADAMTS1*, 5, 6, 7, and 14 were associated with unfavorable OS in stage IV (Table 1). The expression of *ADAMTS1*, 5, 6, 7, 12, 16, 18, and 20 was associated with poor OS and did not significantly differ based on gender. Although the expression levels of *ADAMTS2*, 4, 14, and 15 were related to poor OS among men, only the expression level of *ADAMTS9* was associated

FIGURE 8: Continued.

FIGURE 8: The postprogression survival value for the expression of the ADAMTS family based on chip data. (a) 222486_s_at ADAMTS1, (b) 226311_at ADAMTS2, (c) 214913_at ADAMTS3, (d) 235368_at ADAMTS5, (e) 237411_at ADAMTS6, (f) 228911_at ADAMTS7, (g) 235649_at ADAMTS8, (h) 1554697_at ADAMTS9, (i) 232133_at ADAMTS10, (j) 221421_s_at ADAMTS12, (k) 223844_at ADAMTS13, (l) 230167_at ADAMTS14, (m) 1553427_at ADAMTS15, (n) 238125_at ADAMTS16, (o) 1552726_at ADAMTS17, (p) 230040_at ADAMTS18, (q) 1553179_at ADAMTS19, and (r) 220717 ADAMTS20.

TABLE 1: Correlation between genes expression of ADAMTS family and OS in gastric cancer patients of clinical stage.

ADAMTSs gene chip	Clinical stages I (n = 69); II (n = 145); III (n = 319); IV (n = 152)	HR (95% CI)	P value	ADAMTSs RNA-seq	Clinical stages I (n = 50); II (n = 111); III (n = 149); IV (n = 38)	HR (95% CI)	P value
ADAMTS1	I	1.95 (0.60–6.40)	0.26	ADAMTS1	I	1.08 (0.32–3.64)	0.90
	II	1.57 (0.83–2.97)	0.16		II	2.00 (0.98–4.05)	0.05
	III	1.11 (0.77–1.62)	0.57		III	1.12 (0.69–1.80)	0.65
	IV	1.52 (1.02–2.27)	0.04		IV	1.12 (0.49–2.56)	0.78
ADAMTS2	I	0.26 (0.07–0.96)	0.03	ADAMTS2	I	2.11 (0.61–7.25)	0.22
	II	0.62 (0.33–1.19)	0.15		II	1.62 (0.80–3.26)	0.17
	III	1.56 (1.07–2.28)	0.02		III	1.36 (0.84–2.19)	0.21
	IV	1.29 (0.87–1.91)	0.20		IV	1.28 (0.55–2.96)	0.57
ADAMTS3	I	1.18 (0.43–3.23)	0.75	ADAMTS3	I	0.60 (0.18–2.07)	0.42
	II	1.52 (0.83–2.76)	0.17		II	1.26 (0.63–2.51)	0.51
	III	1.13 (0.85–1.5)	0.40		III	1.35 (0.83–2.18)	0.22
	IV	1.46 (1.00–2.15)	0.05		IV	1.12 (0.48–2.58)	0.80
ADAMTS4(none)	—			ADAMTS4	I	1.42 (0.41–4.91)	0.57
					II	2.04 (0.99–4.19)	0.05
					III	1.24 (0.76–2.01)	0.39
					IV	1.20 (0.52–2.78)	0.67

TABLE 1: Continued.

ADAMTSs gene chip	Clinical stages I (n = 69); II (n = 145); III (n = 319); IV (n = 152)	HR (95% CI)	P value	ADAMTSs RNA-seq	Clinical stages I (n = 50); II (n = 111); III (n = 149); IV (n = 38)	HR (95% CI)	P value
ADAMTS5	I	0.48 (0.16–1.49)	0.19	ADAMTS5	I	0.69 (0.22–2.16)	0.52
	II	1.66 (0.86–3.18)	0.13		II	1.20 (0.61–2.36)	0.61
	III	1.81 (1.24–2.64)	<0.01		III	1.44 (0.89–2.33)	0.14
	IV	1.51 (1.02–2.25)	0.04		IV	1.32 (0.57–3.08)	0.51
ADAMTS6	I	0.41 (0.12–1.33)	0.12	ADAMTS6	I	1.15 (0.36–3.65)	0.81
	II	1.40 (0.74–2.63)	0.29		II	1.50 (0.75–3.00)	0.25
	III	1.36 (0.93–1.98)	0.11		III	1.68 (1.04–2.72)	0.03
	IV	1.31 (0.88–1.95)	0.18		IV	3.37 (1.30–8.69)	<0.01
ADAMTS7	I	2.56 (0.77–8.56)	0.11	ADAMTS7	I	2.82 (0.75–10.63)	0.11
	II	1.35 (0.71–2.57)	0.35		II	1.10 (0.55–2.17)	0.79
	III	1.77 (1.21–2.57)	<0.01		III	1.44 (0.89–2.33)	0.14
	IV	1.76 (1.18–2.63)	<0.01		IV	1.19 (0.51–2.76)	0.68
ADAMTS8	I	1.87 (0.56–6.23)	0.30	ADAMTS8	I	1.89 (0.57–6.32)	0.29
	II	1.52 (0.80–2.88)	0.19		II	1.16 (0.59–2.28)	0.66
	III	1.2 (0.82–1.74)	0.34		III	1.55 (0.96–2.52)	0.07
	IV	1.4 (0.94–2.09)	0.10		IV	1.13 (0.49–2.59)	0.77
ADAMTS9	I	0.86 (0.29–2.57)	0.78	ADAMTS9	I	1.11 (0.34–3.65)	0.87
	II	1.39 (0.74–2.61)	0.31		II	1.08 (0.55–2.13)	0.82
	III	1.36 (0.94–1.98)	0.10		III	1.22 (0.75–1.99)	0.41
	IV	1.58 (1.06–2.34)	0.02		IV	1.35 (0.58–3.12)	0.49
ADAMTS10	I	2.12 (0.69–6.48)	0.18	ADAMTS10	I	1.10 (0.33–3.62)	0.88
	II	0.89 (0.48–1.65)	0.71		II	1.72 (0.86–3.43)	0.12
	III	1.15 (0.79–1.67)	0.47		III	1.25 (0.77–2.02)	0.37
	IV	0.76 (0.51–1.13)	0.17		IV	1.46 (0.63–3.39)	0.38
ADAMTS12	I	1.06 (0.40–2.84)	0.91	ADAMTS12	I	1.41 (0.45–4.44)	0.56
	II	1.67 (0.92–3.05)	0.09		II	1.72 (0.85–3.46)	0.12
	III	1.33 (1.00–1.77)	0.05		III	1.43 (0.89–2.32)	0.14
	IV	1.13 (0.77–1.65)	0.54		IV	1.57 (0.67–3.66)	0.29

TABLE 1: Continued.

ADAMTSs gene chip	Clinical stages I (n = 69); II (n = 145); III (n = 319); IV (n = 152)	HR (95% CI)	P value	ADAMTSs RNA-seq	Clinical stages I (n = 50); II (n = 111); III (n = 149); IV (n = 38)	HR (95% CI)	P value
ADAMTS13	I	1.75 (0.57–5.37)	0.32	ADAMTS13	I	0.39 (0.12–1.31)	0.11
	II	1.33 (0.71–2.49)	0.37		II	0.99 (0.50–1.94)	0.97
	III	1.11 (0.77–1.62)	0.57		III	0.91 (0.56–1.46)	0.69
	IV	0.86 (0.58–1.28)	0.47		IV	0.88 (0.39–2.01)	0.76
ADAMTS14	I	0.83 (0.26–2.67)	0.75	ADAMTS14	I	0.66 (0.21–2.09)	0.47
	II	1.19 (0.62–2.25)	0.60		II	0.76 (0.38–1.49)	0.42
	III	1.42 (0.98–2.07)	0.06		III	1.36 (0.84–2.20)	0.21
	IV	2.10 (1.40–3.16)	<0.01		IV	0.96 (0.42–2.23)	0.93
ADAMTS15	I	1.13 (0.38–3.39)	0.82	ADAMTS15	I	0.69 (0.21–2.26)	0.54
	II	1.33 (0.71–2.49)	0.36		II	2.14 (1.05–4.35)	0.03
	III	1.10 (0.76–1.60)	0.62		III	1.32 (0.82–2.13)	0.26
	IV	1.31 (0.89–1.95)	0.17		IV	1.78 (0.76–4.17)	0.18
ADAMTS16	I	1.25 (0.42–3.73)	0.69	ADAMTS16	I	1.10 (0.35–3.50)	0.87
	II	1.00 (0.53–1.87)	0.99		II	2.60 (1.26–5.36)	<0.01
	III	1.06 (0.73–1.54)	0.76		III	1.13 (0.70–1.81)	0.62
	IV	1.20 (0.81–1.77)	0.37		IV	1.66 (0.71–3.88)	0.24
ADAMTS17	I	0.89 (0.29–2.70)	0.84	ADAMTS17	I	1.22 (0.38–3.87)	0.74
	II	1.88 (0.99–3.58)	0.05		II	0.53 (0.26–1.08)	0.08
	III	1.17 (0.81–1.70)	0.40		III	0.71 (0.44–1.14)	0.16
	IV	0.97 (0.65–1.43)	0.86		IV	1.58 (0.69–3.64)	0.27
ADAMTS18	I	1.32 (0.42–4.16)	0.64	ADAMTS18	I	2.51 (0.75–8.42)	0.12
	II	2.35 (1.20–4.59)	0.01		II	2.41 (1.18–4.82)	0.01
	III	1.06 (0.73–1.55)	0.74		III	1.49 (0.92–2.41)	0.11
	IV	1.22 (0.82–1.81)	0.32		IV	1.00 (0.43–2.33)	1.00
ADAMTS19	I	3.17 (0.97–10.37)	0.05	ADAMTS19	I	4.89 (1.32–18.14)	<0.01
	II	1.07 (0.57–2.01)	0.82		II	0.52 (0.26–1.06)	0.07
	III	1.06 (0.73–1.54)	0.74		III	1.30 (0.80–2.09)	0.29
	IV	0.97 (0.65–1.44)	0.88		IV	1.52 (0.65–3.52)	0.33

TABLE 1: Continued.

ADAMTSs gene chip	Clinical stages I (n = 69); II (n = 145); III (n = 319); IV (n = 152)	HR (95% CI)	P value	ADAMTSs RNA-seq	Clinical stages I (n = 50); II (n = 111); III (n = 149); IV (n = 38)	HR (95% CI)	P value
ADAMTS20	I	1.56 (0.55–4.37)	0.40	ADAMTS20	I	1.58 (0.48–5.21)	0.45
	II	1.36 (0.75–2.46)	0.31		II	1.44 (0.73–2.85)	0.29
	III	1.42 (1.07–1.89)	0.02		III	0.87 (0.54–1.41)	0.58
	IV	1.25 (0.85–1.82)	0.26		IV	1.04 (0.44–2.45)	0.92

TABLE 2: Correlation between genes expression of ADAMTS family and OS in gastric cancer patients of differentiation.

ADAMTSs gene chip	Differentiation poorly (n = 166); moderately (n = 67); well (n = 32)	HR (95% CI)	P value	ADAMTSs RNA-seq	Differentiation I (n = 12); II (n = 134); III (n = 218)	HR (95% CI)	P value
ADAMTS1	Poorly differentiated	0.85 (0.57–1.27)	0.43	ADAMTS1	Grade I	1.29 × 10 ⁹ (0–inf)	0.23
	Moderately differentiated	0.96 (0.50–1.83)	0.90		Grade II	1.26 (0.72–2.20)	0.43
	Well differentiated	—			Grade III	1.37 (0.91–2.07)	0.13
ADAMTS2	Poorly differentiated	1.17 (0.78–1.74)	0.45	ADAMTS2	Grade I	1.29 × 10 ⁹ (0–inf)	0.23
	Moderately differentiated	1.31 (0.69–2.52)	0.41		Grade II	1.35 (0.76–2.37)	0.30
	Well differentiated	—			Grade III	1.30 (0.86–1.97)	0.21
ADAMTS3	Poorly differentiated	1.23 (0.82–1.83)	0.32	ADAMTS3	Grade I	1.29 × 10 ⁹ (0–inf)	0.23
	Moderately differentiated	1.35 (0.71–2.59)	0.36		Grade II	0.79 (0.45–1.40)	0.43
	Well differentiated	1.52 (0.64–3.61)	0.34		Grade III	1.33 (0.88–2.01)	0.17
ADAMTS5	Poorly differentiated	1.16 (0.72–1.88)	0.54	ADAMTS5	Grade I	1.29 × 10 ⁹ (0–inf)	0.23
	Moderately differentiated	0.96 (0.50–1.84)	0.91		Grade II	0.98 (0.56–1.73)	0.95
	Well differentiated	—			Grade III	1.26 (0.83–1.91)	0.27
ADAMTS6	Poorly differentiated	1.07 (0.66–1.73)	0.78	ADAMTS6	Grade I	3.9 × 10 ⁹ (0–inf)	0.44
	Moderately differentiated	1.49 (0.78–2.84)	0.23		Grade II	1.45 (0.81–2.58)	0.20
	Well differentiated	—			Grade III	1.93 (1.27–2.93)	<0.01
ADAMTS7	Poorly differentiated	0.97 (0.65–1.44)	0.86	ADAMTS7	Grade I	1.41 (0.08–23.57)	0.81
	Moderately differentiated	2.39 (1.24–4.59)	<0.01		Grade II	1.12 (0.63–1.99)	0.69
	Well differentiated	—			Grade III	1.58 (1.03–2.41)	0.04
ADAMTS8	Poorly differentiated	1.10 (0.68–1.79)	0.69	ADAMTS8	Grade I	3.56 × 10 ⁹ (0–inf)	0.06
	Moderately differentiated	1.43 (0.75–2.74)	0.28		Grade II	1.00 (0.57–1.75)	0.99
	Well differentiated	—			Grade III	1.07 (0.71–1.61)	0.76

TABLE 2: Continued.

ADAMTSs gene chip	Differentiation poorly (n = 166); moderately (n = 67); well (n = 32)	HR (95% CI)	P value	ADAMTSs RNA-seq	Differentiation I (n = 12); II (n = 134); III (n = 218)	HR (95% CI)	P value
ADAMTS9	Poorly differentiated	1.07 (0.66–1.74)	0.77	ADAMTS9	Grade I	1.29 × 109 (0- inf)	0.23
	Moderately differentiated	1.34 (0.70–2.55)	0.38		Grade II	0.85 (0.48–1.49)	0.57
	Well differentiated	—			Grade III	1.50 (0.98–2.29)	0.06
ADAMTS10	Poorly differentiated	1.08 (0.67–1.74)	0.76	ADAMTS10	Grade I	1.29 × 109 (0- inf)	0.23
	Moderately differentiated	0.81 (0.42–1.55)	0.52		Grade II	1.31 (0.75–2.30)	0.35
	Well differentiated	—			Grade III	1.41 (0.93–2.14)	0.10
ADAMTS12	Poorly differentiated	1.30 (0.87–1.94)	0.21	ADAMTS12	Grade I	0.71 (0.04–11.79)	0.81
	Moderately differentiated	1.00 (0.52–1.92)	0.99		Grade II	1.50 (0.85–2.66)	0.16
	Well differentiated	0.95 (0.40–2.23)	0.90		Grade III	1.54 (1.02–2.34)	0.04
ADAMTS13	Poorly differentiated	0.87 (0.54–1.41)	0.58	ADAMTS13	Grade I	1.62 × 109 (0- inf)	0.32
	Moderately differentiated	1.06 (0.55–2.04)	0.87		Grade II	0.46 (0.25–0.85)	0.01
	Well differentiated	—			Grade III	0.88 (0.59–1.33)	0.56
ADAMTS14	Poorly differentiated	1.28 (0.78–2.08)	0.32	ADAMTS14	Grade I	1.29 × 109 (0- inf)	0.23
	Moderately differentiated	1.80 (0.93–3.47)	0.08		Grade II	0.81 (0.46–1.44)	0.48
	Well differentiated	—			Grade III	0.99 (0.66–1.49)	0.95
ADAMTS15	Poorly differentiated	0.97 (0.59–1.59)	0.91	ADAMTS15	Grade I	2.60 × 109 (0- inf)	0.09
	Moderately differentiated	1.34 (0.69–2.59)	0.38		Grade II	1.23 (0.70–2.17)	0.47
	Well differentiated	—			Grade III	1.59 (1.05–2.41)	0.03
ADAMTS16	Poorly differentiated	0.69 (0.42–1.14)	0.14	ADAMTS16	Grade I	0.71 (0.04–11.79)	0.81
	Moderately differentiated	1.43 (0.75–2.73)	0.28		Grade II	0.95 (0.54–1.68)	0.86
	Well differentiated	—			Grade III	1.56 (1.03–2.36)	0.03
ADAMTS17	Poorly differentiated	0.95 (0.59–1.53)	0.83	ADAMTS17	Grade I	1.41 (0.08–23.57)	0.81
	Moderately differentiated	1.14 (0.59–2.18)	0.70		Grade II	0.98 (0.55–1.74)	0.93
	Well differentiated	—			Grade III	0.77 (0.51–1.17)	0.22
ADAMTS18	Poorly differentiated	1.18 (0.73–1.91)	0.50	ADAMTS18	Grade I	1.29 × 109 (0- inf)	0.23
	Moderately differentiated	1.27 (0.66–2.43)	0.47		Grade II	1.27 (0.72–2.24)	0.41
	Well differentiated	—			Grade III	1.84 (1.20–2.83)	<0.01

TABLE 2: Continued.

ADAMTSs gene chip	Differentiation poorly (n = 166); moderately (n = 67); well (n = 32)	HR (95% CI)	P value	ADAMTSs RNA-seq	Differentiation I (n = 12); II (n = 134); III (n = 218)	HR (95% CI)	P value
ADAMTS19	Poorly differentiated	0.79 (0.49–1.28)	0.34	ADAMTS19	Grade I	1.86 × 109 (0–inf)	0.16
	Moderately differentiated	1.25 (0.65–2.38)	0.50		Grade II	1.12 (0.63–1.98)	0.71
	Well differentiated	—			Grade III	1.20 (0.79–1.81)	0.39
ADAMTS20	Poorly differentiated	1.10 (0.74–1.64)	0.63	ADAMTS20	Grade I	0 (0–inf)	0.09
	Moderately differentiated	0.95 (0.5–1.82)	0.88		Grade II	1.21 (0.68–2.15)	0.52
	Well differentiated	1.92 (0.81–4.57)	0.13		Grade III	1.20 (0.80–1.81)	0.38

TABLE 3: Correlation between genes expression of ADAMTS family and OS in gastric cancer patients of different genders.

ADAMTSs gene chip	Gender female (n = 244); male (n = 567)	HR (95% CI)	P value	ADAMTSs RNA-seq	Gender female (n = 133); male (n = 238)	HR (95% CI)	P value
ADAMTS1	Female	1.63 (1.05–2.50)	0.03	ADAMTS1	Female	1.37 (0.76–2.47)	0.29
	Male	1.63 (1.21–2.20)	<0.01		Male	1.39 (0.94–2.06)	0.10
ADAMTS2	Female	1.27 (0.83–1.95)	0.27	ADAMTS2	Female	1.59 (0.87–2.89)	0.13
	Male	1.68 (1.24–2.26)	<0.01		Male	1.16 (0.79–1.72)	0.45
ADAMTS3	Female	1.56 (1.09–2.22)	0.01	ADAMTS3	Female	1.27 (0.86–1.88)	0.23
	Male	1.24 (1.00–1.53)	0.05		Male	1.65 (0.90–3.02)	0.10
ADAMTS4 (Not find)	—	—		ADAMTS4	Female Male	0.86(0.48–1.54) 1.62(1.09–2.43)	0.60 0.02
ADAMTS5	Female	1.75 (1.13–2.70)	0.01	ADAMTS5	Female	1.65 (0.91–2.98)	0.10
	Male	1.47 (1.09–1.98)	0.01		Male	1.01 (0.68–1.49)	0.97
ADAMTS6	Female	1.57 (1.02–2.42)	0.04	ADAMTS6	Female	2.36 (1.28–4.35)	<0.01
	Male	1.76 (1.31–2.38)	<0.01		Male	1.47 (0.99–2.18)	0.05
ADAMTS7	Female	1.64 (1.07–2.52)	0.02	ADAMTS7	Female	1.35 (0.74–2.45)	0.32
	Male	1.88 (1.40–2.53)	<0.01		Male	1.48 (0.99–2.21)	0.05
ADAMTS8	Female	1.98 (1.28–3.08)	<0.01	ADAMTS8	Female	1.04 (0.58–1.86)	0.91
	Male	1.31 (0.97–1.76)	0.07		Male	1.16 (0.79–1.72)	0.45
ADAMTS9	Female	1.58 (1.03–2.43)	0.03	ADAMTS9	Female	1.72 (0.94–3.14)	0.08
	Male	1.27 (0.95–1.70)	0.11		Male	1.43 (0.97–2.13)	0.07
ADAMTS10	Female	1.08 (0.70–1.65)	0.73	ADAMTS10	Female	1.66 (0.91–3.01)	0.09
	Male	0.94 (0.70–1.26)	0.67		Male	1.26 (0.85–1.86)	0.25

TABLE 3: Continued.

ADAMTSs gene chip	Gender female (n = 244); male (n = 567)	HR (95% CI)	P value	ADAMTSs RNA-seq	Gender female (n = 133); male (n = 238)	HR (95% CI)	P value
ADAMTS12	Female	1.67 (1.17–2.37)	<0.01	ADAMTS12	Female	2.02 (1.10–3.70)	0.02
Male	1.30 (1.05–1.61)	0.02	Male	1.63 (1.00–2.66)	<0.05		
ADAMTS13	Female	0.94 (0.62–1.45)	0.79	ADAMTS13	Female	0.63 (0.35–1.15)	0.13
	Male	1.15 (0.86–1.54)	0.34		Male	0.87 (0.59–1.29)	0.49
ADAMTS14	Female	1.23 (0.80–1.89)	0.35	ADAMTS14	Female	1.09 (0.61–1.97)	0.76
	Male	1.80 (1.33–2.43)	<0.01		Male	1.00 (0.67–1.47)	0.98
ADAMTS15	Female	1.22 (0.79–1.87)	0.37	ADAMTS15	Female	0.86 (0.48–1.55)	0.61
	Male	1.23 (0.92–1.65)	0.16		Male	1.87 (1.25–2.79)	<0.01
ADAMTS16	Female	1.34 (0.87–2.05)	0.18	ADAMTS16	Female	1.83 (1.01–3.32)	0.04
	Male	1.08 (0.80–1.45)	0.61		Male	1.87 (1.25–2.79)	<0.01
ADAMTS17	Female	1.31 (0.85–2.01)	0.22	ADAMTS17	Female	0.78 (0.43–1.41)	0.41
	Male	1.12 (0.83–1.50)	0.45		Male	0.87 (0.59–1.29)	0.49
ADAMTS18	Female	1.32 (0.86–2.02)	0.21	ADAMTS18	Female	2.25 (1.23–4.10)	<0.01
	Male	1.43 (1.06–1.92)	0.02		Male	1.36 (0.92–2.02)	0.13
ADAMTS19	Female	1.22 (0.79–1.87)	0.36	ADAMTS19	Female	0.89 (0.49–1.60)	0.70
	Male	0.96 (0.72–1.30)	0.81		Male	1.26 (0.85–1.86)	0.25
ADAMTS20	Female	1.98 (1.38–2.83)	<0.01	ADAMTS20	Female	1.14 (0.64–2.06)	0.65
	Male	1.67 (1.35–2.07)	<0.01		Male	1.12 (0.75–1.65)	0.58

with poor OS among women (Table 2). Table 3 contains the prognosis of ADAMTS in the tumor stratification group. The expression of genes such as *ADAMTS6*, 7, 12, 15, 16, and 18 was closely related to poor OS in grade III.

3.3. Validation of *ADAMTS12* mRNA Expression Level. By screening 38 GC chips from the GEO database (a total of 1559 GC samples and 883 control samples), we found that the expression level of *ADAMTS12* was significantly increased in GC (SMD = 0.73, 95% CI (0.32–1.14) ($P < 0.05$) (Figure 9)). Thereafter, by performing qRT-PCR using 20 pairs of clinical samples, we found that the expression level of *ADAMTS12* in GC tissues was significantly higher than that in adjacent tissues ($P = 0.024$) (Figure 10).

3.4. *ADAMTS12* Functional Enrichment Analysis. By analyzing the chip data in the Coexpedia database, we found

that the *ADAMTS12* expression-related genes were *COL5A2*, *COL5A1*, *COL1A2*, *SPARC*, *COL3A1*, *COL6A3*, *ADAMTS12*, *ADAMTS2*, *FBN1*, *COL1A1*, and other related genes (Figure 11(a)). Similarly, by analyzing genes related to *ADAMTS12* expression using the RNA-seq data, we found that *ADAMTS2*, *COL5A2*, *FAP*, *COL1A2*, *COL1A1*, *SPARC*, *SULF1*, *COL5A1*, *COL3A1*, and *COL12A1* were the genes closely related to its expression (Figure 11(b)). Through functional enrichment analysis of the top 100 expression-related genes, we found that these genes were closely related to signaling pathways such as protein digestion and absorption, ECM-receptor interaction, and focal adhesion (Figure 12). Through GO analysis, we found that *ADAMTS12* and other related genes were significantly enriched in the ECM organization (biological process), ECM structural constituent (molecular function), and ECM (cellular component) (Figure 13).

FIGURE 9: Forest plot of *ADAMTS12* expression level. Meta-analysis revealed that the expression level of *ADAMTS12* in the cancer patients was higher than that in noncancer patients (standardized mean difference (SMD): 0.73 (0.32–1.14), $P < 0.001$).

4. Discussion

In the present study, we sought to explore the significance of the expression of the ADAMTS family of genes in the prognosis of GC by using a K-M plotter. Thus, the expression of this family of genes in GC was analyzed. According to the RNA-seq and gene chip data, most genes in the ADAMTS family (6, 7, 12, 15, and 18) were found to be closely related to the prognosis of GC. In addition, their high expression was found to be associated with poor prognosis and survival time. By exploring gene mutations of the ADAMTS family, we recognized that it was indeed the most frequent cause of gene alterations. Furthermore, through PCR analysis, we could confirm that the expression level of *ADAMTS12* in GC was consistent with the sequencing data.

Based on the studies conducted on the ADAMTS family, we recognized that most were related to osteoarthritis, blood vessels, and diseases of the platelet [13]. In the present study, however, we primarily discussed prior researches and the role of the ADAMTS family of genes in cancers.

ADAMTS1 is highly expressed in pancreatic cancer and ovarian cancer [14, 15]. Although a study found that its expression level was elevated in GC [6], other studies have found that it displays a low expression level in this malignancy [16, 17]. As these findings align with the sequencing data obtained in our study, its expression level in GC is yet to be clarified. Through a study of *ADAMTS1* in breast cancer, its expression level was found to be low [18–21]. However, another study found that it can promote breast cancer metastasis [22].

FIGURE 10: The expression of *ADAMTS12* in gastric cancer patients. The expression level of *ADAMTS12* was higher in GC tissues compared to that in adjacent noncancerous tissues ($P < 0.05$).

FIGURE 11: Coexpression networks. The coexpressed genes of *ADAMTS12* in (a) RNA-seq data and (b) chip data. In both network diagrams, color and size represent the degree of correlation, with the blue and large circles depicting a high degree of correlation.

ADAMTS2 is highly expressed in GC and colon cancer and is closely related to distant metastasis and the prognosis of colon cancer. However, it is an independent prognostic factor for GC [23, 24]. We found that the overexpression of *ADAMTS2* was associated with unfavorable OS in GC patients, which aligns with the findings of the abovementioned study.

ADAMTS3 is involved in the alteration of osteosarcoma matrices [25], and the single nucleotide polymorphism (SNP) of the *ADAMTS3* gene is an independent prognostic biomarker for cutaneous melanoma [26].

ADAMTS4 is highly expressed in colon cancer and is associated with poor prognosis [27]. *ADAMTS4* was also found to be mutated in early-onset familial colorectal cancer [28].

The role of *ADAMTS5* in digestive system tumors is unclear, but its expression level was confirmed to be low in GC, colon cancer, and liver cancer [29–31]. A study found that *ADAMTS5* overexpresses colon cancer cells to inhibit their invasion and migration [30]. However, Yu et al. found that it is highly expressed in colon cancer and is associated with lymph node metastasis [32]. A high expression level of *ADAMTS5* promotes the invasion and migration of human glioblastoma, non-small-cell lung cancer, and HNSCC cells [33, 34].

ADAMTS6 is highly expressed in esophageal cancer and is an independent marker of its prognosis [35]. Notably, *ADAMTS6* displays the opposite trend in expression in colon and rectal cancer [36, 37]. In the current analysis, we found that a high expression level of

FIGURE 12: (a) KEGG analysis of the coexpressed genes from the RNA-seq data revealed that protein digestion and absorption was the most significantly enriched pathway. (b) KEGG analysis of the coexpressed genes from the chip data also revealed that the pathway of protein digestion and absorption was significantly enriched.

ADAMTS6 was significantly correlated with favorable survival in GC patients.

Presently, only few reports exist on the direction of *ADAMTS7* tumors. Among the genetic mutations in liver cancer, the Asian *ADAMTS7* mutation is only found in Asian Americans [38].

A study revealed that the *ADAMTS8* gene is highly methylated in GC while its mRNA expression level is significantly reduced [7]. *ADAMTS8* is underexpressed in liver

cancer and affects its progression by targeting the ERK signaling pathway [39]. Conversely, it is highly expressed in head and neck squamous cell carcinoma [40]. Porter et al. demonstrated that the expression level of *ADAMTS8* is a predictor of poor OS [41].

ADAMTS9 is poorly expressed in breast cancer, colorectal cancer, and GC and is associated with the hypermethylation of their promoters [42–44]. *ADAMTS9* can also inhibit tumor progression by inhibiting angiogenesis [45].

FIGURE 13: GO functional analysis of the coexpressed genes. The coexpressed genes of the (a) RNA-seq data and (b) chip data. The top 10 genes for biological process, molecular function, and cellular component term are displayed.

ADAMTS12 is associated with ovarian and renal cancer metastases, and its expression level is significantly elevated in metastatic tumors compared to primary tumors [46, 47]. *ADAMTS12* is also highly expressed in esophageal squamous cell carcinoma [48] but has low expression in colon cancer [49]. A previous study revealed that *ADAMTS12* has anti-tumor-growth and angiogenesis effects [50]. In the present study, the data clearly indicate that a higher expression level of *ADAMTS12* was significantly correlated with poor prognosis in patients with GC.

Garam et al. found that *ADAMTS13* is a prognostic risk factor for colon and liver cancer [51]. The single nucleotide polymorphism of *ADAMTS14* is closely related to liver cancer and oral cancer [52, 53].

ADAMTS15 inhibits tumor cell migration and angiogenesis in breast cancer and serves as an independent prognostic factor of this cancer type [41, 54]. *ADAMTS15* also has a negative correlation with the degree of tumor tissue differentiation in colorectal cancer and inhibits the growth and invasion ability of colon cancer cells [55]. According to our research, *ADAMTS15* has a low expression level in GC and may also exhibit an inhibitory effect.

The DNA of *ADAMTS16* displays a hypermethylation state in colorectal tissue and can inhibit tumor proliferation [56]. However, *ADAMTS16* was found to be highly expressed in the esophagus, and its knockout inhibited tumor invasion [57].

ADAMTS18 is a gene that has received considerable attention. In GC, colorectal cancer, pancreatic cancer, lung cancer, breast cancer, esophageal cancer, and nasopharyngeal carcinoma, its low expression level was found to be associated with the abnormal methylation of their promoter [58–63]. Similar studies have also found that *ADAMTS19* is highly methylated in colorectal cancer [64]. Evidently, *ADAMTS20* causes somatic mutation in GC patients with liver metastases [65]. In the present study, *ADAMTS18* was found to be correlated with unfavorable prognosis in GC patients.

In conclusion, the present study was carried out to determine the mRNA expression levels of the ADAMTS family of genes in GC and the prognostic value of their expression using a KM survival curve. As a result, the expression of the ADAMTS family of genes was found to be closely related to the poor prognosis of GC patients. Such findings could enable a better understanding of the

prognostic function of the ADAMTS family of genes in GC. Furthermore, these findings may serve as a favorable predictor of GC prognosis, ultimately contributing to the design of strategies.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Authors' Contributions

Liang Liang and Jin-hui Zhu equally contributed to this work.

Acknowledgments

The present study was supported by grants from the Key Research and Development Program of Science and Technology Department of Guangxi (grant no. 2017AB45153), the Innovation Project of Guangxi Graduate Education (grant no. YCBZ2019043), and the Education Department Foundation for Innovation Team of Guangxi Zhuang Autonomous Region.

References

- [1] F. Bray, J. Ferlay, I. Soerjomataram, R. L. Siegel, L. A. Torre, and A. Jemal, "Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries," *CA: A Cancer Journal for Clinicians*, vol. 68, no. 6, pp. 394–424, 2018.
- [2] J. A. Ajani, T. A. D'Amico, K. Almhanna et al., "Gastric cancer, version 3.2016, NCCN clinical practice guidelines in oncology," *Journal of the National Comprehensive Cancer Network: JNCCN*, vol. 14, no. 10, pp. 1286–1312, 2016.
- [3] H. Wilke, K. Muro, E. Van Cutsem et al., "Ramucirumab plus paclitaxel versus placebo plus paclitaxel in patients with previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (RAINBOW): a double-blind, randomised phase 3 trial," *The Lancet Oncology*, vol. 15, no. 11, pp. 1224–1235, 2014.
- [4] M. J. Binder, S. McCoombe, E. D. Williams, D. R. McCulloch, and A. C. Ward, "The extracellular matrix in cancer progression: role of hyaluronan proteoglycans and ADAMTS enzymes," *Cancer Letters*, vol. 385, pp. 55–64, 2017.
- [5] S. Cal and C. López-Otín, "ADAMTS proteases and cancer," *Matrix Biology*, vol. 44–46, pp. 77–85, 2015.
- [6] M. O. Kilic, B. Aynekin, A. Kara, D. Icen, and K. Demircan, "Differentially regulated ADAMTS1, 8, and 18 in gastric adenocarcinoma," *Bratislava Medical Journal*, vol. 118, no. 02, pp. 71–76, 2017.
- [7] J. Chen, J. Zhang, X. Li et al., "Downregulation of ADAMTS8 by DNA hypermethylation in gastric cancer and its clinical significance," *BioMed Research International*, vol. 2016, Article ID 5083841, 9 pages, 2016.
- [8] S. Cal, J. M. Argüelles, P. L. Fernández, and C. López-Otín, "Identification, characterization, and intracellular processing of ADAM-TS12, a novel human disintegrin with a complex structural organization involving multiple thrombospondin-1 repeats," *Journal of Biological Chemistry*, vol. 276, no. 21, pp. 17932–17940, 2001.
- [9] Z. Tang, C. Li, B. Kang, G. Gao, C. Li, and Z. Zhang, "GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses," *Nucleic Acids Research*, vol. 45, no. W1, pp. W98–W102, 2017.
- [10] E. Cerami, J. Gao, U. Dogrusoz et al., "The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data: figure 1," *Cancer Discovery*, vol. 2, no. 5, pp. 401–404, 2012.
- [11] A. M. Szász, A. Lánckzy, Á. Nagy et al., "Cross-validation of survival associated biomarkers in gastric cancer using transcriptomic data of 1,065 patients," *Oncotarget*, vol. 7, no. 31, pp. 49322–49333, 2016.
- [12] S. Yang, C. Y. Kim, S. Hwang et al., "COEXPEDIA: exploring biomedical hypotheses via co-expressions associated with medical subject headings (MeSH)," *Nucleic Acids Research*, vol. 45, no. D1, pp. D389–D396, 2017.
- [13] E. Zampeli, F. N. Skopouli, and H. M. Moutsopoulos, "Polyserositis in a patient with active systemic lupus erythematosus: a case of pseudo-pseudo meigs syndrome," *The Journal of Rheumatology*, vol. 45, no. 6, pp. 877–878, 2018.
- [14] M. O. Kilic, B. Aynekin, M. Bozer et al., "Differentially regulated ADAMTS1, 8, 9, and 18 in pancreas adenocarcinoma," *Przegląd Gastroenterologiczny*, vol. 12, no. 4, pp. 262–270, 2017.
- [15] M. A. Lima, L. Dos Santos, J. A. Turri et al., "Prognostic value of ADAMTS proteases and their substrates in epithelial ovarian cancer," *Pathobiology*, vol. 83, no. 6, pp. 316–326, 2016.
- [16] J. Chen, C. Zhang, X. Xu, X. Zhu, and D. Dai, "Down-regulation of A disintegrin and metalloproteinase with thrombospondin motif type 1 by DNA hypermethylation in human gastric cancer," *Molecular Medicine Reports*, vol. 12, no. 2, pp. 2487–2494, 2015.
- [17] J. Chen, Y. Zhi, X. Chang, S. Zhang, and D. Dai, "Expression of ADAMTS1 and its correlation with angiogenesis in primary gastric cancer and lymph node metastasis," *Digestive Diseases and Sciences*, vol. 58, no. 2, pp. 405–413, 2013.
- [18] M. Li, L. Liu, W. Zang et al., "miR-365 overexpression promotes cell proliferation and invasion by targeting ADAMTS-1 in breast cancer," *International Journal of Oncology*, vol. 47, no. 1, pp. 296–302, 2015.
- [19] E. Martino-Echarri, R. Fernández-Rodríguez, F. J. Rodríguez-Baena et al., "Contribution of ADAMTS1 as a tumor suppressor gene in human breast carcinoma. Linking its tumor inhibitory properties to its proteolytic activity on nidogen-1 and nidogen-2," *International Journal of Cancer*, vol. 133, no. 10, pp. 2315–2324, 2013.
- [20] V. M. Freitas, J. B. do Amaral, T. A. Silva et al., "Decreased expression of ADAMTS-1 in human breast tumors stimulates migration and invasion," *Molecular Cancer*, vol. 12, no. 1, p. 2, 2013.
- [21] S. Malvia, S. A. R. Bagadi, P. Pradhan et al., "Study of gene expression profiles of breast cancers in Indian women," *Scientific Reports*, vol. 9, no. 1, p. 10018, 2019.
- [22] I. A. Tan, K. Frewin, C. Ricciardelli, and D. L. Russell, "ADAMTS1 promotes adhesion to extracellular matrix proteins and predicts prognosis in early stage breast cancer

- patients," *Cellular Physiology and Biochemistry: International Journal of Experimental Cellular Physiology, Biochemistry, and Pharmacology*, vol. 52, no. 6, pp. 1553–1568, 2019.
- [23] C. Jiang, Y. Zhou, Y. Huang, Y. Wang, W. Wang, and X. Kuai, "Overexpression of ADAMTS-2 in tumor cells and stroma is predictive of poor clinical prognosis in gastric cancer," *Human Pathology*, vol. 84, pp. 44–51, 2019.
 - [24] C. Kirana, L. Peng, R. Miller et al., "Combination of laser microdissection, 2D-DIGE and MALDI-TOF MS to identify protein biomarkers to predict colorectal cancer spread," *Clinical Proteomics*, vol. 16, p. 3, 2019.
 - [25] A. T. Aydemir, M. Alper, and F. Kockar, "SP1-mediated downregulation of ADAMTS3 gene expression in osteosarcoma models," *Gene*, vol. 659, pp. 1–10, 2018.
 - [26] Y. Xu, Y. Wang, H. Liu et al., "Genetic variants in the metzincin metallopeptidase family genes predict melanoma survival," *Molecular Carcinogenesis*, vol. 57, no. 1, pp. 22–31, 2018.
 - [27] J. Chen, Y. Luo, Y. Zhou et al., "Promotion of tumor growth by ADAMTS4 in colorectal cancer: focused on macrophages," *Cellular Physiology and Biochemistry*, vol. 46, no. 4, pp. 1693–1703, 2018.
 - [28] N. Rao, Z. Ke, H. Liu et al., "ADAMTS4 and its proteolytic fragments differentially affect melanoma growth and angiogenesis in mice," *International Journal of Cancer*, vol. 133, no. 2, pp. 294–306, 2013.
 - [29] J. Huang, Y. Sun, H. Chen et al., "ADAMTS5 acts as a tumor suppressor by inhibiting migration, invasion and angiogenesis in human gastric cancer," *Gastric Cancer*, vol. 22, no. 2, pp. 287–301, 2019.
 - [30] J. Li, Y. Liao, J. Huang et al., "Epigenetic silencing of ADAMTS5 is associated with increased invasiveness and poor survival in patients with colorectal cancer," *Journal of Cancer Research and Clinical Oncology*, vol. 144, no. 2, pp. 215–227, 2018.
 - [31] C. Li, Y. Xiong, X. Yang et al., "Lost expression of ADAMTS5 protein associates with progression and poor prognosis of hepatocellular carcinoma," *Drug Design, Development and Therapy*, vol. 9, pp. 1773–1783, 2015.
 - [32] L. Yu, Y. Lu, X. Han et al., "microRNA -140-5p inhibits colorectal cancer invasion and metastasis by targeting ADAMTS5 and IGFBP5," *Stem Cell Research & Therapy*, vol. 7, no. 1, p. 180, 2016.
 - [33] J. Gu, J. Chen, J. Feng et al., "Overexpression of ADAMTS5 can regulate the migration and invasion of non-small cell lung cancer," *Tumor Biology*, vol. 37, no. 7, pp. 8681–8689, 2016.
 - [34] J. Zhang, X. Qin, Q. Sun et al., "Transcriptional control of PAX4-regulated miR-144/451 modulates metastasis by suppressing ADAMs expression," *Oncogene*, vol. 34, no. 25, pp. 3283–3295, 2015.
 - [35] L. Liu, Z. Yang, W. Ni, and Y. Xuan, "ADAMTS-6 is a predictor of poor prognosis in patients with esophageal squamous cell carcinoma," *Experimental and Molecular Pathology*, vol. 104, no. 2, pp. 134–139, 2018.
 - [36] Y. Xie, Q. Gou, K. Xie, Z. Wang, Y. Wang, and H. Zheng, "ADAMTS6 suppresses tumor progression via the ERK signaling pathway and serves as a prognostic marker in human breast cancer," *Oncotarget*, vol. 7, no. 38, pp. 61273–61283, 2016.
 - [37] W. H. Xiao, X. L. Qu, X. M. Li et al., "Identification of commonly dysregulated genes in colorectal cancer by integrating analysis of RNA-Seq data and qRT-PCR validation," *Cancer Gene Therapy*, vol. 22, no. 5, pp. 278–284, 2015.
 - [38] S. Sun, C. Johnson, Q. Hu et al., "Differences in somatic mutation landscape of hepatocellular carcinoma in Asian American and European American populations," *Oncotarget*, vol. 7, no. 26, pp. 40491–40499, 2016.
 - [39] X. Zhao, C. Yang, J. Wu, and Y. Nan, "ADAMTS8 targets ERK to suppress cell proliferation, invasion, and metastasis of hepatocellular carcinoma," *OncoTargets and Therapy*, vol. 11, pp. 7569–7578, 2018.
 - [40] A. Stokes, J. Joutsa, R. Ala-aho et al., "Expression profiles and clinical correlations of degradome components in the tumor microenvironment of head and neck squamous cell carcinoma," *Clinical Cancer Research*, vol. 16, no. 7, pp. 2022–2035, 2010.
 - [41] S. Porter, P. N. Span, F. C. G. J. Sweep et al., "ADAMTS8 and ADAMTS15 expression predicts survival in human breast carcinoma," *International Journal of Cancer*, vol. 118, no. 5, pp. 1241–1247, 2006.
 - [42] B. Shao, Y. Feng, H. Zhang et al., "The 3p14.2 tumour suppressor ADAMTS9 is inactivated by promoter CpG methylation and inhibits tumour cell growth in breast cancer," *Journal of Cellular and Molecular Medicine*, vol. 22, no. 2, pp. 1257–1271, 2018.
 - [43] L. Chen, J. Tang, Y. Feng et al., "ADAMTS9 is silenced by epigenetic disruption in colorectal cancer and inhibits cell growth and metastasis by regulating Akt/p53 signaling," *Cellular Physiology and Biochemistry*, vol. 44, no. 4, pp. 1370–1380, 2017.
 - [44] W. Du, S. Wang, Q. Zhou et al., "ADAMTS9 is a functional tumor suppressor through inhibiting AKT/mTOR pathway and associated with poor survival in gastric cancer," *Oncogene*, vol. 32, no. 28, pp. 3319–3328, 2013.
 - [45] P. H. Y. Lo, H. L. Lung, A. K. L. Cheung et al., "Extracellular protease ADAMTS9 suppresses esophageal and nasopharyngeal carcinoma tumor formation by inhibiting angiogenesis," *Cancer Research*, vol. 70, no. 13, pp. 5567–5576, 2010.
 - [46] T. H. Ho, D. J. Serie, M. Parasramka et al., "Differential gene expression profiling of matched primary renal cell carcinoma and metastases reveals upregulation of extracellular matrix genes," *Annals of Oncology*, vol. 28, no. 3, pp. 604–610, 2017.
 - [47] A. Mariani, C. Wang, A. L. Oberg et al., "Genes associated with bowel metastases in ovarian cancer," *Gynecologic Oncology*, vol. 154, no. 3, pp. 495–504, 2019.
 - [48] X. Li, X. Xiao, R. Chang, and C. Zhang, "Comprehensive bioinformatics analysis identifies lncRNA HCG22 as a migration inhibitor in esophageal squamous cell carcinoma," *Journal of Cellular Biochemistry*, vol. 121, no. 1, pp. 468–481, 2019.
 - [49] D. Wang, T. Zhu, F.-B. Zhang, and C. He, "Expression of ADAMTS12 in colorectal cancer-associated stroma prevents cancer development and is a good prognostic indicator of colorectal cancer," *Digestive Diseases and Sciences*, vol. 56, no. 11, pp. 3281–3287, 2011.
 - [50] M. El Hour, A. Moncada-Pazos, S. Blacher et al., "Higher sensitivity of Adamts12-deficient mice to tumor growth and angiogenesis," *Oncogene*, vol. 29, no. 20, pp. 3025–3032, 2010.
 - [51] N. Garam, É. Maláti, G. Sinkovits et al., "Platelet count, ADAMTS13 activity, von Willebrand factor level and survival in patients with colorectal cancer: 5-year follow-up study," *Thrombosis and Haemostasis*, vol. 118, no. 1, pp. 123–131, 2018.
 - [52] M. J. Sheu, M. J. Hsieh, Y. E. Chou et al., "Effects of ADAMTS14 genetic polymorphism and cigarette smoking on the clinicopathologic development of hepatocellular carcinoma," *PLoS One*, vol. 12, no. 2, Article ID e0172506, 2017.
 - [53] S. C. Su, M. J. Hsieh, Y. F. Liu et al., "ADAMTS14 gene polymorphism and environmental risk in the development of

- oral cancer," *PLoS One*, vol. 11, no. 7, Article ID e0159585, 2016.
- [54] R. Kelwick, L. Wagstaff, J. Decock et al., "Metalloproteinase-dependent and -independent processes contribute to inhibition of breast cancer cell migration, angiogenesis and liver metastasis by a disintegrin and metalloproteinase with thrombospondin motifs-15," *International Journal of Cancer*, vol. 136, no. 4, pp. E14–E26, 2015.
 - [55] C. G. Vilorio, A. J. Obaya, M.-A. Pazos et al., "Genetic inactivation of ADAMTS15 metalloprotease in human colorectal cancer," *Cancer Research*, vol. 69, no. 11, pp. 4926–4934, 2009.
 - [56] F. Moncada-Pazos, J. Kolarova, S. Schafmayer et al., "Aberrant DNA methylation of ADAMTS16 in colorectal and other epithelial cancers," *BMC Cancer*, vol. 18, no. 1, p. 796, 2018.
 - [57] N. Sakamoto, N. Oue, T. Noguchi et al., "Serial analysis of gene expression of esophageal squamous cell carcinoma: ADAMTS16 is upregulated in esophageal squamous cell carcinoma," *Cancer Science*, vol. 101, no. 4, pp. 1038–1044, 2010.
 - [58] Y. Zhang, H. Xu, J. Mu et al., "Inactivation of ADAMTS18 by aberrant promoter hypermethylation contribute to lung cancer progression," *Journal of Cellular Physiology*, vol. 234, no. 5, pp. 6965–6975, 2019.
 - [59] H. Xu, Q. Xiao, Y. Fan et al., "Epigenetic silencing of ADAMTS18 promotes cell migration and invasion of breast cancer through AKT and NF- κ B signaling," *Cancer Medicine*, vol. 6, no. 6, pp. 1399–1408, 2017.
 - [60] I. Guilleret, L. Losi, S. T. Chelbi et al., "DNA methylation profiling of esophageal adenocarcinoma using Methylation Ligation-dependent Macroarray (MLM)," *Biochemical and Biophysical Research Communications*, vol. 479, no. 2, pp. 231–237, 2016.
 - [61] B. Xu, L. Zhang, C. Luo et al., "Hypermethylation of the 16q23.1 tumor suppressor gene ADAMTS18 in clear cell renal cell carcinoma," *International Journal of Molecular Sciences*, vol. 16, no. 1, pp. 1051–1065, 2015.
 - [62] Z. Li, W. Zhang, Y. C. Shao et al., "High-resolution melting analysis of ADAMTS18 methylation levels in gastric, colorectal and pancreatic cancers," *Medical Oncology*, vol. 27, no. 3, pp. 998–1004, 2010.
 - [63] H. Zhang, X. Wang, J. Ying et al., "Epigenetic identification of ADAMTS18 as a novel 16q23.1 tumor suppressor frequently silenced in esophageal, nasopharyngeal and multiple other carcinomas," *Oncogene*, vol. 26, no. 53, pp. 7490–7498, 2007.
 - [64] S. Alonso, B. Gonzalez, T. Ruiz-Larroya et al., "Epigenetic inactivation of the extracellular matrix metalloproteinase ADAMTS19 gene and the metastatic spread in colorectal cancer," *Clinical Epigenetics*, vol. 7, p. 124, 2015.
 - [65] N. Ikari, A. Serizawa, S. Mitani, M. Yamamoto, and T. Furukawa, "Near-comprehensive resequencing of cancer-associated genes in surgically resected metastatic liver tumors of gastric cancer," *The American Journal of Pathology*, vol. 189, no. 4, pp. 784–796, 2019.