

Research Article

Clinical Validation of Adjusted Corneal Power in Patients with Previous Myopic Lasik Surgery

Vicente J. Camps,¹ David P. Piñero,^{1,2,3} Veronica Mateo,¹ Celia García,¹
Alberto Artola,³ Rafael Pérez-Cambrodi,³ and Pedro Ruiz-Fortes³

¹Department of Optics, Pharmacology and Anatomy, University of Alicante, 03690 San Vicente del Raspeig, Spain

²Foundation for the Visual Quality (FUNCAVIS), 03016 Alicante, Spain

³Department of Ophthalmology, Oftalmar, Medimar International Hospital, 03016 Alicante, Spain

Correspondence should be addressed to Vicente J. Camps; vicente.camps@ua.es

Received 24 July 2015; Revised 2 September 2015; Accepted 17 September 2015

Academic Editor: Vasilios F. Diakonis

Copyright © 2015 Vicente J. Camps et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. To validate clinically a new method for estimating the corneal power (P_c) using a variable keratometric index (n_{kadj}) in eyes with previous laser refractive surgery. **Setting.** University of Alicante and Medimar International Hospital (Oftalmar), Alicante, (Spain). **Design.** Retrospective case series. **Methods.** This retrospective study comprised 62 eyes of 62 patients that had undergone myopic LASIK surgery. An algorithm for the calculation of n_{kadj} was used for the estimation of the adjusted keratometric corneal power (P_{kadj}). This value was compared with the classical keratometric corneal power (P_k), the True Net Power (TNP), and the Gaussian corneal power (P_{cGauss}). Likewise, P_{kadj} was compared with other previously described methods. **Results.** Differences between P_{cGauss} and P_c values obtained with all methods evaluated were statistically significant ($p < 0.01$). Differences between P_{kadj} and P_{cGauss} were in the limit of clinical significance ($p < 0.01$, loA [-0.33,0.60] D). Differences between P_{kadj} and TNP were not statistically and clinically significant ($p = 0.319$, loA [-0.50,0.44] D). Differences between P_{kadj} and previously described methods were statistically significant ($p < 0.01$), except with $P_{cHaigisL}$ ($p = 0.09$, loA [-0.37,0.29] D). **Conclusion.** The use of the adjusted keratometric index (n_{kadj}) is a valid method to estimate the central corneal power in corneas with previous myopic laser refractive surgery, providing results comparable to $P_{cHaigisL}$.

1. Introduction

The precise measurement of corneal power after myopic laser refractive surgery is still currently an issue under debate. Several methods have been proposed during the last years which are classified as methods requiring historical clinical data and methods not requiring historical data. Among those methods requiring previous clinical data, some of them are based on a correction of the corneal power using the refracting change achieved [1–3] and others on performing such correction by adjusting the keratometric index [4–7]. The main disadvantage of all these methods is that they are infeasible if previous clinical patient's data are not available. For this reason, other methods that do not require patient's historical data have been developed [3, 5, 8–11]. In this line, our research group has recently proposed a new method for estimating with enough accuracy the corneal power using

the keratometric approach that has been found to be valid in both healthy [12] and post-LASIK eyes [13]. This algorithm based on a variable keratometric index was named adjusted keratometric index (n_{kadj}) and it has been prevalidated clinically in a sample of 32 eyes that had undergone previously myopic LASIK surgery [13]. The aim of the current study is to validate clinically this algorithm for the estimation of the corneal power in eyes with previous myopic LASIK in a larger population including also a larger range of intended refractive corrections.

2. Methods

2.1. Patients and Examination. This retrospective study comprised 62 eyes of 62 patients that had undergone previous correction of a myopic refractive error by means of laser in

situ keratomileusis (LASIK) surgery. All LASIK surgeries had been performed using the Pulzar Z1 solid-state laser (CustomVis Laser Pty Ltd., Osborne Park, Australia, currently CV Laser Pty Ltd.) at the Department of Ophthalmology (Oftalmar) of the Vithas Medimar International Hospital (Alicante, Spain). All surgeries had been performed by one experienced surgeon (AA) between October 2012 and December 2013. For this study, only one eye from each subject was chosen according to a random number sequence (dichotomic sequence, 0 and 1). A comprehensive ophthalmologic examination was performed in all cases at least 3 months after surgery, which included refraction, corrected distance visual acuity (CDVA), slit lamp biomicroscopy, Goldman tonometry, fundus evaluation, and the analysis of the corneal structure by means of a Scheimpflug photography-based tomographer, the Pentacam system (Oculus Optikgeräte GmbH, Germany, software version 1.14r01). All patients were informed after surgery about this retrospective study and signed an informed consent document in accordance with the Helsinki Declaration.

2.2. Corneal Power Calculation. Our research group recently proposed the use of a variable keratometric index (n_{kadj}) for the estimation of the corneal power (P_c) using the keratometric approach in patients with previous myopic LASIK surgery [13]. The following expression was defined for n_{kadj} considering the ocular conditions of the Gullstrand eye model and the range of anterior and posterior curvature that is commonly found in this kind of patients [13]:

$$n_{kadj} = -0.0064286r_{1c} + 1.37688, \quad (1)$$

where r_{1c} is the postoperative anterior corneal radius.

Furthermore, adjusted keratometric corneal power (P_{kadj}) was defined as follows [13]:

$$P_{kadj} = \frac{n_{kadj} - 1}{r_{1c}}. \quad (2)$$

For comparison purposes, the keratometric corneal power was also calculated using the classical keratometric index $n_k = 1.3375$ ($P_{k(1.3375)}$).

The Gaussian corneal power was calculated using the following expression:

$$\begin{aligned} P_{cGauss} &= P_{1c} + P_{2c} - \delta P_{1c} P_{2c} \\ &= \frac{n_c - n_a}{r_{1c}} + \frac{n_{ha} - n_c}{r_{2c}} - \frac{e_c}{n_c} \cdot \frac{n_c - n_a}{r_{1c}} \cdot \frac{n_{ha} - n_c}{r_{2c}}, \end{aligned} \quad (3)$$

where P_{cGauss} is the Gaussian total corneal power, P_{1c} is the anterior corneal power, P_{2c} is the posterior corneal power, r_{1c} is the anterior corneal radius, r_{2c} is the posterior corneal radius, n_a is the refractive index of air, n_c is the refractive index of the cornea, n_{ha} is the refractive index of the aqueous humour, and e_c is the central corneal thickness.

Likewise, the True Net Power (TNP) was also recorded, which is the corneal power provided by the Pentacam system (Oculus) based on the anterior (r_{1c}) and posterior (r_{2c}) corneal radius and calculated by using the Gaussian equation

(P_{cGauss}) with the Gullstrand eye model, but neglecting the corneal thickness (e_c):

$$\begin{aligned} \text{True net Power (TNP)} \\ &= \frac{1.376 - 1}{r_{1c}} \cdot 1000 + \frac{1.336 - 1.376}{r_{2c}} \cdot 1000. \end{aligned} \quad (4)$$

Besides this, corneal power was also estimated by using other methods described previously for such purpose in eyes with previous myopic laser refractive surgery:

(1) Methods requiring previous clinical data:

(a) Awwad method [3]:

$$P_{cAwwad} = P_c - 0.23\Delta SE; \quad (5)$$

(b) Camellin method [4]:

$$P_{cCamellin} = \frac{[(1.3319 + 0.00113\Delta SE) - 1]}{(r_{1cpost}/1000)}; \quad (6)$$

(c) Clinical History method:

$$P_{cpost} = P_{cpre} + \Delta SE; \quad (7)$$

(d) Jarade method [4]:

$$P_{cJarade} = \frac{[(1.3375 + 0.0014\Delta SE) - 1]}{(r_{1cpost}/1000)}; \quad (8)$$

(e) Savini method [5, 6]:

$$P_{cSavini} = \frac{[(1.338 + 0.0009856\Delta SE) - 1]}{(r_{1cpost}/1000)}. \quad (9)$$

(2) Methods not requiring previous data:

(a) Haigis-L method:

$$P_{cHaigisL} = -5.1625r_{1cpost} + 82.2603 - 0.35; \quad (10)$$

(b) Shammas method [8]:

$$P_{cShammas} = 1.14(P_c - 6.8); \quad (11)$$

(c) Seitz method [6]:

$$P_{cSeitz} = 1.114(P_c - 4.98), \quad (12)$$

where $\Delta SE = SE_{pre} - SE_{post}$, SE_{pre} and SE_{post} being the pre- and postsurgery spherical equivalents, r_{1cpost} is the postsurgery anterior corneal radius, and P_{cpre} is the presurgery keratometric corneal power.

For the clinical validation of P_{kadj} , it was compared with P_{cGauss} and TNP. Likewise, the different methods mentioned above were also compared with P_{kadj} and P_{cGauss} in order to demonstrate which was the most accurate approach.

2.3. Statistical Analysis. Statistical analysis was performed using the software SPSS version 19.0 for Windows (SPSS, Chicago, Illinois, USA). Normality of all data distributions was first confirmed by means of the Kolmogorov-Smirnov test. Specifically, the paired Student t -test or Wilcoxon test was used for comparing the different methods of P_c calculation depending on whether the normality condition could be assumed or not. The Bland-Altman analysis [14] was used for evaluating the agreement and interchangeability of the different methods for obtaining the corneal power.

3. Results

This study comprised 62 eyes of 62 patients (34 women [54.8%]), with a mean age of 33.42 ± 7.16 years (range 21 to 52 years) and with preoperative myopia between -0.25 and -6.8 D. The sample comprised 31 left eyes (50%). Mean ocular features of the eyes evaluated in the current study can be seen in Table 1. Table 2 shows the values of corneal power estimated with the previously published methods (Awwad, Camellin, Clinical History, Haigis-L, Jarade, Savini, Seitz, and Shammas methods).

3.1. Clinical Validation of P_{kadj} . As shown in Table 3, there were significant differences ($p < 0.01$, paired Student's t -test) between P_{kadj} and P_{cGauss} , but not ($p = 0.319$, paired Student's t -test) between P_{kadj} and TNP. The Bland-Altman analysis showed that differences between P_{kadj} and P_{cGauss} were barely clinically significant (mean difference: 0.14 ± 0.24 ; limits of agreement (loA): $[-0.33, 0.60$ D]) and that differences between P_{kadj} and TNP were not clinically significant (mean difference 0.03 ± 0.24 ; loA: $[-0.50, 0.44$ D]). A very strong and statistically significant correlation was found between P_{kadj} and P_{cGauss} ($r = 0.994$, $p < 0.01$) as well as between P_{kadj} and TNP ($r = 0.994$, $p < 0.01$).

3.2. Comparison between P_{kadj} and Corneal Power Values Estimated with Other Methods. As shown also in Table 3, differences between P_{kadj} and the rest of the methods for estimation of P_c were statistically significant ($p < 0.01$), except for the difference between P_{kadj} and $P_{cHaigisL}$ ($p = 0.09$). The Bland-Altman analysis confirmed that all these statistically significant differences were also clinically relevant as the ranges of agreement were quite large ($> \pm 0.5$ D). Only differences between P_{kadj} and $P_{cHaigisL}$ (mean: -0.04 ± 0.17 D) did not reach clinical significance (loA: $[-0.37$ to 0.29 D]).

Table 4 summarizes the results of the comparison between the corneal power calculated considering the curvature of the two corneal surfaces as well as corneal thickness (P_{cGauss}) and those values obtained with the other previously published methods for corneal power estimation in corneas with previous myopic laser refractive surgery. As shown, all differences between P_{cGauss} and P_c values obtained with such methods were statistically significant ($p < 0.01$). Considering the Bland-Altman analysis, P_{kadj} and $P_{cHaigisL}$ provided the lower mean differences with P_{cGauss} (-0.14 ± 0.24 D and -0.18 ± 0.21 D, resp.) and the smaller ranges of agreement ($[-0.6$ to $0.33]$ and $[-0.60$ to $0.23]$, resp.). $P_{cCamellin}$ was also close

TABLE 1: Mean ocular features of the eyes evaluated in the current study.

Parameter	Mean \pm SD	Range
SE_{pre} (D)	-3.0 ± 1.6	-6.8 to 0.0
SE_{post} (D)	0.1 ± 0.3	0.0 to 1.0
r_{1cpre} (mm)	7.70 ± 0.25	7.2 to 8.3
r_{1cpost} (mm)	8.18 ± 0.34	7.5 to 9.3
r_{2cpre} (mm)	6.34 ± 0.24	5.9 to 6.9
r_{2cpost} (mm)	6.37 ± 0.24	5.8 to 7.0
e_c (μm)	512 ± 37	407 to 590

SE_{pre} = preoperative spherical equivalent; SE_{post} = postoperative spherical equivalent; r_{1cpre} = preoperative radius of curvature of the anterior corneal surface; r_{1cpost} = postoperative radius of curvature of the anterior corneal surface; r_{2cpre} = preoperative radius of curvature of the posterior corneal surface; r_{2cpost} = postoperative radius of curvature of the posterior corneal surface; e_c = corneal thickness.

TABLE 2: Corneal power measured with different methods.

P_c calculation method	Mean \pm SD	Range
P_{kadj} (D)	39.70 ± 1.86	34.21 to 43.59
P_{cGauss} (D)	39.84 ± 1.72	34.58 to 43.38
TNP (D)	39.73 ± 1.71	34.49 to 43.26
$P_{k(1.3375)}$ (D)	41.31 ± 1.66	36.34 to 44.79
<i>Methods requiring previous data</i>		
P_{cAwwad} (D)	40.60 ± 1.92	34.84 to 44.53
$P_{cCamellin}$ (D)	40.25 ± 1.74	35.07 to 43.90
P_{cCHM} (D)	40.79 ± 2.11	33.94 to 44.80
$P_{cJarade}$ (D)	40.81 ± 1.80	35.45 to 44.58
$P_{cSavini}$ (D)	41.00 ± 1.77	35.72 to 44.71
<i>Methods not requiring previous data</i>		
$P_{cHaigisL}$ (D)	39.66 ± 1.74	34.03 to 43.01
P_{cSeitz} (D)	41.04 ± 1.85	35.56 to 44.92
$P_{cShammas}$ (D)	40.29 ± 1.90	34.68 to 44.26

P_{kadj} = adjusted keratometric corneal power; P_{cGauss} = Gaussian corneal power; TNP = True Net Power; $P_{k(1.3375)}$ = keratometric corneal power using $n_k = 1.3375$; P_{cAwwad} = corneal power obtained using Awwad formula; $P_{cCamellin}$ = corneal power obtained using Camellin formula; P_{cCHM} = corneal power obtained using Clinical History method; $P_{cHaigisL}$ = corneal power obtained using Haigis-L formula; $P_{cJarade}$ = corneal power obtained using Jarade formula; $P_{cSavini}$ = corneal power obtained using Savini formula; P_{cSeitz} = corneal power obtained using Seitz formula; $P_{cShammas}$ = corneal power obtained using Shammas formula.

to P_{cGauss} (mean difference: 0.42 ± 0.14 D), but the range of agreement was larger than that found for P_{kadj} and $P_{cHaigisL}$ ($[0.15$ to 0.68 D]).

4. Discussion

In the current study, the use of P_{kadj} as a method to estimate the corneal power in corneas with previous myopic laser refractive surgery has been validated clinically. Furthermore, the results show that previously reported methods for this estimation are not better than the calculation of P_{kadj} . We have confirmed the clinical prevalidation we conducted in 32 eyes that had undergone myopic LASIK surgery where

TABLE 3: Bland and Altman analysis outcomes of the comparison between $P_{k_{\text{adj}}}$ and the corneal power obtained with other methods.

	$\Delta P_c \pm \text{SD}$ (D)	LoA (D)	Range (D)	p value
$P_{c\text{Gauss}} - P_{k_{\text{adj}}}$ (D)	0.14 ± 0.24	-0.33 to 0.60	-0.43 to 0.70	<0.01
TNP - $P_{k_{\text{adj}}}$ (D)	0.03 ± 0.24	-0.50 to 0.44	-0.55 to 0.61	0.319
$P_{k(1.3375)} - P_{k_{\text{adj}}}$ (D)	1.61 ± 0.19	1.23 to 1.99	1.20 to 2.18	<0.01
<i>Methods requiring previous data</i>				
$P_{c\text{Awwad}} - P_{k_{\text{adj}}}$ (D)	0.90 ± 0.32	0.26 to 1.53	0.04 to 1.65	<0.01
$P_{c\text{Camellin}} - P_{k_{\text{adj}}}$ (D)	0.56 ± 0.20	0.15 to 0.96	0.04 to 1.02	<0.01
$P_{c\text{CHM}} - P_{k_{\text{adj}}}$ (D)	1.09 ± 0.84	-0.55 to 2.73	-1.13 to 3.96	<0.01
$P_{c\text{Jarade}} - P_{k_{\text{adj}}}$ (D)	1.11 ± 0.23	0.67 to 1.55	0.49 to 1.65	<0.01
$P_{c\text{Savini}} - P_{k_{\text{adj}}}$ (D)	1.30 ± 0.19	0.93 to 1.67	0.80 to 1.72	<0.01
<i>Methods not requiring previous data</i>				
$P_{c\text{HaigisL}} - P_{k_{\text{adj}}}$ (D)	-0.04 ± 0.17	-0.37 to 0.29	-0.58 to 0.12	0.09
$P_{c\text{Seitz}} - P_{k_{\text{adj}}}$ (D)	1.34 ± 0.00	1.33 to 1.35	1.32 to 1.35	<0.01
$P_{c\text{Shammas}} - P_{k_{\text{adj}}}$ (D)	0.59 ± 0.04	0.52 to 0.67	0.48 to 0.67	<0.01

$P_{k_{\text{adj}}}$ = adjusted keratometric corneal power; $P_{c\text{Gauss}}$ = Gaussian corneal power; TNP = True Net Power; $P_{k(1.3375)}$ = keratometric corneal power using $n_k = 1.3375$; $P_{c\text{Awwad}}$ = corneal power obtained using Awwad formula; $P_{c\text{Camellin}}$ = corneal power obtained using Camellin formula; $P_{c\text{CHM}}$ = corneal power obtained using Clinical History method; $P_{c\text{HaigisL}}$ = corneal power obtained using Haigis-L formula; $P_{c\text{Jarade}}$ = corneal power obtained using Jarade formula; $P_{c\text{Savini}}$ = corneal power obtained using Savini formula; $P_{c\text{Seitz}}$ = corneal power obtained using Seitz formula; $P_{c\text{Shammas}}$ = corneal power obtained using Shammas formula.

TABLE 4: Bland and Altman analysis outcomes of the comparison between $P_{c\text{Gauss}}$ and the corneal power obtained with other methods.

	$\Delta P_c \pm \text{SD}$ (D)	LoA (D)	Range (D)	p value
$P_{k_{\text{adj}}} - P_{c\text{Gauss}}$ (D)	-0.14 ± 0.24	-0.60 to 0.33	-0.70 to 0.43	<0.01
TNP - $P_{c\text{Gauss}}$ (D)	-0.11 ± 0.01	-0.13 to -0.09	-0.08 to -0.13	<0.01
$P_{k(1.3375)} - P_{c\text{Gauss}}$ (D)	1.47 ± 0.19	1.10 to 1.84	1.07 to 1.97	<0.01
<i>Methods requiring previous data</i>				
$P_{c\text{Awwad}} - P_{c\text{Gauss}}$ (D)	0.76 ± 0.29	0.18 to 1.34	-0.28 to 1.26	<0.01
$P_{c\text{Camellin}} - P_{c\text{Gauss}}$ (D)	0.42 ± 0.14	0.15 to 0.68	0.04 to 0.78	<0.01
$P_{c\text{CHM}} - P_{c\text{Gauss}}$ (D)	0.95 ± 0.83	-0.67 to 2.58	-1.55 to 3.76	<0.01
$P_{c\text{Jarade}} - P_{c\text{Gauss}}$ (D)	0.97 ± 0.18	0.63 to 1.32	0.40 to 1.36	<0.01
$P_{c\text{Savini}} - P_{c\text{Gauss}}$ (D)	1.16 ± 0.14	0.88 to 1.45	0.70 to 1.49	<0.01
<i>Methods not requiring previous data</i>				
$P_{c\text{HaigisL}} - P_{c\text{Gauss}}$ (D)	-0.18 ± 0.21	-0.60 to 0.23	-0.71 to 0.40	<0.01
$P_{c\text{Shammas}} - P_{c\text{Gauss}}$ (D)	0.45 ± 0.26	-0.06 to 0.97	-0.18 to 1.04	<0.01
$P_{c\text{Seitz}} - P_{c\text{Gauss}}$ (D)	1.20 ± 0.23	0.74 to 1.66	0.64 to 1.67	<0.01

$P_{k_{\text{adj}}}$ = adjusted keratometric corneal power; $P_{c\text{Gauss}}$ = Gaussian corneal power; TNP = True Net Power; $P_{k(1.3375)}$ = keratometric corneal power using $n_k = 1.3375$; $P_{c\text{Awwad}}$ = corneal power obtained using Awwad formula; $P_{c\text{Camellin}}$ = corneal power obtained using Camellin formula; $P_{c\text{CHM}}$ = corneal power obtained using Clinical History method; $P_{c\text{HaigisL}}$ = corneal power obtained using Haigis-L formula; $P_{c\text{Jarade}}$ = corneal power obtained using Jarade formula; $P_{c\text{Savini}}$ = corneal power obtained using Savini formula; $P_{c\text{Seitz}}$ = corneal power obtained using Seitz formula; $P_{c\text{Shammas}}$ = corneal power obtained using Shammas formula.

no statistically significant differences between $P_{k_{\text{adj}}}$ and TNP were found, with a mean difference value of 0.00 D and limits of agreement of -0.45 and +0.46 D [13].

In the current series, a mean difference of 0.03 ± 0.24 D between TNP and $P_{k_{\text{adj}}}$ ($p = 0.319$) and a range of agreement from -0.50 to 0.44 D have been found. Besides this, the comparison of $P_{k_{\text{adj}}}$ with $P_{c\text{Gauss}}$ has been also performed. It should be considered that $P_{c\text{Gauss}}$ takes into account neither the curvature of the two corneal surfaces nor the corneal thickness. In this comparison, the mean difference between $P_{k_{\text{adj}}}$ and $P_{c\text{Gauss}}$ was 0.14 ± 0.24 D ($p < 0.01$), with a range

of agreement from -0.33 to 0.60 D. There were only 4 cases out of 66 that showed differences between $P_{k_{\text{adj}}}$ and $P_{c\text{Gauss}}$ of more than ± 0.5 D. All these results are consistent with the theoretical predictions reported previously that estimated maximum differences between $P_{k_{\text{adj}}}$ and $P_{c\text{Gauss}}$ of ± 0.7 D [13]. Therefore, $P_{k_{\text{adj}}}$ is an acceptable method for estimating the corneal power of corneas with previous myopic laser refractive surgery as 100% of estimations were within ± 0.7 D if $P_{c\text{Gauss}}$ is taken as reference. When $P_{k_{\text{adj}}}$ was compared with other different methods of corneal power estimation, statistically significant differences were found ($p < 0.01$),

except for the comparison between $P_{k_{adj}}$ and $P_{c_{HaigisL}}$ ($p = 0.09$). The comparison between $P_{c_{HaigisL}}$ and $P_{k_{adj}}$ showed a mean of difference of -0.04 ± 0.17 and a range of agreement from -0.37 to 0.29 D, confirming that $P_{k_{adj}}$ and $P_{c_{HaigisL}}$ were interchangeable.

Considering that $P_{c_{Gauss}}$ is the most exact method of calculation of the central corneal power in paraxial optics and that $P_{k_{adj}}$, $P_{c_{HaigisL}}$, and $P_{c_{Gauss}}$ can be considered interchangeable according to the results of our study, $P_{k_{adj}}$ and $P_{c_{HaigisL}}$ can be considered appropriate methods for estimating the corneal power in corneas with previous myopic laser refractive surgery when posterior corneal surface data are unknown in clinical practice. The remaining methods (independently if previous historical data are required or not) were found to overestimate $P_{c_{Gauss}}$ (Table 4). The classical keratometric approach for corneal power estimation based on calculations performed only considering the anterior corneal radius ($P_{k(1.3375)}$) induces significant overestimations, ranging from 1.07 to 1.97 D. However, this was not the method providing the poorest performance. The Clinical History method was the most variable procedure, providing corneal power estimations that differed from $P_{c_{Gauss}}$ in a range going from -1.55 to 3.76 D. Among methods requiring previous historical clinical data, $P_{c_{Camellin}}$ was the method providing lower overestimations of corneal power, with differences compared to $P_{c_{Gauss}}$ ranging from 0.04 to 0.78 D. Furthermore, this corneal power estimation method could be clinically valid by using a correction factor. The rest of the methods evaluated led to overestimations of more than 1 D.

Among methods requiring previous historical clinical data, $P_{c_{Camellin}}$, $P_{c_{Savini}}$, and $P_{c_{Jarade}}$ methods are based on a variation of the keratometric refractive index depending on the induced refractive change (ΔSE). This keratometric approach for corneal power estimation is more appropriate than the use of a single value of keratometric index in all cases, but it is still associated with some level of limitation, as has been shown in the current study. A specific keratometric index value for each cornea must be calculated if the keratometric approach is intended to be used for an estimation of corneal power, as demonstrated by our research group in a previous study [13]. This keratometric index was named as exact keratometric index ($n_{k_{exact}}$) and its calculation requires the measurement of anterior and posterior corneal curvature [13]. As devices measuring the posterior corneal curvature are not available in all clinical settings, our group developed the concept of adjusted keratometric index ($n_{k_{adj}}$), which is a clinically valid method for estimating an appropriate keratometric index allowing calculation of the corneal power with enough accuracy. This method only requires the measurement of the anterior corneal radius postoperatively, which can be easily obtained in any clinical setting.

Among methods not requiring previous historical clinical data, the methods evaluated in the current study led to overestimations of more than 1 D except for $P_{c_{HaigisL}}$, as commented previously. These findings confirm the relevance of the curvature of the posterior corneal surface in the error associated with the corneal power estimation when the keratometric approach is used. This is consistent with previous scientific evidence remarking that there is an error

in considering the $k = r_{2c}/r_{1c}$ ratio as a constant when corneal power is estimated in eyes after refractive surgery using the keratometric approach [15]. This is also the reason explaining the inaccuracy of methods of corneal power estimation based on considering the refractive change induced. Our results also confirm that the use of an inappropriate method for corneal power estimation in eyes with previous laser refractive surgery mainly leads to an overestimation of corneal power and consequently to an underestimation of the IOL power required when cataract surgery is planned in this type of cases, resulting in hyperopic residual refractive errors postoperatively.

In conclusion, the use of the adjusted keratometric index ($n_{k_{adj}}$) is a valid and easy method to estimate the central corneal power in corneas with previous myopic laser refractive surgery, improving the accuracy of methods described previously for such purpose. Only $P_{c_{HaigisL}}$ has been found to be comparable to our method and therefore leading to differences compared to the Gaussian corneal power which are clinically acceptable. The advantage of the use of $P_{k_{adj}}$ method is that the estimation of corneal power can be performed with the only requirement of measuring the postoperative anterior corneal radius and with an associated error within ± 0.7 D and only 6% of cases showing differences with the Gaussian power out of the range of ± 0.5 D.

5. Conclusion

With this paper, a new method for calculating corneal power after myopic refractive surgery was validated clinically, with the advantage that only postoperative refractive surgery parameters are required.

Disclosure

All the authors have full control of all primary data and they agree to allow Journal of Cataract and Refractive Surgery to review the data of the current study if requested.

Conflict of Interests

The authors have no financial or proprietary interest in a product, method, or material described herein.

Acknowledgment

This study has been funded by the Project no. GRE12-33 of the University of Alicante.

References

- [1] J. T. Holladay, "Standardizing constants for ultrasonic biometry, keratometry, and intraocular lens power calculations," *Journal of Cataract and Refractive Surgery*, vol. 23, no. 9, pp. 1356–1370, 1997.
- [2] A. M. Hamed, L. Wang, M. Misra, and D. D. Koch, "A comparative analysis of five methods of determining corneal refractive power in eyes that have undergone myopic laser in situ keratomileusis," *Ophthalmology*, vol. 109, no. 4, pp. 651–658, 2002.

- [3] S. T. Awwad, C. Manasseh, R. W. Bowman et al., "Intraocular lens power calculation after myopic laser in situ keratomileusis: estimating the corneal refractive power," *Journal of Cataract and Refractive Surgery*, vol. 34, no. 7, pp. 1070–1076, 2008.
- [4] M. Camellin and A. Calossi, "A new formula for intraocular lens power calculation after refractive corneal surgery," *Journal of Refractive Surgery*, vol. 22, no. 2, pp. 187–199, 2006.
- [5] G. Savini, M. Carbonelli, P. Barboni, and K. J. Hoffer, "Clinical relevance of radius of curvature error in corneal power measurements after excimer laser surgery," *Journal of Cataract & Refractive Surgery*, vol. 36, no. 1, pp. 82–86, 2010.
- [6] G. Savini, K. J. Hoffer, M. Carbonelli, and P. Barboni, "Intraocular lens power calculation after myopic excimer laser surgery: clinical comparison of published methods," *Journal of Cataract and Refractive Surgery*, vol. 36, no. 9, pp. 1455–1465, 2010.
- [7] E. F. Jarade and K. F. Tabbara, "New formula for calculating intraocular lens power after laser in situ keratomileusis," *Journal of Cataract and Refractive Surgery*, vol. 30, no. 8, pp. 1711–1715, 2004.
- [8] H. J. Shammas, M. C. Shammas, A. Garabet, J. H. Kim, A. Shammas, and L. Labree, "Correcting the corneal power measurements for intraocular lens power calculations after myopic laser in situ keratomileusis," *American Journal of Ophthalmology*, vol. 136, no. 3, pp. 426–432, 2003.
- [9] L. Wang, M. A. Booth, and D. D. Koch, "Comparison of intraocular lens power calculation methods in eyes that have undergone laser-assisted in-situ keratomileusis," *Transactions of the American Ophthalmological Society*, vol. 102, pp. 189–197, 2004.
- [10] G. Ferrara, G. Cennamo, G. Marotta, and E. Loffredo, "New formula to calculate corneal power after refractive surgery," *Journal of Refractive Surgery*, vol. 20, no. 5, pp. 465–471, 2004.
- [11] W. Haigis, "Intraocular lens calculation after refractive surgery for myopia: Haigis-L formula," *Journal of Cataract and Refractive Surgery*, vol. 34, no. 10, pp. 1658–1663, 2008.
- [12] V. J. Camps, D. P. P. Llorens, D. de Fez et al., "Algorithm for correcting the keratometric estimation error in normal eyes," *Optometry and Vision Science*, vol. 89, no. 2, pp. 221–228, 2012.
- [13] V. J. Camps, D. P. Piñero, V. Mateo et al., "Algorithm for correcting the keratometric error in the estimation of the corneal power in eyes with previous myopic laser refractive surgery," *Cornea*, vol. 32, no. 11, pp. 1454–1459, 2013.
- [14] J. M. Bland and D. G. Altman, "Statistical methods for assessing agreement between two methods of clinical measurement," *The Lancet*, vol. 1, no. 8476, pp. 307–310, 1986.
- [15] C. Edmund, "Posterior corneal curvature and its influence on corneal dioptric power," *Acta Ophthalmologica*, vol. 72, no. 6, pp. 715–720, 1994.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

