

## Research Article

# Prenatal and Postnatal Polycyclic Aromatic Hydrocarbon Exposure, Airway Hyperreactivity, and Beta-2 Adrenergic Receptor Function in Sensitized Mouse Offspring

Sophie Chu,<sup>1</sup> Hanjie Zhang,<sup>1</sup> Christina Maher,<sup>1</sup> Jacob D. McDonald,<sup>2</sup>  
Xiang Zhang,<sup>3</sup> Shuk-Mei Ho,<sup>3</sup> Beizhan Yan,<sup>4</sup> Steven Chillrud,<sup>4</sup> Frederica Perera,<sup>5</sup>  
Phillip Factor,<sup>6</sup> and Rachel L. Miller<sup>1,5,7</sup>

<sup>1</sup> Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, Columbia University College of Physicians and Surgeons, PH8E-101B, 630 W. 168th Street, New York, NY 10032, USA

<sup>2</sup> Department of Toxicology, Lovelace Respiratory Research Institute, Albuquerque, NM 87101, USA

<sup>3</sup> Department of Environmental Health Sciences, University of Cincinnati College of Medicine, Cincinnati, OH 45201, USA

<sup>4</sup> Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA

<sup>5</sup> Department of Environmental Health, Columbia University Mailman School of Public Health, New York, NY 10032, USA

<sup>6</sup> Department of Medicine, University of Arizona, Tucson, AZ, 85721, USA

<sup>7</sup> Division of Pediatric Allergy and Immunology, Department of Pediatrics, Columbia University College of Physicians and Surgeons, New York, NY 10032, USA

Correspondence should be addressed to Rachel L. Miller; [rlm14@cumc.columbia.edu](mailto:rlm14@cumc.columbia.edu)

Received 10 September 2013; Accepted 8 November 2013

Academic Editor: Lucio Guido Costa

Copyright © 2013 Sophie Chu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Despite data associating exposure to traffic-related polycyclic aromatic hydrocarbons (PAH) in asthma, mechanistic support has been limited. We hypothesized that both prenatal and early postnatal exposure to PAH would increase airway hyperreactivity (AHR) and that the resulting AHR may be insensitive to treatment with a  $\beta_2$ AR agonist drug, procaterol. Further, we hypothesized that these exposures would be associated with altered  $\beta_2$ AR gene expression and DNA methylation in mouse lungs. Mice were exposed prenatally or postnatally to a nebulized PAH mixture versus negative control aerosol 5 days a week. Double knockout  $\beta_2$ AR mice were exposed postnatally only. Prenatal exposure to PAH was associated with reduced  $\beta_2$ AR gene expression among nonsensitized mice offspring, but not increases in DNA methylation or AHR. Postnatal exposure to PAH was borderline associated with increased AHR among sensitized wildtype, but not knockout mice. In the first study that delivers PAH aerosols to mice in a relatively physiological manner, small effects on AHR and  $\beta_2$ AR gene expression, but not  $\beta_2$ AR agonist drug activity, were observed. If confirmed, the results may suggest that exposure to PAH, common ambient urban pollutants, affects  $\beta_2$ AR function, although the impact on the efficacy of  $\beta_2$ AR agonist drugs used in treating asthma remains uncertain.

## 1. Introduction

Exposure to traffic-related air pollution has been associated with exacerbations of respiratory symptoms, decreased lung function, and the development of asthma [1–5]. Incomplete combustion of diesel exhaust particles emitted by motor vehicle engines produces a complex mixture of pollutants that includes significant concentrations of polycyclic aromatic hydrocarbons (PAH). Previously, our group at the Columbia

Center for Children's Environmental Health (CCCEH) and others have shown that exposure to PAH was associated with asthma in children [1, 4, 6, 7]. Notably, the prenatal time window of exposure to PAH has been implicated in the development of childhood asthma, particularly in the presence of exposure to secondhand smoke [4, 8]. Also, repeated prenatal and early childhood exposure to pyrene, the predominant PAH in the NYC CCCEH local environment, has been associated with asthma regardless of exposure to

secondhand smoke or seroatopy in the CCCEH cohort [1]. Despite the emergence of data associating exposure of PAH to asthma-related outcomes, mechanistic support has been limited.

Inhaled  $\beta_2$ -adrenergic agonists are common treatments for reactive airway diseases and used for short-term and long-term alleviation of bronchoconstriction [9]. They are believed to function by binding to the active site of  $\beta_2$ ARs on airway epithelial and smooth muscle cells, leading to bronchodilation via activation of adenylyl cyclase, generation of intracellular cAMP, and the associated signaling events [10]. Despite their wide usage, inhaled  $\beta_2$ AR agonists have been associated with serious asthma-related complications. Specifically, the extended use of short-acting agonists has been associated with a loss of their protective bronchodilatory action and severe asthma exacerbations [11]. Certain populations, like children, African Americans, and those with particular  $\beta_2$ AR genotypes (e.g., homozygous for arginine at  $\beta_2$ AR-16 [Arg/Arg]), appear more susceptible to the morbidity and even mortality associated with long-term  $\beta_2$  agonist use [12–14]. Mechanisms proposed for this increase in morbidity and loss of efficacy have included suppression of symptoms associated with early but not more advanced inflammation and tolerance to the bronchodilatory activity, possibly via receptor desensitization [11].

More recent evidence suggests that  $\beta_2$ AR function and signaling could be impaired by exposure to PAH from ambient urban air. Specifically, we produced a PAH mixture that mimicked the proportional distribution of individual PAH observed in NYC air and found that exposure to the PAH mixture reduced the expression and function of  $\beta_2$ AR in primary human and mouse cell systems [15]. This observation provided the first evidence that environmentally relevant concentrations of PAH can impede  $\beta_2$ AR-mediated airway relaxation. Another study indicated that the exposure of adipocytes to PAHs also impaired the function of  $\beta_2$ AR, without reducing membrane-bound receptor numbers [16].

The primary objective of this study was to determine if exposure to traffic-related PAH alters airway  $\beta_2$ AR function following in utero and early life exposures *in vivo*. We hypothesized that both prenatal and early postnatal exposure to PAH would increase AHR in ovalbumin-sensitized offspring mice and that this AHR may not improve following administration of a  $\beta_2$ AR agonist drug. Further, we hypothesized that these exposures would be associated with altered  $\beta_2$ AR gene expression and DNA methylation in mouse lungs. The key to our approach was the administration of an aerosol of PAH with levels that mimicked the proportional distribution of individual PAH observed in NYC air inhaled by pregnant women [8, 17] that was delivered in a manner more physiological than reported in previous murine studies [18].

## 2. Materials and Methods

**2.1. Animals.** Seven-week-old C57/Bl6 mice were obtained from Charles River. The mice were housed in a conventional animal facility in a private room under a 12 hour light/dark cycle with free access to food and water. Double knockout

TABLE 1: Components of PAH aerosol.

PAH	Proportion (%)	Mean (ng/m <sup>3</sup> )
Benzo[a]anthracene	3.99	0.27
Benzo[a]pyrene	5.13	0.42
Benzo[b]fluoranthene	7.62	0.59
Benzo[k]fluoranthene	1.79	0.15
Benzo[g,h,i]perylene	13.02	1.12
Chrysene	4.82	0.35
Dibenzo[a,h]anthracene	0.89	0.06
Indeno[1,2,3-cd]pyrene	7.42	0.64
Pyrene	52.59	3.69

$\beta_2$ AR DKO  $\beta_2$ AR<sup>tm1Bkk</sup>  $\beta_2$ AR<sup>tm1Bkk</sup>/J mice (<http://jaxmice.jax.org/strain/003810.html>) with complete absence of  $\beta_2$ AR function (null for adrenergic receptor 1 and 2 genes) were obtained from Jackson laboratory. Published data showed that these mice do not respond to administration of  $\beta$  agonist drugs [19]. Mice were mated at 8 weeks old. These mice were housed and fed ad lib breeder chow (Labdiet, St. Louis, MO) starting the first day of mating until postnatal day (PND) 28 when it was replaced with regular chow. The experimental protocol was approved by the IACUC.

**2.2. PAH Exposure.** The PAH mixture was produced by the Lovelace Respiratory Research Institute to replicate the proportional distribution of individual PAH that was measured among a cohort of pregnant women participating at the CCCEH birth cohort [8, 17] as shown in Table 1. The negative control aerosol solution consisted of 99.97% purified water, 0.02% Tween 80, and 0.01% Antifoam A (Sigma-Aldrich, St. Louis, MO). 100  $\mu$ L of PAH solution was added to yield the final concentration of 7.29 ng/m<sup>3</sup>. The aerosols were tested for bacterial growth annually using aerobic and anaerobic cultures (Charles River, MA). Fresh solutions were prepared weekly.

Mice were exposed prenatally or postnatally to PAH versus negative control aerosol, as summarized in Figure 1. The 15 mL solutions were delivered via nebulizers (Unomedical Inc., McAllen, TX). Filtered compressed air was connected to the nebulizer. Prenatal aerosol was delivered beginning from gestational days (GD) 1–3 (the Monday after mating) until GD 19–21 or day of delivery. Postnatal aerosol was administered beginning from postnatal day (PND) 2 until PND 19–21.

Two cages without filter tops were placed in each (PAH, normal air) exposure chamber for five hours a day, five days a week, as shown in Figure 2. The exposure chambers were set to achieve a flow of 12.5 to 13.0 liters per minute (LPM). Pressure gauges on the panel were set to 20 psi. At the beginning of each week of the exposure, a 25 mm Pall Flex filter (Pall Lifesciences, Port Washington, NY) with an Amberlite XAD-4 coating was combined with a noncoated filter in front to provide support to the brittle XAD coated filter. The filters were replenished weekly and stored in –20 F freezer until analyzed at Lamont-Doherty Earth Observatory, Columbia University. Three weekly filters with PAHs were extracted together by mixture of CH<sub>2</sub>Cl<sub>2</sub> and CH<sub>3</sub>OH (9 : 1 v : v) 3 times




FIGURE 1: Algorithm of experimental design. Mice were exposed to PAH or normal air either on GD 3 to 19–21 or PND 2 to 19–21. Offspring were sensitized to either OVA or PBS on PND 24 and then again on PND 31. The mice were OVA challenged on PNDs 38–40. Airway resistance was measured and biospecimens were collected on PND 41.




FIGURE 2: PAH chamber. Mice were exposed to PAH or normal air in two side-by-side chambers delivered via aerosol (only one shown). Filters were collected as depicted and used to measure ambient concentrations.

under sonication and then concentrated under the gentle flow of  $N_2$ . Levels of PAHs, including pyrene which was used as an indicator of PAH aerosolization, were measured by a Varian (now Agilent) 1200L gas chromatography-tandem mass spectrometer (GC/MS/MS).

**2.3. Sensitization to Ovalbumin (OVA) and Measurement of OVA-IgE.** Pups were sensitized to either ovalbumin (OVA) or phosphate buffered saline (PBS) as a negative control beginning from PND 24 using a 0.1 mg solution of Grade V chicken egg albumin (Sigma-Aldrich, St. Louis, MO, A5503-10G) in water and 4 mg Imject alum (Pierce, Rockford, IL, number 77161) intraperitoneal (i.p., 200  $\mu$ L). The mice were redosed one week later. All the mice were then challenged with aerosolized OVA (3% OVA in sterile PBS) for 30 minutes on days 38, 39, and 40.

Anti-OVA-immunoglobulin E (OVA-IgE) was measured by enzyme-linked immune sorbent assay (ELISA) in all available blood specimens using the Mouse Serum Anti-OVA IgE Antibody Assay Kit (Chondrex, Redmond, WA) according to the manufacturer's protocol. Sera were diluted from 1:5 to 1:10. The data were analyzed using a log-log model equation (<http://readerfit.com/>).

**2.4. Measurement of AHR.** Half of the mice were pre-treated with 3 mM of the  $\beta_2$ -adrenergic agonist procaterol (Sigma-Aldrich, St. Louis, MO) by aerosol for 30 minutes immediately prior to AHR measurement using a smaller chamber connected to a nebulizer. Airway reactivity was measured between PND 41 and 45 of age as shown in Figure 1. Airway resistance was measured using a computer controlled, small animal ventilator (SAV, SCIREQ flexiVent,

Montreal, QC, Canada). Mice were sedated (pentobarbital, 30–40 mg/kg i.p.), tracheotomized with an 18 g × 1 cm metal cannula, paralyzed (pancuronium, 50 mg/kg i.p.), and ventilated using a quasisinusoidal ventilatory mode at 150 breaths per minute (bpm) with 3 cm H<sub>2</sub>O of positive end expiratory pressure. Body temperature was maintained with a water-perfused heating pad and cardiac rhythm was monitored continuously using a SCIREQ EKG. Lung volume was then standardized twice using a constant flow, pressure-limited mode of ventilation (tidal volume = 10 mL/kg, pressure at airway opening limited to 30 cm-H<sub>2</sub>O at 150 bpm), and occlusion of the expiratory port of the ventilator for 10 breaths. Mice were then allowed to equilibrate for 5 minutes during which they were ventilated with a quasisinusoidal pattern. Following equilibration, PBS was aerosolized using a SCIREQ zero-dead space ultrasonic nebulizer that was positioned in the inspiratory limb of the ventilator circuit. Airway resistance was measured using an 8-second forced oscillation, composed of nonoverlapping, nonharmonic frequencies that were used to produce an index of central airways resistance,  $R_n$  (Newtonian resistance). Incremental doses of methacholine (8, 16, 32, and 64 mg/mL in PBS) were then aerosolized. For each animal, peak airway resistance following each dose of methacholine was normalized to  $R_n$  following PBS and plotted. All measures were graded A, B, C, or D according to prespecified criteria: (A) an ideal curve exhibiting a strong initial peak of resistance and then a return to baseline following each dose of methacholine, (B) as in (A), but one methacholine doses without clearly defined peak, (C) as in (B), but more than methacholine doses without clearly defined peak, or many exclusions noted by the software, or (D) flat  $R_n$  or loss of cardiac rhythm during procedure. Only data graded A–C were used. At the end of the procedure, all mice were euthanized by cardiac puncture. The lungs and spleen were flash-frozen and stored in –80°C freezer until further analysis. The serum was isolated from the blood obtained by the cardiac puncture and stored in –80°C freezer until IgE testing.

**2.5. Bronchoalveolar Lavage (BAL).** A subset of mice ( $n = 42$ ) distributed across each experimental group was chosen randomly for BAL instead of AHR measurement. The BAL fluid was centrifuged at 4°C at 1500 rpm for 5 minutes. Cell pellets were resuspended in 1 mL of PBS. Slides were prepared using a cytocentrifuge (Cytospin; Shandon GMI, Ramsey, MN) at 500 rpm for 5 minutes and then stained with Wright-Giemsa stain (Sigma-Aldrich, St. Louis, MO). Total 100 cells were counted for each sample from 10 randomly chosen viewing fields and total eosinophil, lymphocyte, macrophage, and neutrophil counts were quantified by a masked reader.

**2.6.  $\beta_2$ AR Gene Expression.** Tissues for gene expression were derived from prenatal and postnatal PAH exposed mice across the four experimental groups. The prenatal specimens were derived from either the nonperfused lung that was ligated during the BAL ( $n = 13$ ) or from lungs of mice that did not undergo AHR or BAL ( $n = 25$ ) and were preserved in RNAlater (Ambion, Austin, TX) in the presence of Trizol

reagent (Invitrogen, Carlsbad, CA). The postnatal specimens were derived from the nonperfused lung that was ligated during BAL ( $n = 22$ ). To account for possible systemic changes in gene expression, corresponding spleens were collected from the same animals. Lungs and splenic DNA samples were pooled first by experimental group, and then the gene expression experiments in the lungs were repeated in individual samples. Precellys 24 homogenizer (Bertin Technologies, France) was used to homogenize mouse lung and spleen tissues at 5000 rpm for 15 sec and repeated twice. The RNA was extracted using SuperScript III Reverse Transcriptase (Invitrogen, Grand Island, NY) according to the manufacturer's (Invitrogen, Grand Island, NY) suggested protocol. Primers for real-time RT-PCR were designed by using Primer-BLAST program (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). The sequences of  $\beta_2$ AR forward and reverse primers are 5'-CTT GCT GGC ACC CAA CGG AA-3' and 5'-ATG CCC ACA ACC CAC GCT TC-3', respectively, with the amplicon size 78 bp. Mouse  $\beta_2$  microglobulin ( $\beta_2m$ ) was used as the housekeeping gene. The sequences of  $\beta_2m$  forward and reverse primers are 5'-GCC TGT ATG CTA TCC AGA AAA CCC C-3' and 5'-TGA GGC GGG TGG AAC TGT GT-3', respectively, with the amplicon size 112 bp. Real-time RT-PCR was performed in a 7900HT Fast Real-Time PCR System (ABI, Foster City, CA) under standard mode. The RT-PCR was performed in triplicate by using three  $\mu$ L 1:10 diluted cDNA in a total of 10  $\mu$ L reaction. The  $2^{-\Delta\Delta C_t}$  method was used to calculate the relative expression level of  $\beta_2$ AR transcripts as we performed previously [20].

**2.7.  $\beta_2$ AR DNA Methylation.** Lungs and splenic DNA samples were pooled by experimental group. Genomic DNA was extracted from mouse lung and splenic tissue by DNeasy Blood & Tissue kit (Qiagen, Valencia, CA) in the presence of RNase A. EZ DNA Methylation kit (Zymo Research, Irvine, CA) was used to bisulfite convert 500 ng of genomic DNA in each reaction and then eluted with 40  $\mu$ L buffer.  $\beta_2$ AR gene promoter flanking transcription start site was selected for the methylation analysis. Methprimer [21] was used to design primers for bisulfite PCR. The forward and reverse primers for  $\beta_2$ AR bisulfite PCR are 5'-TTG GGT AAT TTT TTT AAA GTT TGG T-3' and 5'-ACA TAA AAA TAA CCA TAC CCA CAA C-3', respectively. The 390 bp amplicon (–40 tp +350) covers 32 CpG sites in the CpG island located in the region from +1 to +473.

PCR was performed by adding 2  $\mu$ L bisulfite modified DNA as template in a 25  $\mu$ L reaction. Platinum Taq DNA polymerase (Invitrogen, Carlsbad, CA) was added to the reaction according to manufacturer's instructions. The PCR together with the negative control were performed for 40 cycles with the annealing temperature set at 55°C. Gel-purified amplicons were then TA-cloned into pGEM T easy vector (Promega, Madison, WI). The plasmids in *E. coli* were directly amplified by TempliPhi DNA amplification kits (GE Healthcare, Indianapolis, IN) and submitted for sequencing (Macrogen USA, Rockville, MD). For each sample, a total of 12 clones were picked. The methylation status in each CpG site was analyzed by BiQ Analyzer [22].

TABLE 2: Pyrene concentrations (ng/m<sup>3</sup>) in exposure chamber.

Exposure	N	Mean ± SE	Range
Normal air	11	3.03 ± 0.90	0–9.00
PAH	11	23.24 ± 3.05	7.38–40.00

N represents the number of extractions of 3 filters, each one collecting pyrene levels over a 5-day period of aerosolization. The single level obtained from 3 consecutive filters was designed to capture the average level over a single exposure period. 11 filters were chosen from 13 rounds of mice representing both exposure groups evenly. Measured pyrene concentration in the PAH exposure group averaged higher than that in normal air exposure group ( $P < 0.005$ , two-tailed  $t$ -test).

**2.8. Statistics.** Differences between experimental groups (levels of pyrene collected on filters, Newtonian resistance) were assessed by Student's  $t$ -test and means and standard error values displayed. One-way ANOVA was used to analyze the AHR data. Generalized estimating equations (GEE) were used to evaluate  $R_n$  by experimental condition across all methacholine doses. Two-tailed  $P$  values  $< 0.05$  were considered statistically significant in all models.

### 3. Results

**3.1. PAH Were Well-Aerosolized in Exposure Chamber.** In order to verify proper aerosolization of the PAH in the exposure chamber, particulate fractions of the ambient PAH versus negative control aerosol were measured from the filters over 5-day periods and then averaged. Measured pyrene concentration in the PAH exposure group averaged higher than those in the negative control group ( $23.24 \pm 3.05$  versus  $3.03 \pm 0.90$ ,  $P < 0.005$ ) (Table 2). The experimental PAH concentration was on average 33-fold higher than the background PAH concentration in the animal housing facility (see Supplemental Table 1 available online at <http://dx.doi.org/10.1155/2013/603581>), documenting aerosolization of the PAH during experimental procedures. Time weighted concentrations for the experimentally exposed mice were approximately 74-fold higher than levels measured in the housing facility.

**3.2. OVA-IgE and Airway Eosinophilia.** In order to determine whether the mice were properly sensitized to OVA, anti-OVA IgE levels were measured from serum collected prior to sensitization and 10 days following the last i.p. dose. OVA sensitized mice had higher fold increases than those of the PBS sensitized mice ( $12.69 \pm 2.71$ ,  $1.69 \pm 0.29$ , mean ± SE for OVA, PBS control, resp.,  $P = 0.003$ ).

In the limited number of mice sampled ( $n = 19$ , selected randomly from 4 rounds of exposure), a greater percent of eosinophils were detected among the OVA sensitized mice when compared to the nonsensitized mice ( $10\% + 1.14$  versus  $4\% + 0.90$ ,  $P = 0.003$ ). There was no difference between cell numbers across all experimental groups of the postnatal exposed mice (data not shown).

**3.3. Effects of Prenatal PAH Exposure in Wildtype Mice.** Airway reactivity data were analyzed by normalizing Newtonian




FIGURE 3: AHR following prenatal PAH exposure in the presence and absence of procaterol (Proc). Newtonian resistance ( $H_2O/s/mL$ ) was normalized to value measured following PBS aerosol exposure. Total sample size, followed by sample size for mice that completed highest dose, is listed in parentheses.

resistance ( $H_2O/s/mL$ ) to the value measured following PBS aerosol exposure. Significant differences were not detected across any of the experimental groups ( $P = NS$ , ANOVA) (Figure 3). Experimental groups did not exhibit improvement in AHR following administration of procaterol ( $P = NS$ ).

**3.4. Effects of Early Postnatal PAH Exposure in Wildtype Mice.** PAH/OVA mice exhibited borderline higher airway resistance when compared to the negative control aerosol/OVA exposure group ( $P = 0.055$ , two-tailed  $t$ -test) (Figure 4). This modest finding was confirmed in GEEs that showed a highly significant  $\beta$  value among the PAH/OVA that exceeded the value calculated from the other experimental groups (Supplemental Table 2). There was no significant resolution of AHR among mice following administration of procaterol, regardless of PAH exposure ( $P = NS$ ).

**3.5. Effects of Early Postnatal PAH Exposure Airway Results in  $\beta_2AR^{-/-}$  Double Knockout (DKO) Mice.** Significant differences in AHR were not detected across experimental groups in  $\beta_2AR^{-/-}$  DKO mice ( $P = NS$ , ANOVA) (Figure 5). Further, significant improvements were not detected in any of the experiment groups following pretreatment with procaterol ( $P = NS$ ).

**3.6.  $\beta_2AR$  Gene Expression and DNA Methylation.** Finally, we sought to determine whether  $\beta_2AR$  molecular function was impaired, even in the absence of detectable impairment of  $\beta_2AR$  agonist physiological activity (i.e., improvement




FIGURE 4: AHR following postnatal PAH exposure in the presence and absence of procaterol (Proc). Newtonian resistance ( $H_2O/s/mL$ ) was normalized to value measured following PBS aerosol exposure. Total sample size, followed by sample size for mice that completed highest dose, is listed in parentheses. The PAH/OVA mice showed borderline greater airway resistance when compared to the normal air/OVA group at the 64 mg/mL dose ( $P$  value = 0.055, two-tailed  $t$ -test). Clear improvement across experimental groups following administration of procaterol is not evident.

in AHR) with administration of procaterol. This question was examined through determination of whether aerosolized PAH altered  $\beta_2AR$  gene expression and/or DNA methylation. We found that mice prenatally exposed to PAH (regardless of sensitization) exhibited borderline lower  $\beta_2AR$  gene expression in the lung, but not spleen, when compared to the negative control aerosol mice (i.e.,  $P = 0.054$ ) (Table 3). Differences may have been more pronounced among the non-sensitized mice where the PAH/PBS mice showed lower  $\beta_2AR$  gene expression than that of the negative control aerosol/PBS mice ( $P = 0.048$ ) (Table 3). There were no other differences in gene expression between groups of prenatal exposed mice ( $P = NS$ ). Also, no differences across experimental groups following postnatal exposure to PAH were detected ( $P = NS$ ). There were no differential effects on DNA methylation in the lungs or spleen (unmethylated in all groups, data not shown).

#### 4. Discussion

Using a novel chamber to determine the effects of a more physiological method of mouse exposure to ambient PAH, we found that prenatal exposure to PAH was associated with reduced  $\beta_2AR$  gene expression in the mouse lung, and postnatal exposure to PAH was borderline associated with increased AHR. While these effects were modest or borderline, they may suggest a possible novel mechanism




FIGURE 5: AHR following postnatal PAH exposure in the presence and absence of procaterol in  $\beta_2AR^{(-/-)}$  DKO mice. There were no significant differences between the OVA sensitized/OVA challenged and the PBS sensitized/OVA challenged mice in either PAH or normal air exposure ( $P = NS$ ). Clear improvement across experimental groups following administration of procaterol is not evident.

TABLE 3:  $\beta_2AR$  gene expression in lungs.

	Mean $\pm$ SE		$P$ value
PAH versus normal air	0.65 $\pm$ 0.05 ( $N = 15$ )	0.91 $\pm$ 0.11 ( $N = 18$ )	0.054
PAH/PBS versus normal air/PBS	0.55 $\pm$ 0.06 ( $N = 7$ )	1.00 $\pm$ 0.18 ( $N = 9$ )	0.048
PAH/OVA versus normal air/OVA	0.75 $\pm$ 0.07 ( $N = 8$ )	0.83 $\pm$ 0.13 ( $N = 9$ )	0.620

$N$  refers to the number of mice per group. Values were normalized by setting a mean expression level from the normal air/OVA group and comparing values with the other groups. PAH exposed mice exhibited lower  $\beta_2AR$  gene expression than that of normal air exposed mice, regardless of sensitization ( $P = 0.054$ , two-tailed  $t$ -test). Differences in gene expression appeared more pronounced among the nonsensitized mice ( $P = 0.048$ , two-tailed  $t$ -test).

for loss of  $\beta_2AR$  efficacy following ambient PAH exposure, a common urban pollutant. These results also extend our previous work in primary cells [15] and suggest the first *in vivo* demonstration of altered  $\beta_2AR$  function in the lung following air pollution exposure. One major strength of this study includes the innovative delivery of ambient PAH exposure using individual PAH concentrations proportionate to exposure in the NYC setting [15] and not yet reported in other studies [18]. Other strengths include the assessment of the effects of PAH during the prenatal versus postnatal time periods, documentation of aerosolization through measurement of levels extracted from filters placed in the chamber

and comparison with levels following nonexperimental (i.e., mouse housing) levels, and direct assessment of AHR.

The primary outcome of interest, namely, AHR, was only suggested following early (starting within 3 days of birth) postnatal PAH exposure. Clearly these results were of borderline statistical significance. In contrast, differences in AHR were not found following prenatal exposure, unlike previous human cohort studies [4, 6, 8]. We suspect that the effects of early postnatal PAH may mirror what we found in human cohorts where the combination of prenatal and early postnatal exposure to pyrene, the predominant local airborne PAH, was associated with childhood asthma and related symptoms [1]. Interesting, these results are the opposite of Singh and colleagues' findings following exposure of mice to secondhand smoke. In their model, prenatal, but not postnatal, secondhand smoke exposure increased methacholine-induced AHR in sensitized animals and upregulated allergic immune responses [23]. Fedulov and colleagues also showed that diesel exhaust particles during pregnancy induced greater AHR in the mice offspring [18].

It is not surprising that PAH-induced alterations in AHR may have been observed following postnatal exposure in mice, whereas PAH-induced asthma has been associated with prenatal exposure in some human cohorts [4]. In humans, lung alveolarization continues through age of 7 years, whereas, in the mouse, lung development starts prenatally and continues for several weeks after birth [18]. Hence, the period when rapidly proliferating cells may be most susceptible to toxic environmental exposures occurs during different times of exposure across species and may be later for mice. In addition, the mechanics of breathing differ between mouse and human. Small mammals, like mice, breathe faster [24]. Presumably higher ventilation rates may affect the dose of toxin inhaled.

Another possibility is that other toxicants or chemical compounds are responsible for the previously observed adverse associations following exposure to PAHs. These may include oxygenated PAH (that may increase with photooxidation processes when ambient ozone concentrations are high) and nitroPAHs that were not provided in the aerosol and exposure chamber. Reactive oxygen species may be present in ambient particulate matter or generated upon inhalation of PAHs in the lungs and can be converted to hydroxyl radicals that can induce mitochondrial damage and apoptosis [25, 26]. Also, several families of electrophilic metabolites (diol epoxides, radical cations, and reactive and redox active *o*-quinones) possess cytotoxic properties and have been implicated in PAH-induced carcinogenesis [24].

To ascertain whether the possible association with AHR was mediated by PAH-induced impairment of  $\beta_2$ AR function, we conducted parallel experiments in  $\beta_2$ AR KO mice that would not be expected to exhibit any  $\beta_2$ AR function. With this model, we were unable to implicate the  $\beta_2$ AR pathway as being causal. This result arguably was limited by the insufficient induction of AHR across the experimental groups, documented to occur in some  $\beta_2$ AR KO mouse models [27, 28], which also made the determination of procaterol-induced effects difficult. Also, the intensity of AHR can vary

by wildtype mouse strain, possibly attributable to strain-related differences in the expression of phospholipase C- $\beta_1$  [27], airway inflammation [25], or calcium sensitivity [19].

Despite the absence of an effect of prenatal exposure to PAH on AHR and sensitivity to  $\beta_2$  adrenergic agonist drugs, differences across exposure groups in  $\beta_2$ AR gene expression in the lung were evident. We had predicted that changes in gene expression could occur despite the absence of differences across groups in procaterol-induced resolution of AHR. This finding was more statistically sound when comparing the PAH versus negative control aerosol exposure groups among the nonsensitized mice, suggesting greater susceptibility to PAH exposure when allergic immune responses are not induced. Even though this finding occurred without downstream effects on airway bronchodilation as measured by AHR, the greater susceptibility in the absence of sensitization is consistent with our previous publication. In this work, exposure to pyrene was most associated with asthma and asthma-related symptoms among the nonatopic young children [1]. A similar pattern was observed in other human cohorts where exposure to secondhand smoke was associated with asthma symptoms in nonatopic children [29, 30]. Further, this finding is consistent with other examples in the literature that prenatal measures of several environmental exposures, including tobacco smoke, farming environment, and methyl-rich diet, affect asthma gene expression in the offspring [10, 31–33]. Epigenetic regulation of  $\beta_2$ AR gene expression in T helper 1 and 2 cells has been detected previously [33]. However, the differences in our experiments were not attributable to alterations in DNA methylation, suggesting that an alternative method influenced gene expression. These effects also appeared end organ specific, as they were not detected in the spleen.

We acknowledge several limitations. Differences across mouse strains were not tested, limiting the interpretation of the results from KO mice and comparisons across groups. The small sample sizes may have underpowered us to detect small differences across experimental groups, perhaps explaining the weak results following administration of procaterol. The experimental conditions may have been insufficient to induce more physiological or molecular effects. Finally, we were unable to compare AHR measures with gene expression measures because different mice necessarily were used to avoid the effects of AHR-related procedures on gene expression.

In conclusion, this is the first study to deliver PAH aerosols to mice in a relatively physiological manner and measure their effects on airway and molecular outcomes. This may be the first *in vivo* demonstration that a mixture of common traffic-related air pollutants may impair  $\beta_2$ AR function on some level prenatally and postnatally, as the results begin to suggest that postnatal PAH exposure may induce AHR. Although the reduction in  $\beta_2$ AR gene expression that was present following prenatal PAH exposure, particularly in the absence of OVA sensitization, was not associated with altered AHR or loss of reversibility after administration of a  $\beta_2$  agonist, hence, its clinical significance is unclear. If the molecular effects are confirmed and associated with other outcomes in future studies, they may help drive a novel paradigm for urban asthma morbidity.

## Authors' Contribution

Sophie Chu and Hanjie Zhang contributed equally in the preparation of the paper.

## Acknowledgment

This work was supported by National Institute of Environmental Health Sciences at the National Institutes of Health (Grant no. P50ES015905).

## References

- [1] K. H. Jung, B. Yan, S. I. Hsu et al., "Prenatal and early postnatal exposure to pyrene and nonvolatile polycyclic aromatic hydrocarbons and asthma at ages 5-6 yrs: effect of seroatopy," *Annals of Allergy, Asthma and Immunology*, vol. 109, no. 4, pp. 249-254, 2012.
- [2] J. J. Kim, S. Smorodinsky, M. Lipsett, B. C. Singer, A. T. Hodgson, and B. Ostro, "Traffic-related air pollution near busy roads: the East Bay Children's Respiratory Health Study," *American Journal of Respiratory and Critical Care Medicine*, vol. 170, no. 5, pp. 520-526, 2004.
- [3] M. M. Patel and R. L. Miller, "Air pollution and childhood asthma: recent advances and future directions," *Current Opinion in Pediatrics*, vol. 21, no. 2, pp. 235-242, 2009.
- [4] M. J. Rosa, K. H. Jung, M. S. Perzanowski et al., "Prenatal exposure to polycyclic aromatic hydrocarbons, environmental tobacco smoke and asthma," *Respiratory Medicine*, vol. 105, no. 6, pp. 869-876, 2011.
- [5] A. J. Venn, S. A. Lewis, M. Cooper, R. Hubbard, and J. Britton, "Living near a main road and the risk of wheezing illness in children," *American Journal of Respiratory and Critical Care Medicine*, vol. 164, no. 12, pp. 2177-2180, 2002.
- [6] W. Jedrychowski, A. Galas, A. Pac et al., "Prenatal ambient air exposure to polycyclic aromatic hydrocarbons and the occurrence of respiratory symptoms over the first year of life," *European Journal of Epidemiology*, vol. 20, no. 9, pp. 775-782, 2005.
- [7] K. Nadeau, C. McDonald-Hyman, E. M. Noth et al., "Ambient air pollution impairs regulatory T-cell function in asthma," *Journal of Allergy and Clinical Immunology*, vol. 126, no. 4, pp. 845-852, 2010.
- [8] R. L. Miller, R. Garfinkel, M. Horton et al., "Polycyclic aromatic hydrocarbons, environmental tobacco smoke, and respiratory symptoms in an inner-city birth cohort," *Chest*, vol. 126, no. 4, pp. 1071-1078, 2004.
- [9] National Heart, Lung, and Blood Institute. National Asthma Education and Prevention Program, Expert Panel Report 3: guidelines for the diagnosis and management of asthma, 2007.
- [10] M. Cazzola, C. O. Page, P. Rogliani, and M. G. Matera, "B2-agonist therapy in lung disease," *American Journal of Respiratory and Critical Care Medicine*, vol. 187, no. 7, pp. 690-696, 2013.
- [11] H. S. Nelson, "Is there a problem with inhaled long-acting  $\beta$ -adrenergic agonists?" *Journal of Allergy and Clinical Immunology*, vol. 117, no. 1, pp. 3-16, 2006.
- [12] G. A. Hawkins, K. Tantisira, D. A. Meyers et al., "Sequence, haplotype, and association analysis of ADR $\beta$ 2 in a multiethnic asthma case-control study," *American Journal of Respiratory and Critical Care Medicine*, vol. 174, no. 10, pp. 1101-1109, 2006.
- [13] E. Israel, J. M. Drazen, S. B. Liggett et al., "The effect of polymorphisms of the  $\beta$ 2-adrenergic receptor on the response to regular use of albuterol in asthma," *American Journal of Respiratory and Critical Care Medicine*, vol. 162, no. 1, pp. 75-80, 2000.
- [14] A. W. McMahon, M. S. Levenson, B. W. McEvoy, A. D. Mosholder, and D. Murphy, "Age and risks of FDA-approved long-acting  $\beta$ 2-adrenergic receptor agonists," *Pediatrics*, vol. 128, no. 5, pp. e1147-e1154, 2011.
- [15] P. Factor, A. T. Akhmedov, J. D. McDonald et al., "Polycyclic aromatic hydrocarbons impair function of  $\beta$ 2- adrenergic receptors in airway epithelial and smooth muscle cells," *The American Journal of Respiratory Cell and Molecular Biology*, vol. 45, no. 5, pp. 1045-1049, 2011.
- [16] P. Irigaray, V. Ogier, S. Jacquenet et al., "Benzo[a]pyrene impairs  $\beta$ -adrenergic stimulation of adipose tissue lipolysis and causes weight gain in mice: a novel molecular mechanism of toxicity for a common food pollutant," *FEBS Journal*, vol. 273, no. 7, pp. 1362-1372, 2006.
- [17] F. P. Perera, V. Rauh, W.-Y. Tsai et al., "Effects of transplacental exposure to environmental pollutants on birth outcomes in a multiethnic population," *Environmental Health Perspectives*, vol. 111, no. 2, pp. 201-205, 2003.
- [18] A. V. Fedulov, A. Leme, Z. Yang et al., "Pulmonary exposure to particles during pregnancy causes increased neonatal asthma susceptibility," *The American Journal of Respiratory Cell and Molecular Biology*, vol. 38, no. 1, pp. 57-67, 2008.
- [19] Y. Xiang, E. Devic, and B. Kobilka, "The PDZ binding motif of the  $\beta$ 1 adrenergic receptor modulates receptor trafficking and signaling in cardiac myocytes," *The Journal of Biological Chemistry*, vol. 277, no. 37, pp. 33783-33790, 2002.
- [20] X. Zhang, Y.-K. Leung, and S.-M. Ho, "AP-2 regulates the transcription of estrogen receptor (ER)- $\beta$  by acting through a methylation hotspot of the 0N promoter in prostate cancer cells," *Oncogene*, vol. 26, no. 52, pp. 7346-7354, 2007.
- [21] L.-C. Li, "Designing PCR primer for DNA methylation mapping," *Methods in Molecular Biology*, vol. 402, pp. 371-383, 2007.
- [22] C. Bock, S. Reither, T. Mikeska, M. Paulsen, J. Walter, and T. Lengauer, "BiQ Analyzer: visualization and quality control for DNA methylation data from bisulfite sequencing," *Bioinformatics*, vol. 21, no. 21, pp. 4067-4068, 2005.
- [23] S. P. Singh, S. Gundavarapu, J. C. Peña-Philippides et al., "Prenatal secondhand cigarette smoke promotes Th2 polarization and impairs goblet cell differentiation and airway mucus formation," *The Journal of Immunology*, vol. 187, no. 9, pp. 4542-4552, 2011.
- [24] T. M. Penning, M. E. Burczynski, C.-F. Hung, K. D. McCoull, N. T. Palackal, and L. S. Tsuruda, "Dihydrodiol dehydrogenases and polycyclic aromatic hydrocarbon activation: generation of reactive and redox active o-quinones," *Chemical Research in Toxicology*, vol. 12, no. 1, pp. 1-18, 1999.
- [25] M. Y. Chung, R. A. Lazaro, D. Lim et al., "Aerosol-borne quinones and reactive oxygen species generation by particulate matter extracts," *Environmental Science and Technology*, vol. 40, no. 16, pp. 4880-4886, 2006.
- [26] T. Xia, P. Korge, J. N. Weiss et al., "Quinones and aromatic chemical compounds in particulate matter induce mitochondrial dysfunction: implications for ultrafine particle toxicity," *Environmental Health Perspectives*, vol. 112, no. 14, pp. 1347-1358, 2004.
- [27] D. W. McGraw, K. F. Almoosa, R. J. Paul, B. K. Kobilka, and S. B. Liggett, "Antithetic regulation by  $\beta$ -adrenergic receptors of Gq

- receptor signaling via phospholipase C underlies the airway  $\beta$ -agonist paradox," *The Journal of Clinical Investigation*, vol. 112, no. 4, pp. 619–626, 2003.
- [28] L. P. Nguyen, R. Lin, S. Parra et al., " $\beta$ 2-adrenoceptor signaling is required for the development of an asthma phenotype in a murine model," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 7, pp. 2435–2440, 2009.
- [29] L. García-Marcos, J. A. Castro-Rodríguez, M. M. Suarez-Varela et al., "A different pattern of risk factors for atopic and non-atopic wheezing in 9–12-year-old children," *Pediatric Allergy and Immunology*, vol. 16, no. 6, pp. 471–477, 2005.
- [30] M. C. McCormack, P. N. Breyse, E. C. Matsui et al., "Indoor particulate matter increases asthma morbidity in children with non-atopic and atopic asthma," *Annals of Allergy, Asthma and Immunology*, vol. 106, no. 4, pp. 308–315, 2011.
- [31] M. J. Ege, C. Bieli, R. Frei et al., "Prenatal farm exposure is related to the expression of receptors of the innate immunity and to atopic sensitization in school-age children," *Journal of Allergy and Clinical Immunology*, vol. 117, no. 4, pp. 817–823, 2006.
- [32] J. W. Hollingsworth, S. Maruoka, K. Boon et al., "In utero supplementation with methyl donors enhances allergic airway disease in mice," *The Journal of Clinical Investigation*, vol. 118, no. 10, pp. 3462–3469, 2008.
- [33] J. W. McAlees, L. T. Smith, R. S. Erbe, D. Jarjoura, N. M. Ponzio, and V. M. Sanders, "Epigenetic regulation of beta2-adrenergic receptor expression in TH1 and TH2 cells," *Brain, Behavior, and Immunity*, vol. 25, no. 3, pp. 408–415, 2011.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

