


Research Article

Molecular Evidence of *Wolbachia* Species in Wild-Caught *Aedes albopictus* and *Aedes aegypti* Mosquitoes in Four States of Northeast India

Sathishkumar Vinayagam,^{1,2} Tulika Nirmolia,¹ Sumi Chetry,¹ Narendran Pradeep Kumar,³ Prasanta Saini,³ Dibya Ranjan Bhattacharyya,¹ Ipsita Pal Bhowmick,¹ Kamaraj Sattu,² and Saurav Jyoti Patgiri ¹

¹ICMR-Regional Medical Research Centre North East Region, Dibrugarh, Assam 786010, India

²Periyar University, Centre for PG & Research Studies, Dharmapuri 635205, India

³ICMR-Vector Control Research Centre, Puducherry 605006, India

Correspondence should be addressed to Saurav Jyoti Patgiri; saurav.patgiri@gmail.com

Received 31 May 2023; Revised 17 August 2023; Accepted 25 August 2023; Published 5 September 2023

Academic Editor: Rajib Chowdhury

Copyright © 2023 Sathishkumar Vinayagam et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Wolbachia, a Gram-negative intracellular bacterium, naturally infects many arthropods, including mosquito vectors responsible for the spread of arboviral diseases such as Zika, chikungunya, and dengue fever. Certain *Wolbachia* strains are involved in inhibiting arbovirus replication in mosquitoes, and this phenomenon is currently being studied to combat disease vectors. A study was conducted in four states in north-eastern India to investigate the presence of natural *Wolbachia* infection in wild-caught *Aedes albopictus* and *Aedes aegypti* mosquitoes, the established vectors of dengue. The detection of a *Wolbachia* infection was confirmed by nested PCR and sequencing in the two mosquito species *Ae. aegypti* and *Ae. albopictus*. Positivity rates observed in *Ae. aegypti* and *Ae. albopictus* pools were 38% (44 of 115) and 85% (41 of 48), respectively, and the difference was significant (chi-square = 28.3174, $p = 0.00000010$). Sequencing revealed that all detected *Wolbachia* strains belonged to supergroup B. Although *Wolbachia* infection in *Ae. aegypti* has been previously reported from India, no such reports are available from north-eastern India. Data on naturally occurring *Wolbachia* strains are essential for selecting the optimal strain for the development of *Wolbachia*-based control measures. This information will be helpful for the future application of *Wolbachia*-based vector control measures in this part of the country.

1. Introduction

Vector-borne diseases cause significant loss of life in terms of morbidity and mortality. In north-eastern (NE) India, dengue fever and malaria are endemic and outbreaks are common in various states in the region [1–4]. Mosquito vectors such as *Ae. aegypti* and *Ae. albopictus* cause many dengue outbreaks each year and traditional measures to combat them have not yielded the expected results [5]. Increasing insecticide resistance among the mosquito population further exacerbates this problem. In India, both mosquito species showed different degrees of resistance to

dichlorodiphenyltrichloroethane (DTT) in most states [6, 7]. Therefore, more effective and biologically active vector control measures are needed to prevent these vector-borne diseases [8]. *Wolbachia*, a class of Alphaproteobacteria, is endosymbiotic in several arthropods and filarial nematodes in the biosphere. There are currently 17 supergroups of endosymbiotic *Wolbachia* (A–S, excluding G and R), most of which infect terrestrial arthropods, particularly insects and arachnids [9–11]. Interestingly, certain *Wolbachia* species have been shown to possess a natural ability to alter the biology of the infected host mosquito, making them less susceptible to infection by arboviruses such as dengue virus

(DENV), chikungunya virus (CHIKV), yellow fever virus (YFV), and Zika virus (ZIKV). This property has been exploited by various groups to transfect *Ae. aegypti* with *Wolbachia* strains such as *wMel*, *wMelPop* – CLA, *wAlbB*, and *wMel/wAlbB* [5]. In Orissa, genetically distinct and unique *Wolbachia* species have been reported in the coastal plains, which show completely different characteristics from the other populations of the country [12]. In addition, both *wAlbA* and *wAlbB* *Wolbachia* endosymbionts were observed in *Ae. albopictus* population from the Andaman and Nicobar Islands [13]. A low prevalence of these *Wolbachia* endosymbionts has been observed in Indian wild mosquitoes of *An. culicifacies* and *An. stephensi* species from Tamil Nadu [14]. However, there can be significant overlap in *Wolbachia* strains infecting one host, and different strains can affect the survival of the other [15]. Geographical and ecological factors must also be taken into account. This could have an important impact on the selection of the optimal strain for transfection in *Wolbachia*-based vector control strategies since a detailed assessment of native strains in mosquito populations is first required. The current study was conducted to detect *Wolbachia* infection in adult *Ae. aegypti* and *Ae. albopictus* mosquitoes collected in four different states (Assam, Arunachal Pradesh, Nagaland, and Meghalaya) in NE India by nested PCR using 16S rRNA-specific primers followed by sequencing.

2. Materials and Methods

We used archived mosquito samples collected as part of the previously conducted project “Vector Surveillance for ZIKV in Selected High-Risk Areas” [16, 17]. *Ae. albopictus* and *Ae. aegypti* mosquitoes (adult) were sampled from February 2018 to February 2019 in urban areas from four dengue-prone regions in four different states in NE India: Guwahati (Kamrup Metro district, Assam), Tura (West Garo Hills district, Meghalaya), Pasighat (East Siang district, Arunachal Pradesh), and Dimapur (Dimapur district, Nagaland). The location of the study sites is shown in Figure 1. Adult mosquitoes were originally collected using suction tubes from indoor and outdoor resting sites such as open water tanks, garages, tire dumps, and leaf axils. Mosquitoes collected in the field were separated according to species, collection site, date, sex, and blood-fed status of the female mosquitoes. A maximum of 20 mosquitoes were pooled in one tube and transported to the laboratory in 50 μ L of TRI Reagent (Molecular Research Center, Inc., USA) at 4°C. Samples were uniformly homogenized, and DNA was extracted using the commercially available DNeasy Blood and Tissue Kit (Qiagen) and stored at –20°C until further processing.

With some minor modifications, ITS2-PCR (Internal Transcribed Spacer-2) was performed to validate *Ae. aegypti* and *Ae. albopictus* species in the collected mosquito pools [18]. The PCR contained 1.0 mM MgCl₂ and the primers ITS2-F and ITS2-R (5'-ATCACTCGGCTCGTGGATCG-3', 5'-ATGCTTAAATTTAGGGGGTAGT-3') at a concentration of 1 μ M each. The PCR settings were as follows: 95°C for 5 minutes (initial denaturation), then 35 cycles of 95°C

for 30 seconds (denaturation), 56°C for 30 seconds (annealing), 72°C for 45 seconds (extension), and 72°C for ten minutes the last extension. Positive controls for *Ae. albopictus* were provided by the Indian Council of Medical Research-Vector Control Research Center (ICMR-VCRC, Puducherry) and internal controls were used for *Ae. aegypti*. *Wolbachia* detection in *Aedes* mosquitoes was performed using nested PCR (nPCR) as described by Shaw et al. [19]. From the extracted individual pools of *Ae. aegypti* and *Ae. albopictus* gDNA, 16S rRNA *Wolbachia* gene was targeted by nested PCR. The initial PCR was performed with *Wolbachia* 16S rRNA-specific primer pairs (W-Specf: 5-CATACCTATTCGAAGGGATAG-3 and W-Specr: 5-AGCTTCGAGTGAACCAATTC-3) in a 25 μ L reaction volume with 5 μ L of gDNA. Then, 5 μ L of the primary PCR amplicon was used as a target in the second round of PCR with the following internal primers: 16SNF (5-GAAGGGATAGGGTCGGTTCG-3) and 16SNR (5-CAATTCCCATGGCGGTGACG-3) in a reaction volume of 50 μ L. The PCR protocol for nested 16S rRNA PCR was as follows: initial denaturation at 95°C for 15 minutes, followed by 35 cycles of 15 seconds at 95°C, 25 seconds at 66°C, and 30 seconds at 72°C; followed by a final extension step at 72°C for 5 minutes [19]. *Wolbachia* control DNA provided by ICMR-VCRC, Puducherry, was used as a PCR positive control, and double distilled water (ddH₂O) was used as a negative control. The secondary PCR product, 412 bp in size, was considered specific for *Wolbachia* and sequenced using Sanger’s technique. The *Wolbachia* 16S and *Aedes* ITS2 sequences obtained were checked for sequence quality and compared using the Bioedit Version 7.2 software [20]. The aligned nucleotide sequences were checked for matches and compared to pre-existing high-similarity sequences downloaded from the NCBI GenBank database. All sequences were aligned with Clustal W and exported to MEGA X software for further genetic analysis [21].

3. Results

The project collected a total of 6,229 adult *Aedes* mosquitoes from dengue-endemic areas in four different Northeast states. Details on the distribution of these mosquitoes can be found elsewhere [17]. In short, it was found that *Ae. aegypti* was the predominant *Aedes* species (63.3%) among all mosquitoes collected in the study. In Guwahati, Dimapur, and Tura, which are predominantly urban areas, *Ae. aegypti* was dominant, while in Pasighat, which is surrounded by forested areas, *Ae. albopictus* was predominant [17]. From a total of 515 pools, 163 representative pools from the four regions were randomly selected for the current study. Since *Wolbachia* infection in *Ae. albopictus* is already established in this region, a larger number of *Ae. aegypti* pools were selected for analysis (115 vs. 48). Of the 163 pools, a total of 85 pools were found positive for *Wolbachia* by nPCR (Table 1, Figure 2). Positivity rates observed in *Ae. aegypti* and *Ae. albopictus* were 38% (44 of 115) and 85% (41 of 48), respectively, and the difference was significant (chi-square = 28.3174, $p = 0.00000010$). A total of 17 *Wolbachia* 16S PCR amplicons (8 from *Ae. aegypti* and 9 from *Ae.*




FIGURE 1: District-level map of NE India (7 states) showing the four study areas (GPS coordinates marked with black * sign) with land use/land cover details. (Data source: IRS P6 LISS 111 satellite data from National Remote Sensing Centre, Indian Space Research Organization, Department of Space, Govt. of India, Balanagar, Hyderabad-500037; image created with Bhuvan (<https://bhuvan.nrsc.gov.in/>) and QGIS version. 3.8.3-Zanzibar).

TABLE 1: Distribution and characteristics of *Wolbachia* positive pools by nPCR.

Locations	Mosquitoes species	Gender	No. of pools processed	<i>Wolbachia</i> positive pools by nPCR	Pool positivity rate (%)
Nagaland (Dimapur)	<i>Aedes aegypti</i>	Male	16	4	25.00
		Female	22	0	0
	<i>Aedes albopictus</i>	Males	1	0	0
		Female	3	0	0
		Total	42	4	9.52
Assam (Guwahati)	<i>Aedes aegypti</i>	Male	19	10	52.60
		Female	22	4	18.10
	<i>Aedes albopictus</i>	Male	3	1	33.30
		Female	3	2	66.60
		Total	47	17	36.17
Arunachal Pradesh (Pasighat)	<i>Aedes aegypti</i>	Male	3	2	66.60
		Female	3	2	66.60
	<i>Aedes albopictus</i>	Male	7	7	100
		Female	31	31	100
		Total	44	42	95.45
Meghalaya (Tura)	<i>Aedes aegypti</i>	Male	11	10	90
		Female	19	12	63.10
	<i>Aedes albopictus</i>	Male	0	0	0
		Female	0	0	0
		Total	30	22	73.33




FIGURE 2: Frequency distribution and characteristics of *Wolbachia* positive pools by nPCR.

albopictus pools) were sequenced in the current study using Sanger's technique. The resulting sequences have been deposited in the NCBI GenBank database (accession numbers: OL477363.1-OL477379.1). The phylogenetic maximum likelihood tree was constructed using the Tamura-Nei model [22]. The tree with the highest log likelihood (-663.46) comprised 27 nucleotide sequences (295 positions each) (Figure 3). MEGA X was used to perform evolutionary analysis [21]. When compared to other known sequences from groups A and B, it was found that all isolates in the current study belonged to *Wolbachia* supergroup B. For the confirmation of the field-collected *Aedes* samples as *Ae. aegypti* and *Ae. albopictus*, an ITS2-PCR was performed with corresponding positive controls (Figure 4). A total of 8 ITS2-PCR amplicons (4 *Ae. aegypti* and 4 *Ae. albopictus*) were sequenced and submitted to the NCBI GenBank (ITS2 sequence accession numbers: OP327745 – OP327752). A total of 25 nucleotide sequences were used to construct the phylogenetic tree using the neighbour joining method and the Kimura-2 parameter in MEGA 11 [23, 24]. Analysis of the phylogenetic tree (Figure 5) showed good agreement of the study samples with *Ae. aegypti* and *Ae. albopictus* [18, 25].

4. Discussion

The current study has shown the prevalence of *Wolbachia* species in two important dengue vectors viz. *Ae. aegypti* and *Ae. albopictus* from four dengue endemic areas spread over four states in northeastern India. In both mosquito species, *Wolbachia* supergroup B was detected. The positivity rate was higher in *Ae. albopictus* compared to that of *Ae. aegypti* (85% vs. 38%), and the difference was significant. Although previous studies from Northeast India have reported

Wolbachia in various mosquito vectors, there are no previous reports of *Wolbachia* infection in *Ae. aegypti* [15]. Traditionally, it was assumed that natural infection of *Ae. aegypti* with *Wolbachia* was not common [15, 26]. Previously, researchers used different sets of primers to detect natural infection of *Wolbachia* in mosquito vectors. *Wolbachia* surface protein (*wsp*)-based primers have been widely used to detect *Wolbachia* superinfections in many arthropods. A comparison between *wsp* primers and 16SrRNA-based primers in *Ae. aegypti* mosquitoes has shown that the highest detection rate was achieved with 16SrRNA primers in the US [27]. Malaria vectors such as Anopheles mosquitoes were also assumed not to be naturally infected with *Wolbachia* until a unique 16SrRNA-based PCR demonstrated that *An. gambiae* carry low-level natural *Wolbachia* infection [19].

Subsequently, researchers from Malaysia (25%), the Philippines (16.8%), and the USA (44.8%) have reported natural *Wolbachia* infection in *Ae. aegypti* using *Wolbachia* 16SrRNA primers [27-29]. In 2019, the natural infection of *Ae. aegypti* with *Wolbachia* supergroup B was detected using 16SrRNA-based primers from Coimbatore, Tamil Nadu, India [5]. The reported strain showed 99% homology with the *wAlbB* strain in *Ae. albopictus* [5]. This is similar to the homology levels observed in our study (98%-99%) compared to published sequences in the NCBI database. The abovementioned studies from Malaysia, Philippines, USA, and India observed *Wolbachia* infection in *Ae. aegypti* mosquitoes by screening individual mosquitoes. However, *Wolbachia* infection in screened pools of *Ae. aegypti* has also been described previously [30, 31]. Coon et al. reported two *Wolbachia* 16S rDNA OTUs (operational taxonomic units) in a pool of 30 *Ae. aegypti* larvae collected in Florida in 2014 [30].




FIGURE 3: Phylogenetic analysis of *Wolbachia* strains isolated in the study (maximum likelihood method). Red dots indicate samples from the current study.




FIGURE 4: PCR amplified ITS2 nuclear gene fragments with respective controls of *Ae. aegypti* and *Ae. albopictus* (lanes 1 and 2 are positive controls of *Ae. aegypti* and *Ae. albopictus*, lane 3: negative control, lane 4: 1 kb marker, and lanes 5–14 are sample pools of different states of northeast India).

Similarly, *Wolbachia* 16S OTUs have been detected in pooled *Ae. aegypti* mosquitoes in Thailand [31].

In most countries, including India, *Ae. albopictus* mosquitoes have traditionally been subjected to *Wolbachia* detection using primers based on *Wolbachia* surface protein (*wsp*). In a study conducted in Orissa, India, 1291 male and

female *Ae. albopictus* mosquitoes collected from 15 districts across the state were tested for *Wolbachia* infection using *wsp* primer-based PCR. Among these, 1281 (99.2%) mosquitoes tested positive for *Wolbachia* infection; most were individually infected with supergroups B and A, and some had mixed infection with A and B [12]. Another study from




FIGURE 5: Neighbour-joining tree showing the taxonomic characterization of *Ae. aegypti* and *Ae. albopictus* (blue and green denotes isolates from the current study).

Orissa found that 5% of *Ae. albopictus* were monoinfected with *wAlbA*, 10% with *wAlbB* and 80% with both *wAlbA* and *wAlbB* [32]. Similarly, Ravikumar H et al. also reported *Wolbachia* infection in eight of twenty mosquito species using *wsp* primer sets, with *Wolbachia* A and B observed in *Ae. albopictus* mosquitoes [33]. From Assam, NE India, Soni et al. showed superinfection with *Wolbachia* A and B from Dibrugarh, Tinsukia, and Sibsagar districts; whereas from Tezpur, in *Ae. albopictus*, only the *Wolbachia* supergroup A was detected. [15]. In the Andaman and Nicobar Islands, a total of 57 *Ae. albopictus* mosquitoes (100%) were found to be infected with *Wolbachia* supergroups A and B by *wsp* primer-based PCR [13].

The *wAlbB* strain has been shown to confer protection against arboviruses in mosquito vectors. However, the route of infection seems to play a crucial role. Whether a particular mosquito species is infected with this strain naturally, through transient or stable transfection, has a major impact on resistance to arboviruses such as dengue and chikungunya [34, 35]. While natural infection of *Ae. albopictus* with *wAlbA* and *wAlbB* showed no antiviral activity, stable transfection with the *wMel* strain has been reported to reduce transmission of DENV and CHIKV [34, 35]. Likewise in the case of *Ae. aegypti*, stable transfection with *wMelPop*, *wAlbB*, and *wMel* strains all showed reduced infection rate, viral load, and transmission rates for DENV and CHIKV compared to *Wolbachia*-free mosquitoes [34, 36]. However, such studies on natural populations are

scarce and large-scale studies across different geographic locations are needed to obtain conclusive evidence for the potential antiviral role of wild-caught, naturally infected *Ae. aegypti* and *Ae. albopictus*. Different types of *Wolbachia* strains provide different survival benefits to host mosquitoes and may also entail fitness costs. One variety can also complement or compete with the other. Experimenting with *Ae. albopictus* triple infections with *wMel*, *wAlbA*, and *wAlbB* has shown that the introduction of new *Wolbachia* strains can sometimes lead to unexpected complications in uninfected or naturally infected mosquito vectors [36–38]. Our observation on the molecular detection of *Wolbachia* using 16SrRNA primer-based PCR and sequencing in *Ae. aegypti* and *Ae. albopictus* could have important implications for future intervention strategies based on the transfection of *Wolbachia* strains on *Aedes* mosquitoes. It may or may not play an important role in reducing arbovirus transmission under natural conditions. However, further validation of the definitive presence of natural infection in these mosquito hosts requires additional molecular tests such as the detection of bacteria in the host tissue and their removal after antibiotic treatment and whole genome sequencing [39]. We propose that natural infection of *Wolbachia* in mosquito vectors needs to be delineated, preferably based on multiple lines of evidence in different geographic regions, before initiating vector control measures based on *Wolbachia*-infected mosquitoes.

Data Availability

All the sequences generated in this study have been submitted to the NCBI GenBank and can be obtained from the database. The relevant Accession Numbers are mentioned in the text.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Authors' Contributions

N.P.K., S.J.P., and D.R.B. contributed to conceptualization of the study. S.V., T.N., S.C., S.J.P., D.R.B., P.S., I.P.B., K.S., and N.P.K. contributed equally to conducting the study, analysis, and validation of the results and drafting and critically reviewing the manuscript.

Acknowledgments

The authors acknowledge the help and cooperation of the respective district administrations of Assam, Meghalaya, Arunachal Pradesh, and Nagaland, where the study was carried out and all the project staff involved in the study. The current study was funded by the Indian Council of Medical Research (ICMR) through Grant no. VIR/9/2017/ECD-1 dated 21.08.17.

References

- [1] S. A. Khan, P. Dutta, R. Topno, M. Soni, and J. Mahanta, "Dengue outbreak in a hilly state of Arunachal Pradesh in Northeast India," *The Scientific World Journal*, vol. 2014, Article ID 584093, 6 pages, 2014.
- [2] B. Borkakoty, M. Das, K. Sarma et al., "Molecular characterisation and phylogenetic analysis of dengue outbreak in Pasighat, Arunachal Pradesh, Northeast India," *Indian Journal of Medical Microbiology*, vol. 36, no. 1, pp. 37–42, 2018.
- [3] S. Chetry, S. A. Khan, B. Apum, and P. Dutta, "Circulation of dengue virus-1 Genotype III during 2015 dengue outbreak in Arunachal Pradesh: a maiden report from Northeast India," *International Journal of Infectious Diseases*, vol. 45, pp. 433–434, 2016.
- [4] D. K. Sarma, P. K. Mohapatra, D. R. Bhattacharyya et al., "Malaria in north-East India: importance and implications in the era of elimination," *Microorganisms*, vol. 7, no. 12, p. 673, 2019.
- [5] S. Balaji, S. Jayachandran, and S. R. Prabakaran, "Evidence for the natural occurrence of Wolbachia in *Aedes aegypti* mosquitoes," *FEMS Microbiology Letters*, vol. 366, no. 6, Article ID fnz055, 2019.
- [6] R. B. Kushwah, C. L. Dykes, N. Kapoor, T. Adak, and O. P. Singh, "Pyrethroid-resistance and presence of two knockdown resistance (kdr) mutations, F1534C and a novel mutation T1520I, in Indian *Aedes aegypti*," *PLoS Neglected Tropical Diseases*, vol. 9, no. 1, p. e3332, 2015.
- [7] R. B. Kushwah, P. K. Mallick, H. Ravikumar et al., "Status of DDT and pyrethroid resistance in Indian *Aedes albopictus* and absence of knockdown resistance (kdr) mutation," *Journal of Vector Borne Diseases*, vol. 52, no. 1, pp. 95–98, 2015.
- [8] P. K. Jangir and A. Prasad, "Spatial distribution of insecticide resistance and susceptibility in *Aedes aegypti* and *Aedes albopictus* in India," *International Journal of Tropical Insect Science*, vol. 42, no. 2, pp. 1019–1044, 2021.
- [9] E. Glowaska, A. Dragun-Damian, M. Dabert, and M. Gerth, "New Wolbachia supergroups detected in quill mites (Acari: syringophilidae)," *Infection, Genetics and Evolution*, vol. 30, pp. 140–146, 2015.
- [10] G.-H. Wang, L.-Y. Jia, J.-H. Xiao, and D.-W. Huang, "Discovery of a new Wolbachia supergroup in cave spider species and the lateral transfer of phage WO among distant hosts," *Infection, Genetics and Evolution*, vol. 41, pp. 1–7, 2016.
- [11] E. Lefoulon, T. Clark, F. Borveto et al., "Pseudoscorpion Wolbachia symbionts: diversity and evidence for a new supergroup S," *BMC Microbiology*, vol. 20, no. 1, p. 188, 2020.
- [12] B. Das, T. Satapathy, S. K. Kar, and R. K. Hazra, "Genetic structure and Wolbachia genotyping in naturally occurring populations of *Aedes albopictus* across contiguous landscapes of Orissa, India," *PLoS One*, vol. 9, no. 4, Article ID e94094, 2014.
- [13] A. Sivan, A. N. Shriram, D. Bhattacharya, and P. Vijayachari, "Wolbachia endobacterium in wild population of *Aedes albopictus* (skuse) (Diptera: Culicidae) and phylogeny from andaman and nicobar Islands, India," *Journal of Vector Borne Diseases*, vol. 51, no. 3, pp. 235–238, 2014.
- [14] S. G. Sankar, T. M. Sundari, and A. A. P. Anand, "Low prevalence of natural Wolbachia in major malarial vectors *Anopheles culicifacies* s.l., and *Anopheles stephensi*: a first report," *bioRxiv*, 2020.
- [15] M. Soni, C. Bhattacharya, J. Sharma, S. A. Khan, and P. Dutta, "Molecular typing and phylogeny of Wolbachia: a study from Assam, North-Eastern part of India," *Acta Tropica*, vol. 176, pp. 421–426, 2017.
- [16] N. P. Kumar, A. Kumar, D. Panneer et al., "Nation-wide vector surveillance on Zika and Dengue did not indicate transmission of the American lineage-pandemic Zika virus in India," *International Journal of Infectious Diseases*, vol. 113, pp. 119–124, 2021.
- [17] S. Chetry, S. J. Patgiri, D. R. Bhattacharyya, P. Dutta, and N. P. Kumar, "Incrimination of *Aedes aegypti* and *Aedes albopictus* as vectors of dengue virus serotypes 1, 2 and 3 from four states of Northeast India," *Access Microbiology*, vol. 2, no. 4, p. acmi000101, 2020.
- [18] K. Gupta, R. Dhawan, M. Kajla et al., "Molecular identification of *Aedes aegypti* mosquitoes from Pilani region of Rajasthan, India," *Journal of Vector Borne Diseases*, vol. 53, no. 2, pp. 149–155, 2016.
- [19] W. R. Shaw, P. Marcenac, L. M. Childs et al., "Wolbachia infections in natural *Anopheles* populations affect egg laying and negatively correlate with Plasmodium development," *Nature Communications*, vol. 7, no. 1, Article ID 11772, 2016.
- [20] T. A. Hall, "BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT," *Nucleic Acids Symposium Series*, vol. 41, pp. 95–98, 1999.
- [21] S. Kumar, G. Stecher, M. Li, C. Knyaz, and K. Tamura, "Mega X: molecular evolutionary genetics analysis across computing platforms," *Molecular Biology and Evolution*, vol. 35, no. 6, pp. 1547–1549, 2018.
- [22] K. Tamura, "Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C-content biases," *Molecular Biology and Evolution*, vol. 9, no. 4, pp. 678–687, 1992.
- [23] M. Nei and S. Kumar, *Molecular Evolution and Phylogenetics*, Oxford University Press, Oxford, UK, 2000.

- [24] K. Tamura, G. Stecher, and S. Kumar, "MEGA11: molecular evolutionary genetics analysis version 11," *Molecular Biology and Evolution*, vol. 38, no. 7, pp. 3022–3027, 2021.
- [25] G. Minard, V. Tran Van, F. H. Tran et al., "Identification of sympatric cryptic species of *Aedes albopictus* subgroup in Vietnam: new perspectives in phyllosymbiosis of insect vector," *Parasites & Vectors*, vol. 10, no. 1, p. 276, 2017.
- [26] A. Gloria-Soria, T. G. Chiodo, and J. R. Powell, "Lack of evidence for natural Wolbachia infections in *Aedes aegypti* (Diptera: Culicidae)," *Journal of Medical Entomology*, vol. 55, no. 5, pp. 1354–1356, 2018.
- [27] A. Kulkarni, W. Yu, J. Jiang et al., "Wolbachia pipientis occurs in *Aedes aegypti* populations in New Mexico and Florida, USA," *Ecology and Evolution*, vol. 9, no. 10, pp. 6148–6156, 2019.
- [28] C. H. J. Teo, P. K. C. Lim, K. Voon, and J. W. Mak, "Detection of dengue viruses and Wolbachia in *Aedes aegypti* and *Aedes albopictus* larvae from four urban localities in Kuala Lumpur, Malaysia," *Tropical Biomedicine*, vol. 34, no. 3, pp. 583–597, 2017.
- [29] T. M. Carvajal, K. Hashimoto, R. K. Harnandika, D. M. Amalin, and K. Watanabe, "Detection of Wolbachia in field-collected *Aedes aegypti* mosquitoes in metropolitan Manila, Philippines," *Parasites & Vectors*, vol. 12, no. 1, p. 361, 2019.
- [30] K. L. Coon, M. R. Brown, and M. R. Strand, "Mosquitoes host communities of bacteria that are essential for development but vary greatly between local habitats," *Molecular Ecology*, vol. 25, no. 22, pp. 5806–5826, 2016.
- [31] P. Thongsripong, J. A. Chandler, A. B. Green et al., "Mosquito vector-associated microbiota: metabarcoding bacteria and eukaryotic symbionts across habitat types in Thailand endemic for dengue and other arthropod-borne diseases," *Ecology and Evolution*, vol. 8, no. 2, pp. 1352–1368, 2018.
- [32] I. Mohanty, A. Rath, S. P. Swain, N. Pradhan, and R. K. Hazra, "Wolbachia population in vectors and non-vectors: a sustainable approach towards dengue control," *Current Microbiology*, vol. 76, no. 2, pp. 133–143, 2019.
- [33] H. Ravikumar, N. Ramachandraswamy, and H. P. Puttaraju, "Molecular strain typing of Wolbachia infection from Indian mosquitoes using wsp gene," *Asian Pacific Journal of Tropical Disease*, vol. 1, no. 2, pp. 106–109, 2011.
- [34] K. N. Johnson, "The impact of Wolbachia on virus infection in mosquitoes," *Viruses*, vol. 7, no. 11, pp. 5705–5717, 2015.
- [35] S. T. Ogunlade, M. T. Meehan, A. I. Adekunle, D. P. Rojas, O. A. Adegboye, and E. S. McBryde, "A review: aedes-borne arboviral infections, controls and wolbachia-based strategies," *Vaccines*, vol. 9, no. 1, p. 32, 2021.
- [36] P. A. Ross, K. L. Robinson, Q. Yang et al., "A decade of stability for wMel Wolbachia in natural *Aedes aegypti* populations," *PLoS Pathogens*, vol. 18, no. 2, Article ID e1010256, 2022.
- [37] X. Liang, J. Liu, G. Bian, and Z. Xi, "Wolbachia inter-strain competition and inhibition of expression of cytoplasmic incompatibility in mosquito," *Frontiers in Microbiology*, vol. 11, p. 1638, 2020.
- [38] E. P. Caragata, H. L. C. Dutra, P. H. F. Sucupira, A. G. A. Ferreira, and L. A. Moreira, "Wolbachia as translational science: controlling mosquito-borne pathogens," *Trends in Parasitology*, vol. 37, no. 12, pp. 1050–1067, 2021.
- [39] E. Chrostek and M. Gerth, "Is *Anopheles gambiae* a natural host of Wolbachia?" *mBio*, vol. 10, no. 3, Article ID e00784-19, 2019.