

Research Article

Catch-All No More: Integrative Systematic Revision of the Genus *Allolobophora* Eisen, 1874 (Crassiclitellata, Lumbricidae) with the Description of Two New Relict Earthworm Genera

Alejandro Martínez Navarro ¹, Sergio Jiménez Pinadero ¹, Thibaud Decaëns ²,
Mickaël Hedde ³, Marta Novo ¹, Dolores Trigo ¹ and Daniel Fernández Marchán ¹

¹Biodiversity, Ecology and Evolution Department, Faculty of Biology, Universidad Complutense de Madrid, Madrid, Spain

²CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

³Eco&Sols, INRAE, IRD, CIRAD, SupAgro Montpellier, Montpellier, France

Correspondence should be addressed to Alejandro Martínez Navarro; alejma21@ucm.es

Received 20 September 2022; Revised 25 December 2022; Accepted 7 January 2023; Published 15 February 2023

Academic Editor: Muhammad Muddassir Ali

Copyright © 2023 Alejandro Martínez Navarro et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The taxonomy of earthworms has been riddled by instability, lack of systematically useful characters, and lax diagnoses of some genera. This has led to the use of some genera such as *Allolobophora* Eisen, 1874 as taxonomic wastebaskets, blurring their evolution and biogeographical history. The implementation of molecular techniques has revolutionized the systematics of the genus; however, some of its species have not been previously included in molecular phylogenetic analyses. Thus, the molecular markers COI, 16S, ND1, 12S, and 28S were sequenced for six endemic species including several taxa of *Allolobophora* and *Aporrectodea* Örely, 1885 (another related catch-all genus). Phylogenetic relationships determined by Bayesian inference and maximum likelihood analyses support the status of two of the six taxa examined (*Allolobophora burgondiae* Bouché, 1972 and *Aporrectodea icterica* Savigny, 1826) as part of *Allolobophora* sensu stricto and a presumed synonymy between *Allolobophora* and *Heraclescolex* Qiu and Bouché, 1998. Branch lengths and average pairwise genetic distances support the transfer of *Allolobophora satchelli* Bouché, 1972 to the genus *Panoniona* Mršić and Šapkarev, 1988 and the emergence of two new genera, *Heraultia* gen. nov. and *Vosgesia* gen. nov., endemic to France, hosting *Allolobophora tiginosa* Bouché, 1972 and *Allolobophora zicsii* Bouché, 1972, respectively. The aforementioned changes of status and the diagnosis for *Heraultia* and *Vosgesia* are presented. These results provided more evolutionarily and biogeographically coherent earthworm groups and highlighted that the Maghreb and the area around the Alps are potential key locations for the diversification of *Allolobophora* and several lineages of Lumbricidae.

1. Introduction

The taxonomy of earthworms has proven to be very unstable over the last two centuries. This is due to a lack of systematically useful morphological characters and the fact that most of them are homoplasious or symplesiomorphic [1]. Within Lumbricidae Rafinesque-Schmaltz 1805, the most speciose family of the Palaearctic, some genera have been inadvertently used as taxonomic wastebaskets due to excessively loose diagnoses. As shown by the molecular phylogenetic analyses of Dominguez et al. [2], such catch-

all taxa, e.g., *Aporrectodea* Örely, 1885, *Allolobophora* Eisen, 1874, or *Helodrilus* Hoffmeister, 1885, contain independent evolutionary lineages which should be described as different genera.

Allolobophora was established by Eisen in 1874 based on external morphological characters and included seven species, but without selecting a type species. Further species were progressively added to a somewhat refined *Allolobophora* (as internal characters were added to its diagnosis), but after the revision of Pop [3], it became a catch-all genus [1]. Later works suggested that *Allolobophora* should be

divided into subgenera or even different genera [4–8], but with little consensus.

Only the designation of *Allolobophora chlorotica* (Savigny, 1826) as its genus type allowed the delimitation of *Allolobophora sensu stricto* and *Allolobophora sensu lato*, which includes most of its species [1]. The inclusion of *All. chlorotica* in the phylogenetic analysis of Domínguez et al. [2] corroborated this view: *All. chlorotica* and other green pigmented species (*Allolobophora dubiosa* Örley, 1881, *Allolobophora moebii* Michaelsen, 1895, and *Allolobophora molleri* Rosa, 1889) formed a clade separated from the species assigned to the subgenus *Allolobophora (Gatesona)* Bouché, 1972 and to *Allolobophora sensu lato* (or *Karpatodinariona* Mršić and Šapkarev, 1988 and *Serbiona* Mršić and Šapkarev, 1988). Further molecular phylogenetic works cemented this separation by establishing the genus *Gatesona* [9] and by amending the genus *Cernosvitovia* Omodeo, 1956 to include several Balkanic species of *Allolobophora sensu lato* [10].

Eophila Rosa, 1893 is in strong conflict with *Allolobophora*, as numerous species have been transferred between the two genera in successive taxonomic revisions. It has also been considered synonymous with *Heraclescolex* Qiu and Bouché, 1998, a genus consisting mainly of green pigmented earthworms with little distinction from the other two genera. The inclusion of the type species of the genus *Eophila*, *Eophila tellinii* Rosa, 1888, in molecular phylogenetic analysis by de Sosa et al. [11] has eliminated the confusion: *Eophila* should be restricted to *Eo. tellinii*, *Eophila gestroi* (Cognetti de Martiis, 1905), and *Eophila crodabepis* [12]. Furthermore, the genotype of *Heraclescolex (All. moebii)* was nested with *All. chlorotica*, thus suggesting the synonymy of the latter genera.

Koinodrillus Qiu and Bouché, 1998 was created to accommodate species previously assigned to *Allolobophora* and/or *Aporrectodea*. Its type species, *Aporrectodea georgii* Michaelsen, 1890, and a few other representatives (*Aporrectodea jassyensis* (Michaelsen, 1891), *Aporrectodea limicola* (Michaelsen, 1890), *Aporrectodea rosea* (Savigny, 1826), and *Allolobophora oliveirae* (Rosa, 1894)) were included in phylogenetic trees by Domínguez et al. [2], but the results did not strongly support this genus. On the contrary, *Ap. georgii* and *Ap. jassyensis* formed a clade, but the rest of the species appeared scattered within other well-supported clades or behaved as rogue taxa (with unstable positions).

Despite the significant advances in the taxonomy and systematics of the French Lumbricidae in recent years, the phylogenetic relationships of some rare endemic species remain uncertain. For example, *Allolobophora burgondiae* (Bouché, 1972) and *Allolobophora satchelli* (Bouché, 1972) were originally placed within *Allolobophora sensu stricto* by Bouché [4] and retained within the genus by Qiu and Bouché [8], and their relationships were not been questioned; yet they were never been included in molecular phylogenetic analyses. *Allolobophora tiginosa* (Bouché, 1972) and *Aporrectodea icterica* (Savigny, 1826) were transferred from *Allolobophora sensu stricto* to *Koinodrillus* and *Heraclescolex*, respectively, in Qiu and Bouché [13], while *Allolobophora zicsii* Bouché, 1972 was transferred from *Allolobophora sensu lato* to *Koinodrillus* [14]. Finally, *Apor-*

rectodea pseudoantipai (Qiu and Bouché, 1998) was originally placed within *Koinodrillus*.

This paper places the aforementioned species in a molecular phylogenetic context to (i) place the last few *incerta sedis* French endemic species into their appropriate genera, (ii) interrogate whether *Allolobophora* and *Heraclescolex* are synonymous, and (iii) test the validity of *Koinodrillus*.

2. Materials and Methods

2.1. Samplings and Studied Specimens. This study follows the methodology of Marchán et al. [9, 15], and the method description partially reproduces their wording. 15 specimens of six species of Lumbricidae (genera *Allolobophora* and *Aporrectodea*) were collected during two sampling surveys in southeastern and northeastern France in spring 2021 and autumn 2021, respectively. The list of species and the sampling localities are given in Table 1. Individuals were collected by digging up the soil and sorting by hand, rinsing with water, and fixing in pure ethanol to allow further molecular analyses. Sampling and handling of specimens were done ethically and in accordance with Directive 2010/63/EU.

Species classification and morphological diagnoses were made under a binocular stereomicroscope using the set of external and internal morphological characters used by Qiu and Bouché [16] and following the format established by Domínguez et al. [17]. The following main external morphological characters were considered: mean length, mean number of segments, mean weight, pigmentation, type of prostomium, setal arrangement, position of papillae, position of first dorsal pore, nephridial pore arrangement, position and development of male pores, position and development of female pores, position of spermathecal pores, position of clitellum, and position of tubercula pubertatis. The main internal anatomical features were position of oesophageal hearts, position and morphology of calciferous glands, position of crop, position of gizzard, type of typhlosole, shape of nephridial bladders, number and position of seminal vesicles, and number and position of spermathecae.

2.2. DNA Extraction and Sequencing. After morphological identification, total genomic DNA was extracted from ventral integument samples of approximately 5 × 5 mm by using the SpeedTools Tissue DNA Extraction kit (Biotools). Regions of the nuclear gene 28S rRNA and mitochondrial 16S rRNA, 12S rRNA, NADH dehydrogenase (ND1), and cytochrome oxidase subunit 1 (COI) were amplified by polymerase chain reaction (PCR), using the primers described in Pérez-Losada et al. [18] and Folmer et al. [19]. PCR reactions were performed using a GeneAmp Multicycler Ep gradient (Eppendorf) under the following conditions: an initial denaturation step (5 min at 94°C); 40 cycles (35 for ND1) consisting of denaturation at 95 °C for 30 s, annealing (between 45°C and 55°C) for 45 s, and extension at 72°C for 1 min; and a final extension step (5 min at 72°C). The amplified PCR products were purified using the Multiscreen PCRµ96 purification kit (Millipore) and sequenced in Macrogen (Spain). DNA sequences

TABLE 1: Sampling information for the target species of this work.

Species	N	Collectors	Sampling location	Date	Coordinates
<i>Allolobophora burgondiae</i>	3	Marchán, Decaëns, Della Vedova	Bouze-lès-Beaune, Côte-d'Or, France	30-Oct-2021	47.062, 4.760
<i>Allolobophora satchelli</i>	3	Marchán, Decaëns, Della Vedova	Saint-Louis, Haut-Rhin, France	04-Nov-2021	47.618, 7.538
<i>Allolobophora tiginosa</i>	1	Hedde, Marsden, Gerard, Beauchesne	Puéchabon, Hérault, France	09-Feb-2021	43.713, 3.605
<i>Aporrectodea icterica</i>	3	Marchán, Decaëns, Della Vedova	Plombières-les-Bains, Vosges, France	02-Nov-2021	47.988, 6.512
<i>Allolobophora zicsii</i>	3	Marchán, Decaëns, Della Vedova	Luxeuil-les-Bains, Haute-Saône, France	02-Nov-2021	47.836, 6.359
<i>Aporrectodea pseudoantipai</i>	2	Marchán, Decaëns, Della Vedova	Trépot, Doubs, France	31-Oct-2021	47.170, 6.168

obtained in this study are available in GenBank, under accession numbers OQ224974–OQ224979 for COI, OQ236700–OQ236705 for ND1, OQ225516–OQ225521 for 28S, OQ225522–OQ225527 for 12S, and OQ225532–OQ225537 for 16S; COI sequences, together with metadata, are publicly available in the dataset DS-ALLREV in BOLD (doi:10.5883/DS-ALLREV).

2.3. Phylogenetic Analyses. Sequences were aligned using MAFFT v.7 [20] with default parameters and concatenated in BioEdit [21], obtaining a sequence of 3,260 base pairs for each species. Sequences reported by Dominguez et al. [2], Domínguez et al. [17], Pérez-Losada et al. [18], Pérez-Losada et al. [22], Pérez-Losada et al. [23], Paoletti et al. [12], de Sosa et al. [11], Bozorgi et al. [24], Jiménez Pinadero et al. [25], and Marchán et al. [26] with representatives of most Lumbricidae genera and two members of the closest families (Hormogastridae and Criodrilidae) were downloaded from GenBank and used as a reference dataset. The included species and their source publications are listed in Supplementary Table 1.

The best-fitting evolutionary model for each partition was selected with jModelTest v. 2.1.3 [27] following the Akaike Information Criterion (AIC; [28]) and the Bayesian Information Criterion (BIC; [29]). GTR+I+G was selected as the best evolutionary model for COI, 28S, and ND1, whereas GTR+G and HKY+I+G were chosen for the molecular markers 12S and 16S, respectively. Phylogenetic relationships were inferred by Bayesian inference analysis (BI) using MrBayes v.3.2.6 [30] implemented in CIPRES Science Gateway V. 3.3 [31]. Parameters were set to 50 million generations and sampled every 5,000th generation (10,000 trees). Two independent runs of four chains each were performed, and 20% of the trees were discarded as burn-in. The remaining trees were combined and summarized on a 50% majority-rule consensus tree. Maximum likelihood phylogenetic inference was performed using RAXML-NG [32], starting from 10 random initial trees and 1000 fast bootstrap replicates. Clade support values over 70% and 90% (for bootstrap and posterior probability, respectively) were considered high.

Uncorrected average pairwise distances between the studied species for the molecular markers COI and 16S were calculated in MEGA 11 [33] to support their status as separate species and to investigate genetic distances within and between genera.

3. Results

3.1. Morphological Study. The studied specimens fitted the diagnoses of their respective species [4, 13, 14].

3.2. Molecular Phylogenetic Analysis. Phylogenetic relationships determined by Bayesian inference (Figure 1) and maximum likelihood (Supplementary Figure 1) were congruent with previous phylogenetic analyses and between them, with the main internal branches resolved receiving high statistical support. The results do not support the monophyly of *Allolobophora* as currently defined, as the

studied species appeared scattered throughout different clades. However, *All. burgondiae* and *Ap. icterica* were recovered in a clade with *All. chlorotica* and the species formerly assigned to *Heraclescolex* (*All. moebii*—designated as its type—*All. molleri*, *Ap. icterica*, and *Ap. dubiosa*). In contrast, the four remaining species were recovered as more distantly related taxa, not sharing the most recent common ancestor with *Allolobophora*. Although with low support, *All. zicsii* was recovered in both phylogenetic analysis as a sister species of *Avelona ligra* (Bouché, 1969), the only known species of the genus *Avelona* Qiu and Bouché, 2000; both species formed a clade sister to the strongly supported genus-level clade *Gatesona*. A similar scenario was found for *All. satchelli*, who was recovered as a sister taxon to *Panionia leoni* Michaelsen, 1891. With respect to *Allolobophora tiginosa*, this taxon was placed in a large clade comprising *Castellodrilus* Qiu and Bouché, 1998; *Zophoscolex* Qiu and Bouché, 1998; *Ethnodrilus* Bouché, 1972; *Proselodrilus* Bouché, 1972; and *Cataladrilus* Qiu and Bouché, 1998, forming a well-supported clade with the last two mentioned genera. However, the precise phylogenetic relationship between them remained unsolved. Finally, the results for *Aporrectodea pseudoantipai* show that it belongs in a clade with the representatives of *Aporrectodea* (except for *Aporrectodea rosea*, which is considered a rogue taxon), with *Aporrectodea handlirschi* (Rosa, 1897) as its closest relative. *Aporrectodea georgii* (type species of *Koinodrilus*) showed no affinity with the species once assigned to *Koinodrilus* (*zicsii*, *pseudoantipai*, *rosea*, *jassyensis*, *limicola*, and *oliveirae*). *Allolobophora bartolii* (Bouché, 1970), although with minor support, does not appear to be related to *Allolobophora sensu stricto*. Instead, it is more closely related to *Eophila* Rosa, 1893 and *Imetescolex* Szederjesi et al., 2022.

The four taxa not placed within the *Allolobophora sensu stricto* clade showed high distance values for both markers with representatives of that clade (including *All. burgondiae* and *All. icterica*), with *All. satchelli* showing the smallest distances with the species of the genus (Table 2). However, *All. satchelli* displayed its lowest distance values for both markers with *P. leoni* (14.71% for COI, the lowest value for this marker, and 6.00% for 16S). For *All. zicsii*, UAPG distances for the molecular markers COI and 16S were high between the species and every taxon included in the analysis, even with its closest relative *Av. ligra* (19.17% for COI and 14.02% for 16S). *All. tiginosa* showed its lowest distance with the type species of the genus *Cataladrilus*, *C. monticola* (10.61% for 16S), while *Ap. pseudoantipai* showed generally lower distances with *Ap. handlirschi* than with any other species of *Allolobophora*.

4. Discussion

4.1. Systematic Implications. The inclusion of *All. burgondiae* and *Ap. icterica* within a well-supported clade with *All. chlorotica*, *All. molleri*, and *All. dubiosa* and the low UAPG distance values among them suggest that both species should be placed within *Allolobophora*. The phylogenetic position of *Ap. pseudoantipai* with the representatives of *Aporrectodea*,




FIGURE 1: Detail of the clades recovered by Bayesian inference that includes the species of study. The phylogenetic analysis was performed using the concatenated sequence of the five molecular markers (COI, ND1, 16S, 12S, and 28S). Vertical grey lines show differences in topology according to a maximum likelihood analysis. Support values are shown besides corresponding nodes (posterior probability, left; bootstrap, right). The taxa included in this work are shown in bold. *Allolobophora* sensu stricto clade is shaded in green. The tree containing all phylogenetic relationships is shown in Supplementary Figure 2.

all of them forming a well-supported genus-level clade, suggests that it should remain within that genus. The possibility that *Eiseniona* Omodeo, 1956 (with *Eiseniona handlirschi* as type species) is a valid genus and that *Ap. pseudoantipai* should belong to it cannot be discarded without sampling additional representatives.

On the other hand, the nonmonophyly of *All. satchelli*, *All. zicsii*, and *All. tiginosa* with the species of *Allolobophora* sensu stricto implies these species should be assigned to other genera. For *All. satchelli*, the Bayesian inference and the UAPG distances support the inclusion of the species within the genus *Panoniona* Mršić and Šapkar, 1988, changing its status to *Panoniona satchelli*. Interestingly, *All. bartolii* was placed by Qiu and Bouché [8] within *Panoniona*, which was not supported by molecular phylogenetics results. This highly polymorphic species (probably being a species complex) requires further study before a new genus can be proposed for it. The high distance values between *All. zicsii* and all the studied species and the long branches that separate this taxon from its closest relative suggest the designation of a new genus for this species. Similarly, *All. tiginosa* has been shown to be closely related to the genera

Prosello-drilus and *Cataladrilus* but without belonging to the same genus-level clade, thus justifying the creation of its own genus.

The nonmonophyly of *Aporrectodea georgii* with the other species included in *Koinodrilus* suggests that it would be a monospecific genus if retained. Also, *Aporrectodea georgii* does not appear to be related to *Aporrectodea* sensu stricto, the closest relatives of *Aporrectodea trapezoides* (Dugès 1828), so it should not be classified within *Aporrectodea*. Therefore, *Koinodrilus* would be a valid name, and a new one would not be necessary. Interestingly, Marchán et al. [34] recovered a close phylogenetic relationship between *Ap. georgii* and *All. chlorotica* when using the novel phylogenomic technique AHE (Anchored Hybrid Enrichment, [35]): this relationship was recovered with low support in the maximum likelihood analysis based on the traditional Sanger-sequenced markers. As the taxonomic sampling was limited (43 species), an increased number of species could confirm or deny such an evolutionary affinity.

Proposed taxonomic changes

Phylum Annelida Lamarck, 1802

Subphylum Clitellata Michaelsen, 1919

Class Oligochaeta Grube, 1850
 Superorder Megadrili Benham, 1890
 Order Haplotaxida Michaelsen, 1900
 Family Lumbricidae Rafinesque-Schmaltz, 1815
 Tribe Proselodrilini Qiu and Bouché, 1998
 Genus *Heraultia* Martínez Navarro and Marchán, 2022
 Type species: *Heraultia tiginosa* stat. nov. (Bouché, 1972)

Material studied: 1 adult specimen; BOLD sample ID: ALL-016; Occitanie, Hérault, Puéchabon; latitude/longitude: 43.713/3.605; 09-Feb-2021; M. Hedde, Marsden, Gerard, Beachesne leg. Specimen deposited in UCM-LT.

Diagnosis:

Small-sized (50-75 mm and 0.2-0.29 gr) Proselodrilini with less than 200 segments and absence of pigmentation. Prostomium is epilobous. Setae are closely paired. First dorsal pore in 4/5. Male pores in 1/2 15 with well-developed porophores. Spermathecal pores in 10/11, 11/12, and 12/13. Nephridial pores aligned. Clitellum in 21-36. *Tubercula pubertatis* in 27-29, winglet shaped. Oesophageal hearts in 6-11; Calciferous glands poorly developed in 13 and 14, without diverticula or dilations. Crop in 15-16. Glizzard in 17-18. Typhlosole is bifid. Three pairs of globular spermathecae in 11, 12, and 13, rarely double. Glandular papillae in 10, 11, and 26-32. Two pairs of seminal vesicles in 11 and 12, poorly developed. Nephridial bladders are U-shaped (incurvate and reclinate). Cross section of longitudinal musculature elementary (sensu Bouché, 1972).

Differential diagnosis:

Heraultia shares with *Proselodrilus* and *Cataladrilus* (its closest relatives) the anterior position of the clitellum and the number and position of seminal vesicles (two pairs in 11 and 12), but it is clearly separated from both by the poorly developed calciferous glands (well-developed and usually with diverticula or dilations in 11 in *Proselodrilus* and *Cataladrilus*) and by the position of the spermathecae (10/11, 11/12, and 12/13 vs. (12/13), 13/14, and 14/15 in *Proselodrilus* and 9/10 and 10/11 in *Cataladrilus*). For the diagnoses of the compared genera, see Table 3.

Other species included: none.

Tribe Lumbricini Qiu and Bouché, 1998

Genus *Allolobophora* Eisen, 1874

Type species: *Allolobophora chlorotica* (Savigny 1826)

Material studied: (a) *Allolobophora burgondiae*: 3 adult specimens; BOLD sample ID: ALL-002, ALL-003, ALL-004; Bourgogne, Côte d'Or, Bouze-lés-Beaune; latitude/longitude: 47.062/4.760; 30-Oct-2021; R. Della Vedova, T. Decaëns, D. Fernández Marchán leg. Specimen deposited in UCM-LT; (b) *Allolobophora icterica*: 3 adult specimens; BOLD sample ID: ALL-010, ALL-011, ALL-012; Grand Est, Vosges, Plombières-les-Bains; latitude/longitude: 47.988/6.512; 02-Nov-2021; R. Della Vedova, T. Decaëns, D. Fernández Marchán leg. Specimen deposited in UCM-LT.

Diagnosis:

Small-to-medium-sized Lumbricini with green pigmentation (sometimes absent). Prostomium is epilobous, no transversal furrows. Setae are closely paired. First dorsal pore in 4/5 (rarely 3/4, 5/6, and 7/8). Male pores in 1/2 15 with well-developed porophores. Presence of spermatophores. Spermathecal pores in variable position and number

between 7/8, 8/9, 9/10, and 10/11, simple. Nephridial pores irregularly distributed (*en solfège*). Clitellum moderately posterior to extremely posterior, starting between 28 and 73 and ending between 36 and 78. Oesophageal hearts in 6-11. Calciferous glands in 10-14 with diverticula in 10. Crop in 15-16. Glizzard in 17-18. Typhlosole is usually trifid (sometimes bifid or with 4 lamellae). Variable number of simple spermathecae, usually 3 or 4 pairs. Four pairs of seminal vesicles in 9, 10, 11, and 12. Nephridial bladders are U-shaped (incurvate and reclinate or proclinate). The main morphological characters of the species of *Allolobophora* sensu stricto (see below) can be found in Table 4.

Differential diagnosis:

Even though *Cernosvitovia* as redefined by Popović et al. [10] is a morphologically heterogeneous genus, *Allolobophora* can be distinguished from it by its pigmentation (if present: green vs. brown) and either the position of the male pore (1/2 15 vs. 25-30) or the position of the spermathecae (variable in 7/8, 8/9, 9/10, and 10/11 vs. 9/10, 10/11, 11/12, and (12/13)).

Allolobophora is clearly separated from *Gatesona* by the absence of pigmentary dots in cephalic segments, by its general pigmentation (if present: green vs. brown), by the absence of very conspicuous genital papillae between segment 9 and the end of the clitellum and by the start of the clitellum (28-73 vs. 23-27). They also differ in the position of the spermathecae (variable in 7/8, 8/9, 9/10, and 10/11 vs. 9/10, 10/11, and (11/12)).

Allolobophora differs from *Panoniona* by the more posterior position of the clitellum, frequently green pigmentation vs. absence of pigmentation, repetition of spermathecae (simple vs. double or multiple), number of seminal vesicles (four pairs in 9, 10, 11, and 12 vs. two pairs in 11 and 12), and shape of nephridial bladders (U-shaped vs. J-shaped).

Allolobophora can be separated from *Eophila* by its generally smaller size, its pigmentation (if present: green vs. purplish-brown in bands or brown), and the position of the spermathecae (variable in 7/8, 8/9, 9/10, and 10/11 vs. 9/10 and 10/11). The comparison of the mentioned genera is shown in Table 3.

Species included the following:

- (i) *Allolobophora chlorotica* (Savigny, 1826)
- (ii) *Allolobophora burgondiae* Bouché, 1972
- (iii) *Allolobophora icterica* (Savigny, 1826)
- (iv) *Allolobophora molleri* Rosa, 1889
- (v) *Allolobophora moebii* Michaelsen, 1895
- (vi) *Allolobophora dubiosa* Örley, 1881

Remarks: the species assigned to *Heraclescolex* by Qiu and Bouché [13] form a remarkably homogenous group in terms of morphology and anatomy. Since four species formerly assigned to *Heraclescolex* (including the type species) have been recovered within *Allolobophora* by phylogenetic analyses, it is likely that the rest will also belong to this genus. The taxonomic status of the *Allolobophora molleri*-

TABLE 2: Uncorrected average pairwise genetic (UAPG) distances between the species of study and representatives of the genus *Allobophora*, *Cataladrilus*, *Proselodrilus*, *Avelona*, *Panoriona*, and *Aporrectodea*, for 16S (above the diagonal) and COI (below the diagonal). The cell could be shown in green (low genetic distance between the compared taxa), yellow (medium), or red (high).

	<i>Cat. monticola</i>	<i>Pr. pyrenaicus</i>	<i>Pr. biauriculatus</i>	<i>P. leoni</i>	<i>All. chlorotica</i>	<i>All. dubiosa</i>	<i>All. molleri</i>	<i>Av. ligra</i>	<i>All. burgondiae</i>	<i>Ap. pseudoantipai</i>	<i>All. zicsii</i>	<i>Ap. ictérica</i>	<i>All. satchelli</i>	<i>All. tiginosa</i>	<i>Ap. handlirschi</i>
<i>Cat. monticola</i>															
<i>Pr. pyrenaicus</i>	11.94		14.67	16.36	13.88	15.36	15.38	16.97	16.03	17.74	17.00	15.69	15.53	15.00	14.19
<i>Pr. biauriculatus</i>	16.01	17.00		14.55	11.07	14.55	15.41	14.36	15.41	17.46	16.72	16.22	12.71	14.91	14.91
<i>P. leoni</i>	20.63	19.97	18.15		9.33	11.63	11.81	13.55	12.81	14.00	14.81	11.30	6.00	14.17	10.65
<i>All. chlorotica</i>	17.99	21.45	19.47	20.46		7.67	8.51	11.06	7.65	13.37	14.02	7.49	7.36	13.52	9.00
<i>All. dubiosa</i>	19.80	20.13	18.98	19.80	17.99		7.82	12.71	10.15	15.68	14.98	6.15	10.67	15.17	9.98
<i>All. molleri</i>	20.87	20.03	18.86	20.37	19.20	15.19		12.23	10.17	14.70	14.45	5.82	11.69	13.52	10.83
<i>Av. ligra</i>	19.31	19.64	18.65	18.65	18.48	17.66	19.37		13.38	17.26	14.02	12.71	11.54	15.69	11.87
<i>All. burgondiae</i>	19.83	19.83	20.33	19.67	19.01	18.68	17.70	17.85		14.86	15.64	9.30	11.52	16.17	11.65
<i>Ap. pseudoantipai</i>	18.68	19.34	17.52	18.68	20.33	18.84	18.56	19.67	19.87		17.82	14.84	12.35	17.17	12.65
<i>All. zicsii</i>	18.84	20.33	19.17	19.83	18.02	19.34	19.73	19.17	17.55	18.02		14.14	14.52	15.78	12.65
<i>Ap. ictérica</i>	19.80	21.95	20.30	20.63	18.32	16.01	16.69	20.13	16.86	18.51	20.83		11.00	14.67	10.15
<i>All. satchelli</i>	17.69	19.50	18.02	14.71	17.36	18.84	18.56	18.18	17.38	18.84	19.34	16.36		12.88	9.18
<i>All. tiginosa</i>	16.53	17.52	16.36	18.02	17.85	18.35	17.89	18.02	17.88	17.69	17.85	17.19	15.04		13.81
<i>Ap. handlirschi</i>	17.16	19.31	17.49	17.99	18.65	19.64	18.03	18.15	20.17	18.35	20.99	20.13	18.02	16.36	

TABLE 3: Comparison of the main morphological characters of the four revised genera and its closest genera.

	<i>Alloblophora</i> Eisen, 1874	<i>Gatesoma</i> Qiu and Bouché, 1998	<i>Eophila</i> Rosa, 1893	<i>Cernovitovia</i> Omodeo, 1856	<i>Panontonia</i> Mršić and Šapkarev, 1988	<i>Vogesia</i> Martínez Navarro and Marchán, 2022	<i>Avelona</i> Qiu and Bouché, 2000	<i>Heraulitia</i> Martínez Navarro and Marchán, 2022	<i>Proselodrilus</i> Bouché, 1972 or tanilobous	<i>Cataladrilus</i> Qiu and Bouché, 1998
Length (mm)	30-150	68-192	100-360	42-250	40-151	50-75	40-80	50-75	35-335	25-220
N. segments	60-210	118-197	139-341	138-304	115-181	125-140	100-125	156-172	130-350	124-384
Pigmentation	Green (sometimes absent)	Pigmentary dots	Purplish- brown in bands or brown	Light grey to brown	Absent	Absent	Brown	Absent	Absent	Absent
Prostomium	Epilobous	Epilobous	Epilobous	Epilobous or prolobous	Epilobous	Epilobous	Epilobous	Epilobous	Epilobous, prolobous or tanilobous	Epilobous
First dorsal pore	(3/4), 4/5, (5/6), (6/7)	4/5-7/8	5/6	9/10, 10/11, 12/13, 13/14	4/5	4/5	8/9	4/5	4/5-6/7	4/5-11/12
Spermathecal pores	Simple in 7/8, 8/9, 9/10, 10/11	Simple in 9/10, 10/11	Paired in 9/10, 10/11	Position and number highly variable, from 9/10 to 19/20	Double or multiple in 9/10, 10/11	Simple or double in 9/10, 10/11	Simple in 9/10, 10/11	Simple in 10/11, 11/12, 12/13	Simple in 12/13, 13/14, and 14/15 or simple, double, or multiple in 14, 15 (19) 20-27 (34)	Simple in 9/10, 10/11
Clitellum	28-78	1/2 22-36	24-44	21-65	23-34	26-34	29-36	21-36	Variable	22-46
Tubercula pubertatis	30-78	27-26	29-40	21-62	28-34	27-33	(31) 32-35 (36)	27-29	Variable	28-45
Oesophageal hearts	6-11	5, 6-11, 12	6-11	5, 6-11, 12	6-11	6-11	6-11	6-11	6, 7-11	6, 7-11
Calciferous glands	10-14 with diverticula in 10	10-12 (14) with diverticula in 10	10-12 with diverticula in 10	10-13	10-14 with diverticula in 10	10-14 with diverticula in 10	10-14 with diverticula in 10	13-14	11-15 with diverticula in 11	11-15 with diverticula in 11
Crop	15-16	15-16	15-16	15-16	15-16	15-16	15-16	15-16	15-16	15-16 (17, 18)
Gizzard	17-18	17-18	17-19	17-19, 20	17-18	17-18	17-18	(17) 18	17-18	17 (18, 19)-18 (21)
Typhlosole	Bifid or trifid	Bifid	Trifid	Trifid	Trifid	Bifid	Bifid	Bifid	Bifid or multifid	Simple, bifid, trifid, or multifid
Seminal vesicles	9-12	(9), (10), 11, 12	9-12	9-12 or 11-12	11-12	11-12	11-12	11-12	11-12	11-12
Spermathecae	Simple in 7/8, 8/9, 9/10, 10/11	Simple, globular in 9, 10	Simple in 9/10, 10/11	Position and number highly variable, from 9/10 to 19/20.	Double or multiple in 10, 11	Simple or double in 10, 11	Simple in 10, 11	Globular in 11, 12, 13	Variable in (12/13), 13/14, 14/15	9/10, 10/11

TABLE 4: List of the species included in *Allolobophora* sensu stricto and its main morphological features.

	<i>All. chlorotica</i>	<i>All. burgondiae</i>	<i>All. icterica</i>	<i>All. molleri</i>	<i>All. moebii</i>	<i>All. dubiosa</i>
Length (mm)	30-80	75-80	70-90	150-200	60-125	126-240
N. segments	60-140	150-170	140-200	145-210	104-122	121-303
Pigmentation	Green (sometimes absent)	Absent/brownish	Absent	Green	Green	Absent
Prostomium	Epilobous	Epilobous	Epilobous	Epilobous	Epilobous	Epilobous
First dorsal pore	4/5 (5/6)	4/5	4/5	4/5 (5/6)	4/5 (5/6)	4/5
Spermathecal pores	8/9, 9/10, 10/11	8/9, 9/10, 10/11	(7/8), 8/9, 9/10, 10/11	7/8, 8/9, 9/10, 10/11	7/8, 8/9, 9/10, 10/11	(7/8), 8/9, 9/10, 10/11
Clitellum	(28) 29-37	(27) 28-38 (39)	(33) 34-43 (44)	(48) 49-59	(52) 53-61 (62)	(36) 38-46 (49)
Tubercula pubertatis	30-36	30-37	(35) 36-42 (43)	50-57	(55) 56-61	(43) 44-47 (48)
Oesophageal hearts	6-11	6-11	6-11	6-11	6-11	6-11
Calciferous glands	10-14 with diverticula in 10	10-14 with diverticula in 10	10-14 with diverticula in 10	10-12 with diverticula in 10	10-12 with diverticula in 10	10-14 with diverticula in 10
Crop	15-16	15-16	15-16	15-16	15-16	15-16
Gizzard	17-18	17-18	17-18	17-18	17-18	17-18
Typhlosole	Bifid	Trifid	Bifid	Trifid	Trifid	Bifid
Nephridial bladders	U-shaped	U-shaped	U-shaped	U-shaped	U-shaped	U-shaped
Seminal vesicles	9, 10, 11, 12	9, 10, 11, 12	9, 10, 11, 12	9, 10, 11, 12	9, 10, 11, 12	9, 10, 11, 12
Spermathecae	9, 10, 11	9, 10, 11	9, 10, 11	7/8, 8/9, 9/10, 10/11	7/8, 8/9, 9/10, 10/11	8/9, 9/10, 10/11

moebii complex and closely related species is very controversial [36–38] and should be addressed by including the different species and subspecies in molecular phylogenetic analyses.

Genus *Panoniona* Mršić and Šapkarev, 1988

Type species: *Allolobophora leoni* Michaelsen, 1891

Material studied: *Panoniona satchelli*, 3 adult specimens; BOLD sample ID: ALL-013, ALL-014, ALL-015; Alsace, Haut-Rhin, Saint-Louis; latitude/longitude: 47.618/7.538; 04-Nov-2021; R. Della Vedova, T. Decaëns, D. Fernández Marchán leg. Specimen deposited in UCM-LT.

Diagnosis:

Small-to-medium-sized Lumbricini with absence of pigmentation. Prostomium is epilobous. Setae are closely paired. First dorsal pore in 4/5. Male pores in 1/2 15 with well-developed porophores. Spermathecal pores in 9/10, 10/11, double or multiple. Nephridial pores aligned or irregularly distributed (*en solfège*). Clitellum between 23 and 34. *Tubercula pubertatis* between 28 and 34. Oesophageal hearts in 6-11. Calciferous glands in 10-14 with diverticula in 10. Crop in 15-16. Glizzard in 17-18. Glandular papillae in 11-13, 16, (17), 26-29, and (34). Typhlosole is trifid or with two large and two small lamellae. Spermathecae double or multiple in 10, 11, globular and intracelomic. Two pairs of seminal vesicles in 11 and 12. Nephridial bladders are J-shaped with or without loop.

Differential diagnosis:

Panoniona can be differentiated from *Allolobophora* by the more anterior position of the clitellum and absence of pigmentation vs. frequently green pigmentation, repetition of spermathecae (double or multiple vs simple), number of seminal vesicles (two pairs in 11 and 12 vs. four pairs in 9, 10, 11, and 12), and shape of nephridial bladders (J-shaped vs. U-shaped).

Even though *Cernosvitovia* as redefined by Popovic et al. [10] is a morphologically heterogeneous genus, *Panoniona* can be distinguished from it by the type of spermathecae (double or multiple vs. simple) and either the position of the male pore (1/2 15 vs. 25-30) or the position of the spermathecae (9/10 and 10/11 vs. 9/10, 10/11, 11/12, and (12/13)).

Panoniona is clearly separated from *Gatesona* by the absence of pigmentary dots in cephalic segments, absence of very conspicuous genital papillae between segment 9 and the end of the clitellum, shape of nephridial bladders (J-shaped vs. sigmoid), repetition of spermathecae (double or multiple vs. simple), and number of seminal vesicles (two pairs in 11 and 12 vs. three to four pairs in (9, 10), 11, and 12).

Panoniona differs from *Eophila* by its generally smaller size, its pigmentation (absent vs. purplish-brown in bands or brown), and repetition of spermathecae (double or multiple vs. simple; Table 3).

Species included the following:

(i) *Panoniona leoni* Michaelsen, 1891

(ii) *Panoniona satchelli* *stat. nov.* Bouché, 1972

Genus *Vosgesia* Martínez Navarro and Marchán, 2022

Type species: *Vosgesia zicsii* *stat. nov.* (Bouché, 1972)

Material studied: 3 adult specimens; BOLD sample ID: ALL-007, ALL-008; ALL-009; Bourgogne; Haute-Saône, Luxeuil-les-Bains, latitude/longitude: 47.836/6.359; 02-Nov-2021; R. Della Vedova, T. Decaëns, D. Fernández Marchán leg. Specimens deposited in UCM-LT.

Diagnosis:

Small-sized (50-75 mm and 0.2-0.29 gr) Lumbricini with less than 150 segments and absence of pigmentation. Prostomium is epilobous. Setae are closely paired. First dorsal pore in 4/5. Male pores in 1/2 15 with well-developed porophores. Spermathecal pores in 9/10, 10/11, simple or double. Nephridial pores aligned. Clitellum in 26-34. *Tubercula pubertatis* in 27-33, with maximum development in 30-33. Oesophageal hearts in 6-11. Calciferous glands in 10-14 with diverticula in 10. Crop in 15-16. Glizzard in 17-18. Glandular papillae in 13, 27, and 30-32. Typhlosole is bifid. Spermathecae simple or double in 10, 11, globular and intracelomic. Two pairs of seminal vesicles in 11 and 12. Nephridial vesicles are J-shaped (incurvate and reclinate).

Differential diagnosis:

Vosgesia shares with *Avelona* (their closest relatives) the position of spermathecae (9/10 and 10/11) and seminal vesicles (11 and 12), typhlosole shape (bifid), and several external character states, but it is clearly separated by the position of the dorsal pore (4/5 vs. 8/9), the position of the clitellum (26-34 vs. 29-36), the nephridial pore disposition (aligned vs. irregular), the type of spermathecae (simple or double vs. simple), and the shape of the nephridial bladders (U-shaped vs. digitoid). The complete diagnoses of both genera are listed in Table 3.

Other species included: none.

4.2. Evolutionary and Biogeographic Implications. Historically, the genus *Allolobophora* includes a large number of species that are often taxonomic unrelated, resulting in a genus with a wide distribution and high diversity (the second largest one within the Lumbricidae), but with weak evolutionary ties. Unfortunately, this has complicated evolutionary and biogeographic interpretations. The gradual implementation of molecular techniques has revolutionized the systematics of the genus, with the elevation of redefined *Cernosvitovia* [10] and *Gatesona* [9], previously considered subgenera of *Allolobophora*, to generic status. The present work has continued this trend by supporting the elevation of *Panoniona* to genus status and the assignment of two of the studied species to newly described genera.

The resulting monophyletic clade comprising the former *Allolobophora* *sensu stricto* (with *All. chlorotica* and *All. burgondiae*) and *Heraclescolex* (comprising *All. moebii* and *All. molleri*, *All. icterica*, and *All. dubiosa*) appears more coherent in its morphological variability and distribution than the loosely defined *Allolobophora* of old. Qiu and Bouché [16] already indicated the strong morphological similarity between *Allolobophora* *sensu stricto* and *Heraclescolex*; in spite of that, surprisingly, they proposed to place them in different tribes. Domínguez et al. [2] showed for the first time that *All. molleri*, *All. moebii*, and *All. dubiosa* are closely related taxa to *All. chlorotica* by molecular phylogenetic

methods. The presence of *All. burgondiae* and *All. icterica* within that clade reinforces the idea that *Allolobophora* and *Heraclescolex* are synonyms. An alternative interpretation of the topology of the phylogenetic trees might be that *Allolobophora* sensu stricto (with *All. chlorotica* and *All. burgondiae*) and *Heraclescolex* do not constitute the same genus but are very closely related, independent genera. Even though both scenarios are likely, their extremely similar morphology (including the widespread green pigmentation) advises to synonymize them. Their main difference, the orientation of the nephridial bladders, is a rather feeble taxonomic character within a strongly variable structure within genera [9, 24]. Examination of the remaining species of *Heraclescolex* and their inclusion in the molecular phylogenetic analysis could recover them within *Allolobophora*, which would further support our hypothesis.

Interestingly, the genotypes of both merged genera (*All. chlorotica* and *All. moebii*) constitute species complexes on opposite poles of the morphological variability spectrum.

Allolobophora chlorotica has been shown to comprise several cryptic lineages [39, 40], some of which are widespread or even cosmopolitan, while a few are endemic to Provence and the Alps [39, 41]. Moreover, several subspecies have been described, but their association with those genetic lineages is still unknown. The integration of both sources of information with the new phylogenetic frame is essential to understand their phylogeography and the geographic origin of the genus.

The *molleri-moebii* species complex is a particularly controversial group that has been assigned to the genera *Allolobophora*, *Eophila*, *Aporrectodea*, or *Heraclescolex*. Trigo et al. [37, 38] found specimens with intermediate character states with respect to the position and extension of the clitellum in a continuum between the previously described species, which blurred the boundary between them. Barros [36] concluded that *All. molleri* and *All. moebii* constitute a single species named *All. molleri*. However, these results were not accepted by Qiu and Bouché [8, 13, 14, 16], who described additional subspecies and closely related species (the Maghrebian *Heraclescolex kionionus* Qiu and Bouché, 1998, *Heraclescolex rifanus* Qiu and Bouché, 1998, and *Heraclescolex postsellis* Qiu and Bouché, 1998). Our results appear to support the former view, as *All. molleri* and *All. moebii* do not have sufficient genetic divergence to be considered separate species. In order to solve the decade-old riddle of the taxonomy of this species complex, a systematic sampling of different populations of *All. molleri*, *All. moebii*, and the other species within this complex (*Allolobophora fernandae* Graff, 1957, *Allolobophora monchicana* Trigo, Mascates, Briones and Diaz Cosin, 1990, and *Allolobophora opisthosellata* Graff, 1961) will be performed in the near future; this will allow integrative species delimitation and will shed light on the evolutionary significance of the extraordinary clitellar variability of this group.

Although several questions remain to be solved, our current knowledge allows us to propose a rough scenario for the evolutionary origin and historical biogeography of *Allolobophora*. All the species assigned to the newly defined *Allolobophora* (plus the currently unstudied *Heraclescolex* species)

are native to the western Iberian Peninsula, France, Switzerland, Italy, and central-eastern Europe. The center of origin could be near the western Alps, as suggested by the presence of “endemic” lineages of *All. chlorotica* as well as *All. burgondiae* and *All. icterica* (Figure 2). The less intuitive distribution corresponds to the *All. moebii-molleri* complex and *Heraclescolex* species, which show a remarkable disjunction. However, the highest diversity of species and subspecies in southwest of the Iberian Peninsula, Morocco, and Algeria suggests an origin of this species group in the AlKaPeCa (Alboran-Kabylian-Peloritan-Calabrian) terrane, whose fragments migrated to their current positions in the Iberian Peninsula and northern Africa [42]. This terrane showed continuity with alpine Corsica and the developing Alps, allowing the dispersal and diversification of the ancestors of the extant *Allolobophora* species to their current range. An increased taxonomic and phylogeographic sampling of the areas and species involved would allow to refine and test this hypothesis, offering valuable insights into the link between the paleogeography of the western Mediterranean and its diversity.

The small, isolated genera *Heraultia* and *Vosgesia* and the occurrence of *Panoniona* in northeastern France and Switzerland also provide interesting biogeographic and evolutionary insights (Figure 2). *Heraultia tiginosa* represents a phylogenetic and biogeographic link between the French *Zophoscolex* and *Ethnodrilus* and the mostly Pyrenean *Proselodrilus* and *Cataladrilus*. While the latter two are remarkably diverse genera, *Heraultia* comprises a single species known from a single location: this suggests that *Heraultia* is a relict genus whose diversity and range might have been reduced by environmental and faunistic changes (as inferred for *Galiciandrilus*, *Compostelandrilus*, and *Vindoboscolex* [15]). Although new species belonging to *Heraultia* might be discovered in the future extending its distribution area, its presence highlights the high conservation value of Herault (and the area around Montpellier) as a hotspot of Mediterranean earthworm diversity.

Similarly, *V. zicsii* appears as an isolated distant relative of *Av. ligra* and *Gatesona* and the most northeastern representative of their clade (Figure 1). Its presence attests to a wider distribution of such lineage in the past and putatively to a greater diversity that is no longer present in France. *Panoniona satchelli* is isolated from its eastern relative *P. leoni* (present in Italy, Slovenia, Croatia, and further east) by the Alps (and about 400 km in a straight line) (Figure 2). In the case of *Panoniona*, the cause of such disjunction could indeed be the progressive elevation of the mountain chain, with an associated change in climate and vegetation, which could have led to a large unsuitable area between the current species. However, there is another cause that could explain the present range of *Vosgesia zicsii* and *P. leoni*. As stated by Bouché [4], endemic species in France appear to be restricted to the south of the Loire River: towards the North, they would have been eliminated by the presence of permafrost during the Last Glacial Maximum (LGM, ca. 33,000-15,000 years ago [43]). Even though most recent models suggest that permafrost would have not been present south of 49° N in France [44], the dominant




FIGURE 2: Sampling localities of the species studied in this work; for all species except *Allobophora icterica*, this corresponds to their known ranges. The main ranges of the closest relatives according to phylogenetic analyses are shown as coloured shapes.

biome in most of France and the rest of Europe would have been steppe-tundra [45] which is unsuitable for most lumbricid earthworms. The almost complete absence of endemic earthworm species (and low diversity) in Germany, Czech Republic, Poland, Slovakia, and countries further north supports this hypothesis. Nevertheless, Bouché [4] noted the presence of a few strictly endemic species in a small region (Val de Saône-Vosges) within the “glaciated” area. *Allobophora burgondiae*, *P. satchelli*, *V. zicsii*, and *Ap. pseudoantipai* (as well as *Ap. gogna* Bouché, 1972 and *Ap. velox* (Bouché 1967)) occur there. Thus, the isolation of these species relative to their closest southern or eastern relatives would be explained by the disappearance of intermediate species during the LGM or other unsuitable climatic events. Interestingly, some models show that the aforementioned region roughly corresponds to the presumed distribution of different vegetation types (such as ombrotrophic bogs) during the LGM [46], which supports the role of the Val de Saône-Vosges area as a glacial refuge for endemic Lumbricidae. Further efforts should be devoted to biodiversity assessment and conservation of this unique hotspot, which has been generally neglected in favour of the southern regions.

The biogeographic inferences drawn from *Allobophora* and the relict species from northeastern France illustrate the relevance of the Alpine region for the evolution of several lineages of Lumbricidae, with the effects of the LGM likely disrupting the remaining evidence. This is probably also true for other genera such as *Aporrectodea* sensu stricto, *Lumbricus* Linnaeus, 1758 or *Octodrilus* Omodeo, 1956. A greater focus on the implementation of phylogeography and molecular phylogenetics to the earthworm faunas of Switzerland, Italy, Austria, Slovenia, Croatia, and Serbia would greatly enhance our understanding of the evolution and historical biogeography of Lumbricidae.

5. Conclusions

Molecular phylogenetic analyses revealed that the representatives of *Allobophora* sensu stricto and *Heraclodescolex* belong to a clearly delimited clade, which suggest their synonymy. On the other hand, other species formerly assigned to *Allobophora* were recovered within *Panonionia*, or as isolated, genus-level branches. Therefore, the revised definitions for *Allobophora* and *Panonionia* were provided, and the new genera *Heraulitia* and *Vosgesia* were described.

Although several species of *Heraclescolex* remain to be included in this molecular phylogenetic framework, it is likely that most of them will be nested within the newly defined *Allolobophora*. If so, the synonymy between both genera would be proven.

Systematic revision of the aforementioned earthworm groups revealed evolutionarily more coherent groups that provide information on their diversification and historical biogeography. The Maghreb and the areas around the Alps were highlighted as relevant targets for expanding our knowledge on this topic, and the importance of the Val de Saône-Vosges (northeastern France) as a glacial refugium with relict earthworm lineages was confirmed.

Data Availability

DNA sequences obtained in this study are available in GenBank with their corresponding accession numbers. COI sequences, together with metadata, are publicly available in BOLD.

Conflicts of Interest

The authors declare that there are no conflicts of interest or personal relationships that could have influenced the work reported in this article.

Acknowledgments

This work was supported by subproject Fauna Ibérica XII. Oligochaeta, Lumbricoidea: Lumbricidae, Hormogastridae #PGC2018-095851-B-C66 from the Spanish Ministry of Sciences, Innovation and Universities. Daniel Fernández Marchán was funded by a María Zambrano Postdoctoral Grant from the Spanish Ministry of Sciences, Innovation and Universities and by a Make Our Planet Great Again Postdoctoral grant from Campus France. Marta Novo was supported by Ramón y Cajal Fellowship (RYC2018-024654-I), and this study was funded by Grant PID2021-122243NB-I00, both from MCIN/AEI/10.13039/501100011033/FEDER, UE. We are thankful to Raphaël Della Vedova for his support in the sampling campaign.

Supplementary Materials

Additional information can be found in the online version of the article on the publisher's website. Table S1: list of species included in the phylogenetic analyses with reference to the original publication from which the sequences were taken. Figure S1: phylogenetic tree estimated by maximum likelihood, based on the concatenated sequence of COI, 16S, 28S, 12S, and ND1. Figure S2: complete Bayesian inference of the phylogenetic relationships of the studied taxa and some Lumbricidae representatives. (*Supplementary Materials*)

References

- [1] C. Csuzdi and A. Zicsi, *Earthworms of Hungary (Annelida: Oligochaeta, Lumbricidae)*, Hungarian Natural History Museum, Budapest, Hungary, 2003.
- [2] J. Dominguez, M. Aira, J. W. Breinholt, M. Stojanovic, S. W. James, and M. Perez-Losada, "Underground evolution: new roots for the old tree of lumbricid earthworms," *Molecular Phylogenetics & Evolution*, vol. 83, pp. 7–19, 2015.
- [3] V. Pop, "Zur phylogenie und Systematik der Lumbriciden," *Zoologische Jahrbücher Abteilung für Systematik Ökologie und Geographie der Tiere*, vol. 74, pp. 487–522, 1941.
- [4] M. B. Bouché, *Lombriciens de France. Ecologie et systématique* (vol. 72), Institut National de la Recherche Agronomique, Paris, France, 1972.
- [5] N. Mršić and J. A. Šapkarev, "Revision of the genus *Allolobophora* Eisen 1874 (sensu Pop 1941) (Lumbricidae, Oligochaeta)," *Prirodonaučen muzej na Makedonija*, vol. 19, pp. 1–38, 1988.
- [6] P. Omodeo, "Oligocheti dell'Indochina e del Mediterraneo Orientale," *Memorie del Museo Civico di Storia Naturale di Verona*, vol. 5, pp. 321–336, 1956.
- [7] T. Perel, "A critical analysis of the Lumbricidae genera system (with key to the USSR fauna genera)," *Révue d'Écologie et Biologie du Sol*, vol. 13, no. 4, pp. 635–643, 1976.
- [8] J. Qiu and M. Bouché, "Le genre *Allolobophora* Eisen, 1874 (Oligochaeta Lumbricidae), ses avatars et sa définition moderne," *Documents pédozoologiques et intégrologiques*, vol. 4, pp. 86–97, 1998.
- [9] D. F. Marchán, S. Jiménez, T. Decaëns, and J. Domínguez, "Systematic revision of *Gatesona* (Crassiclitellata, Lumbricidae), an endemic earthworm genus from the Massif Central (France)," *PLoS One*, vol. 16, no. 9, article e0255978, 2021.
- [10] F. J. Popović, M. M. Stojanović, J. Domínguez, J. M. Sekulić, T. B. Trakić, and D. F. Marchán, "Molecular analysis of five controversial Balkanic species of *Allolobophora* (sensu lato) Eisen, 1873 (Lumbricidae, Clitellata) with emendation of the genus *Cernosvitovia* Omodeo, 1956," *Zootaxa*, vol. 5116, no. 3, pp. 351–372, 2022.
- [11] I. de Sosa, D. J. Díaz Cosín, C. Csuzdi, M. G. Paoletti, and D. F. Marchán, "Placing *Eophila tellinii* (Oligochaeta, Lumbricidae) in a molecular phylogenetic context advances the century-old controversy around the problematic genus," *European Journal of Soil Biology*, vol. 94, article 103114, 2019.
- [12] M. G. Paoletti, R. J. Blakemore, C. Csuzdi et al., "Barcoding *Eophila crodabepis* sp. nov. (Annelida, Oligochaeta, Lumbricidae), a large stripy earthworm from alpine foothills of north-eastern Italy similar to *Eophila tellinii* (Rosa, 1888)," *PLoS One*, vol. 11, no. 3, article e0151799, 2016.
- [13] J. Qiu and M. Bouché, "*Heraclescolex*, un nouveau genre de Lumbricidae (Annelida: Oligochaeta)," *Documents pédozoologiques et intégrologiques*, vol. 4, pp. 140–152, 1998.
- [14] J. Qiu and M. Bouché, "Contribution à la taxonomie des Avelonini trib. nov. (Oligochaeta: Lumbricidae). *Avelona* gen. nov., *Koinodrilus* gen. nov et *Nicodrilus cuendeti* sp. nov.," *Documents pédozoologiques et intégrologiques*, vol. 4, pp. 109–116, 1998.
- [15] D. F. Marchán, C. Csuzdi, T. Decaëns, T. Szederjesi, V. Pizl, and J. Domínguez, "The disjunct distribution of relict earthworm genera clarifies the early historical biogeography of the Lumbricidae (Crassiclitellata, Annelida)," *Journal of Zoological Systematics and Evolutionary Research*, vol. 59, no. 8, pp. 1703–1717, 2021.
- [16] J. Qiu and M. Bouché, "Révision des taxons supraspécifiques de Lumbricoidea," *Documents pédozoologiques et intégrologiques*, vol. 3, pp. 179–216, 1998.

- [17] J. Domínguez, M. Aira, P. G. Porto, D. J. Díaz Cosín, and M. Pérez-Losada, "Multigene phylogeny reveals two new isolated and relic earthworm genera (Oligochaeta: Lumbricidae)," *Zoological Journal of the Linnean Society*, vol. 182, no. 2, pp. 258–274, 2018.
- [18] M. Pérez-Losada, M. Ricoy, J. C. Marshall, and J. Domínguez, "Phylogenetic assessment of the earthworm Aporrectodea caliginosa species complex (Oligochaeta: Lumbricidae) based on mitochondrial and nuclear DNA sequences," *Molecular Phylogenetics and Evolution*, vol. 52, no. 2, pp. 293–302, 2009.
- [19] O. Folmer, M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek, "DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates," *Molecular Marine Biology and Biotechnology*, vol. 3, no. 5, pp. 294–299, 1994.
- [20] K. Katoh and D. M. Standley, "MAFFT multiple sequence alignment software version 7: improvements in performance and usability," *Molecular Biology and Evolution*, vol. 30, no. 4, pp. 772–780, 2013.
- [21] T. A. Hall, "Bio edit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT," *Nucleic Acid Symposium Series*, vol. 41, pp. 95–98, 1999.
- [22] M. Pérez-Losada, J. W. Breinholt, P. G. Porto, M. Aira, and J. Domínguez, "An earthworm riddle: systematics and phylogeography of the Spanish Lumbricid Postandrilus," *PLoS One*, vol. 6, no. 11, article e28153, 2011.
- [23] M. Pérez-Losada, J. W. Breinholt, M. Aira, and J. Domínguez, "An updated multilocus phylogeny of the Lumbricidae (Annelida: Clitellata: Oligochaeta) earthworms," *Journal of Phylogenetics & Evolutionary Biology*, vol. 3, no. 1, 2015.
- [24] F. Bozorgi, M. Seiedy, M. Malek, M. Aira, M. Perez-Losada, and J. Domínguez, "Multigene phylogeny reveals a new Iranian earthworm genus (Lumbricidae: *Philomontanus*) with three new species," *PLoS One*, vol. 14, no. 1, article e0208904, 2019.
- [25] S. Jiménez Pinadero, D. F. Marchán, M. Novo, D. Trigo, J. Domínguez, and D. J. Díaz Cosín, "Sorry atlanticus, you are not my type: molecular assessment splits *Zophoscolex* (Lumbricidae: Crassicitellata) into French and Iberian genera," *Zoological Journal of the Linnean Society*, vol. 194, no. 3, pp. 726–735, 2022.
- [26] D. F. Marchán, T. Decaëns, D. J. Díaz Cosín, M. Hedde, E. Lapied, and J. Domínguez, "French Mediterranean islands as a refuge of relic earthworm species: *Cataladrilus porquerolensis* sp. nov. and *Scherotheca portrosana* sp. nov. (Crassicitellata, Lumbricidae)," *European Journal of Taxonomy*, no. 701, pp. 1–22, 2020.
- [27] D. Darriba, G. L. Taboada, R. Doallo, and D. Posada, "jModelTest 2: more models, new heuristics and parallel computing," *Nature Methods*, vol. 9, no. 8, pp. 772–772, 2012.
- [28] H. Akaike, "Information theory and an extension of the maximum likelihood principle," in *Proceedings of the 2nd International Symposium on Information, bn petrow, f. Czaki Akademiai Kiado*, pp. 267–281, Budapest, Hungary, 1973.
- [29] G. Schwarz, "Estimating the dimension of a model," *The Annals of Statistics*, vol. 6, no. 2, pp. 461–464, 1978.
- [30] F. Ronquist, M. Teslenko, P. van der Mark et al., "MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space," *Systematic Biology*, vol. 61, no. 3, pp. 539–542, 2012.
- [31] M. A. Miller, W. Pfeiffer, and T. Schwartz, "Creating the CIPRES science gateway for inference of large phylogenetic trees," in *2010 Gateway Computing Environments Workshop (GCE)*, pp. 1–8, New Orleans, LA, USA, 2010.
- [32] A. M. Kozlov, D. Darriba, T. Flouri, B. Morel, and A. Stamatakis, "RAXML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference," *Bioinformatics*, vol. 35, no. 21, pp. 4453–4455, 2019.
- [33] K. Tamura, G. Stecher, and S. Kumar, "MEGA11: molecular evolutionary genetics analysis version 11," *Molecular Biology and Evolution*, vol. 38, no. 7, pp. 3022–3027, 2021.
- [34] D. F. Marchán, S. W. James, A. R. Lemmon, E. M. Lemmon, M. Novo, and J. Domínguez, "A strong backbone for an invertebrate group: anchored phylogenomics improves the resolution of genus-level relationships within the Lumbricidae (Annelida, Crassicitellata)," *Organisms Diversity & Evolution*, vol. 22, no. 4, pp. 915–924, 2022.
- [35] A. R. Lemmon, S. A. Emme, and E. M. Lemmon, "Anchored hybrid enrichment for massively high-throughput phylogenomics," *Systematic Biology*, vol. 61, no. 5, pp. 727–744, 2012.
- [36] F. Barros, *Estudio Cariológico y Alozimico del Complejo Allohophora molleri* (Lumbricidae, Oligochaeta), [Ph.D. thesis], Universidade de Santiago de Compostela, 1992.
- [37] D. Trigo, R. Mascato, M. I. Briones, and D. Diaz Cosin, "Lombrices de tierra de Portugal continental. Inventario y citas," *Arquivos do Museu Bocage*, vol. 1, pp. 521–567, 1990.
- [38] D. Trigo, S. Mato, B. F. Souto, and D. J. Diaz Cosin, "Earthworms of continental Portugal. Relationships with soil factors," *Italian Journal of Zoology*, vol. 56, no. 4, pp. 327–331, 1989.
- [39] L. Dupont, F. Lazrek, D. Porco et al., "New insight into the genetic structure of the *Allohophora chlorotica* aggregate in Europe using microsatellite and mitochondrial data," *Pedobiologia*, vol. 54, no. 4, pp. 217–224, 2011.
- [40] R. A. King, A. L. Tibble, and W. O. C. Symondson, "Opening a can of worms: unprecedented sympatric cryptic diversity within British lumbricid earthworms," *Molecular Ecology*, vol. 17, no. 21, pp. 4684–4698, 2008.
- [41] D. F. Marchán, M. Hedde, E. Lapied et al., "Contrasting phylogeographic patterns of earthworms (Crassicitellata, Lumbricidae) on near-shore Mediterranean islands," *European Journal of Soil Biology*, vol. 101, article 103242, 2020.
- [42] R. Leprêtre and D. F. Lamotte, "The Tell-Rif orogenic system (Morocco, Algeria, Tunisia) and the structural heritage of the southern Tethys margin," *BSGF-Earth Sciences Bulletin*, vol. 189, no. 2, p. 10, 2018.
- [43] P. U. Clark, A. S. Dyke, J. D. Shakun et al., "The last glacial maximum," *Science*, vol. 325, no. 5941, pp. 710–714, 2009.
- [44] K. H. Stadelmaier, P. Ludwig, P. Bertran et al., "A new perspective on permafrost boundaries in France during the last glacial maximum," *Climate of the Past*, vol. 17, no. 6, pp. 2559–2576, 2021.
- [45] N. Ray and J. Adams, "A GIS-based vegetation map of the world at the last glacial maximum (25,000–15,000 BP)," *Internet Archaeology*, no. 11, 2001.
- [46] V. Janská, B. Jiménez-Alfaro, M. Chytrý et al., "Palaeodistribution modelling of European vegetation types at the last glacial maximum using modern analogues from Siberia: prospects and limitations," *Quaternary Science Reviews*, vol. 159, pp. 103–115, 2017.