

We investigated the effect of the excitatory amino acid (EAA) receptor agonists L-glutamate, N-methyl-D-aspartate (NMDA), (RS)-a-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) and kainic acid on KCl-induced contractions of rabbit tracheal smooth muscle, as well as the role of epithelium and endogenously produced nitric oxide and prostaglandins on these responses. L-Glutamate decreased KCl-induced contractions up to 30%. This effect was attenuated by epithelium removal, tetrodotoxin, methylene blue and indomethacin but not by *N*<sup>G</sup>-nitro-L-arginine methyl ester. While NMDA, AMPA and kainic acid had no effect, the combination of NMDA + kainic acid decreased KCl-induced contractions. These results suggest that, in rabbit trachea, L-glutamate has, at least in part, an epithelium-dependent effect mediated via prostaglandin formation and that the EAA receptors involved are non-classical.

**Key words:** Airway smooth muscle, Rabbit, L-Glutamate, Prostaglandins, EAA receptors

## Epithelium-dependent effect of L-glutamate on airways: involvement of prostaglandins

Apostolia A. Hatziefthimiou<sup>CA</sup>, Konstantinos I. Gourgoulianis and Paschalidis-Adam Molyvdas

Department of Physiology, Medical School,  
University of Thessaly, Papakiriazi 22, 412 22  
Larissa, Greece

<sup>CA</sup>Corresponding Author

Tel: + 30 41 565005

Fax: + 30 41 565068

E-mail: axatzi@med.uth.gr

## Introduction

In humans, glutamate (in the form of monosodium) could be a provoking factor involved in asthma attacks, although the existence of monosodium glutamate-induced asthma has not been established.<sup>1</sup> Data obtained from animal studies support the hypothesis that the intense exposure to glutamate can be toxic for lungs because the over-activation of excitatory amino acid (EAA) receptors, mainly of the N-methyl-D-aspartate (NMDA) type, is involved in the pathogenesis of a variety of lung lesions like acute injury,<sup>2</sup> pulmonary edema,<sup>3</sup> oxidant tissue injury,<sup>4</sup> airway hyper-responsiveness and inflammation.<sup>5,6</sup> In animal lungs, accumulating data suggest that the localization of EAA receptors are presumably neuronal and L-glutamate exerts its effect via NMDA<sup>2,3</sup> or via non-classical EAA receptors.<sup>7</sup>

The purpose of the present study was first to investigate the effect of main ionotropic EAA receptors agonists such as L-glutamate, NMDA, (RS)-a-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) and kainic acid on KCl-induced contractions of rabbit tracheal smooth muscle. Second, we examined the involvement of epithelium and the endogenous formation of nitric oxide (NO) and prostaglandins in the effect of L-glutamate.

## Materials and methods

Adult rabbits (1–2 kg body weight) of either sex were sacrificed by an overdose of intravenously

administered sodium thiopentone (Abbot, Italy). Exothoracic tracheal tissue was removed and placed in Krebs solution (pH 7.4 at 37°C) with the following composition: Na<sup>+</sup>, 137 mM; Mg<sup>2+</sup>, 1.1 mM; K<sup>+</sup>, 5.9 mM; Cl<sup>-</sup>, 123.0 mM; Ca<sup>2+</sup>, 2 mM; H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, 1.2 mM; HCO<sub>3</sub><sup>-</sup>, 24.9 mM; and glucose, 9.6 mM. The solution was gassed with 95% O<sub>2</sub> and 5% CO<sub>2</sub>.

After the removal of connective tissue, tracheal rings were dissected from tracheas under an SZ30 Olympus stereoscope (Japan). The tracheal strips were 2 mm in width and were cut longitudinally through the cartilage opposite the smooth muscle layer. In experiments with epithelium-denuded tracheal rings, the epithelial layer was removed with a cotton-tipped applicator. Each strip was placed with the superfused luminal side up in a water-jacketed organ bath. One end of the tracheal strip was fixed on the bottom of the organ bath and the other to the transducer tip. The entire strip was continuously perfused with oxygenated Krebs solution at 37°C. The volume of the organ bath was approximately 3 ml and the perfusion rate was 5 ml/min. Tracheal strips were stretched manually to 1 g resting tension and were allowed to equilibrate in the organ bath for at least 60 min. Before starting the experiments, the rings were contracted by elevating the extracellular [K<sup>+</sup>]<sub>0</sub> concentration to 80 mM or by 10<sup>-5</sup> M acetylcholine (ACh) until a constant and reproducible contraction was achieved. Elevating potassium solutions were made by isosmotic substitution with [Na<sup>+</sup>]<sub>0</sub>. The concentration of 10<sup>-5</sup> M ACh was chosen

because, in pilot studies,  $10^{-5}$  M ACh produced 50% of the maximal response (data not shown). Changes in tension were recorded on a Grass FT03C force displacement transducer (Astro Med Inc., USA) and displayed via a Universal oscillograph (Harvard Apparatus, England) recorder. To investigate the effect of L-glutamate on ACh-induced or KCl-induced contractions, increasing concentrations of L-glutamate (range,  $10^{-9}$  to  $10^{-3}$  M) were added in the perfusion medium and, after 15 min, contractions were induced by ACh or KCl. The effect of the main ionotropic EAA receptors agonists NMDA, AMPA and kainic acid (range,  $10^{-9}$  to  $10^{-4}$  M) on KCl-induced contractions was studied in the same way. We also studied the effect of kainic acid (range,  $10^{-9}$  M to  $10^{-4}$  M), in the presence of  $10^{-6}$  M NMDA and the effect of the NMDA (range,  $10^{-9}$  M to  $10^{-4}$  M) in the presence of  $10^{-6}$  M kainic acid. The experiments with NMDA were performed in the presence of  $Mg^{2+}$  and also in  $Mg^{2+}$ -free Krebs solution.

Experiments with epithelium-intact and epithelium-denuded preparations were carried out in parallel. In experiments in which tetrodotoxin (TTX), methylene blue, *N*<sup>G</sup>-nitro-L-arginine methyl ester (L-NAME) and indomethacin were used, the rings were incubated with the inhibitor for 45 min before the exposure to L-glutamate. Contractions were evoked by 80 mM KCl.

Values are expressed as a percentage of reference contraction induced by 80 mM KCl or  $10^{-5}$  M ACh. All data are presented as mean  $\pm$  standard error (SE), and *N* refers to the number of animals. Statistical differences between all the responses were assessed using paired or unpaired *t*-tests, and  $p < 0.05$  was considered to be significant.

ACh, L-glutamate, NMDA, AMPA, kainic acid, TTX, methylene blue, L-NAME and indomethacin were all obtained from Sigma (Germany).

## Results

L-Glutamate evoked a significant concentration-dependent decrease of KCl-induced contractions (Fig. 1). The maximal decrease (30%) was observed in the presence of  $10^{-3}$  M L-glutamate ( $p < 0.001$ , paired *t*-test; Fig. 1). The removal of epithelium, on the contrary, reduced significantly the effect of L-glutamate ( $p < 0.05$ , paired *t*-test; Fig. 1). L-Glutamate had no effect on ACh-induced contractions (Fig. 2). The presence of TTX ( $3 \times 10^{-6}$  M) in the perfusion medium significantly attenuated the effect of L-glutamate on KCl-induced contractions ( $p < 0.01$ , paired *t*-test, for concentrations of L-glutamate between  $10^{-8}$  and  $10^{-3}$  M; Fig. 3). The treatment of preparations with methylene blue ( $10^{-5}$  M) attenuated significantly the effect of L-glutamate on KCl-induced contractions ( $p < 0.01$ , unpaired *t*-test; Fig. 4) and had




FIG. 1. L-Glutamate effect on epithelium-intact (●) and epithelium-denuded preparations (○). The responses are mean  $\pm$  SE (*n* = 5) and are plotted as a percentage of the reference response before the exposure to L-glutamate. \* $p < 0.05$ , \*\*\* $p < 0.001$ , paired *t*-test.




FIG. 2. L-Glutamate effect on ACh-induced contractions (●), and in the presence of methylene blue (○). The responses are mean  $\pm$  SE (*n* = 6) and are plotted as a percentage of the reference response before the exposure to L-glutamate.




FIG. 3. The effect of L-glutamate on KCl-induced contractions (●), and in the presence of TTX (○). The responses are mean  $\pm$ SE ( $n = 6$ ) and are plotted as a percentage of reference response before the exposure to L-glutamate.  $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , paired t-test.




FIG. 5. The effect of L-glutamate (●), AMPA (○), kainic acid (□), NMDA (■) and NMDA in magnesium-free medium (▲). The responses are mean  $\pm$ SE;  $n = 11$  for L-glutamate, and  $n = 5$  for each other agonist. The responses are plotted as a percentage of the reference response before the exposure to agonist.




FIG. 4. The effect of L-glutamate on KCl-induced contractions (●), and in the presence of methylene blue (○), L-NAME (▲) and indomethacin (△). The responses are mean  $\pm$ SE;  $n = 11$  for experiments with L-glutamate,  $n = 7$  for experiments in the presence of methylene blue,  $n = 5$  for experiments in the presence of L-NAME, and  $n = 6$  for experiments in the presence of indomethacin. The responses are plotted as a percentage of the reference response before the exposure to L-glutamate. \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$ , unpaired t-test.




FIG. 6. The effect of NMDA in the presence of  $10^{-6}$  M kainic acid (▲), of kainic acid in the presence of  $10^{-6}$  M NMDA (■), and in magnesium-free medium (○). The responses are mean  $\pm$ SE ( $n = 6$ ) and are plotted as a percentage of the reference response before the exposure to agonists. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  paired t-test.

a small, statistically not significant, potentiating effect on contractions evoked by ACh (Fig. 2). The presence of L-NAME ( $10^{-4}$  M) in the perfusion medium did not influence the effect of L-glutamate (Fig. 4), while the presence of indomethacin ( $10^{-5}$  M) significantly reduced the effect of L-glutamate ( $p < 0.05$ , unpaired *t*-test; Fig. 4).

Finally, NMDA, in the presence or absence of Mg<sup>2+</sup> in the perfusion medium, AMPA or kainic acid had no effect on KCl-induced contractions (Fig. 5). However, the combination of NMDA + kainic acid (NMDA in the presence of  $10^{-6}$  M kainic acid or kainic acid in the presence of  $10^{-6}$  M NMDA) evoked a significant decrease of KCl-induced contractions (Fig. 6). The absence of Mg<sup>2+</sup> from the perfusion medium had no effect on these responses (Fig. 6). The combined effect of the two ionotropic EAA agonists, kainic acid and NMDA, was not as potent as that by L-glutamate alone. L-Glutamate, NMDA, AMPA or kainic acid in concentrations already mentioned had no effect on tracheal strips under resting conditions and in the absence of KCl-induced contractile activity (data not shown).

## Discussion

Our results demonstrate that L-glutamate application on rabbit trachea decreases KCl-induced contractions, without affecting ACh-induced contractions of smooth muscle. KCl in high concentration induces contractions in rabbit trachea mainly via the ACh release;<sup>8</sup> therefore, we assume that L-glutamate, at the dose tested, had no post-junctional effect. The effect of L-glutamate on rabbit tracheal smooth muscle was attenuated by the removal of epithelium, as well as by the presence of TTX, methylene blue or indomethacin in the perfusion medium. These data provide evidence, first, for the release of a neuronal agent that mediates the L-glutamate effect and, second, for an epithelium-dependent and prostaglandin-dependent way of action.

In perfused rat lungs, activation of NMDA receptors, in the presence of the NO precursor L-arginine, stimulates cGMP formation and synthesis of NO.<sup>2,3</sup> To investigate the involvement of cGMP and/or NO in the relaxant effect of L-glutamate, we treated our preparations with methylene blue, an inhibitor of cGMP formation. While methylene blue significantly reduced the effect of L-glutamate, the treatment of our preparations with L-NAME, an inhibitor of NO synthase, had no effect. These results suggest that NO was not involved in the L-glutamate effect on rabbit trachea. Previous studies reported that methylene blue might exert its effect via a mechanism independent of the inhibition of cGMP formation, and may inhibit the synthesis and/or the release of prostaglandins.<sup>9–11</sup> For this reason, we treated our prepara-

tions with indomethacin, an inhibitor of cyclooxygenase. The presence of indomethacin in the perfusion medium attenuated the effect of L-glutamate, thus suggesting that L-glutamate exerts its effect, at least in part, via prostaglandin formation.

Although our results provide evidence for a neuronal localization of L-glutamate receptors, their possible site of localization is not clear. L-Glutamate receptor subunits are present on airways.<sup>12,13</sup> According to the available data, the L-glutamate receptors, mainly of NMDA type,<sup>14</sup> are located on parasympathetic nerves.<sup>15</sup> In guinea pigs, indirect evidence also support the existence of L-glutamate receptors on airways sensory neurons, where they mediate the acute response to capsaicin.<sup>5</sup> Thus, in rat bronchi, activation of glutamate receptors enhances cholinergic contractions.<sup>7</sup> This effect may be mediated indirectly by the release of sensory peptides, because sensory peptides may facilitate cholinergic neurotransmission in airways.<sup>16–18</sup>

In contrast to these previous observations, our findings demonstrate a relaxant effect of L-glutamate in rabbit trachea. The discrepancy may be due to species and/or tissue differences. Supporting this, L-glutamate has no effect on the responsiveness of rabbit airways.<sup>19</sup> In addition, guinea pig airways contain significantly greater amounts of substance P<sup>20</sup> and are more sensitive to NK-1 and NK-2 receptors agonists than rabbit.<sup>21</sup> Capsaicin induces plasma extravasation in guinea pig and rat trachea<sup>22–24</sup> but has no effect on rabbit trachea.<sup>25</sup> Concerning tissue differences, the density and the type of tachykinin receptors, as well as the tissue responsiveness to sensory peptides, alter from proximal to distal airways.<sup>26,27</sup> Considering these observations, L-glutamate receptors may exist on the sensory fibers of rabbit but their activation and the subsequent release of sensory peptides neither provoke direct contractions or enhance ACh release. Direct measurements are necessary to test this hypothesis.

The presence of the NMDA receptor subunit has also been demonstrated by immunocytochemistry in neurons of rat larynx and esophagus containing NO and vasoactive intestinal peptide (VIP),<sup>28</sup> from which the inhibitory nerves supplying airway smooth muscle originate.<sup>29</sup> The presence of a functional non-adrenergic non-cholinergic inhibitory (iNANC) system in rabbits has also been demonstrated,<sup>30,31</sup> but this system seems not to use NO as a neurotransmitter.<sup>32</sup> The other neurotransmitter of the iNANC system (VIP) has been detected in rabbit airways.<sup>33,34</sup> Although VIP exerts its effect via cAMP formation,<sup>35</sup> prostaglandins<sup>36</sup> as well as epithelium<sup>37,38</sup> may be involved in its effect. The release of VIP in rabbit trachea stimulated by L-glutamate could be a reasonable explanation, while L-glutamate failed to induce contractions since VIP has also a modulatory effect on ACh release.<sup>39</sup>

Previous observations in mammalian peripheral tissues,<sup>7,40-42</sup> as well as in airways,<sup>7</sup> suggest the existence of non-classical glutamate receptors. Studies in the central nervous system have reported glutamate receptor subtypes composed of both kainic acid or AMPA and NMDA receptor subunits,<sup>43</sup> or non-classical glutamate receptors blocked by both NMDA and non-NMDA antagonists.<sup>44</sup> In this respect, we tested the effect of the main EAA receptor agonists NMDA, AMPA and kainic acid as well as the combination of the NMDA plus kainic acid. Our results show that L-glutamate exerts its effect via a non-classical EAA receptor, since the classical EAA receptor agonists, NMDA, AMPA and kainic acid, were ineffective. However, the combination of NMDA + kainic acid decreased KCl-induced contractions. The role of Mg<sup>2+</sup> was probably not significant in our system, as the presence or absence of Mg<sup>2+</sup> did not affect the decrease of contraction evoked by kainic acid in the presence of NMDA in the bath medium. The effect of the combinations of EAA agonists tested in the present study was less potent than L-glutamate. Therefore, we cannot exclude the involvement of metabotropic receptors in this effect.

In conclusion, in rabbit trachea, L-glutamate in part has an epithelium-dependent effect, mediated via prostaglandin formation. The EAA receptors involved in this effect are non-classical EAA receptors of the NMDA/non-NMDA type.

## References

- Stevenson DD. Monosodium glutamate and asthma. *J Nutr* 2000; 130: 1067S-1073S.
- Said SI, Berisha HI, Pakbaz H. N-Methyl-D-aspartate receptors outside the central nervous system: activation causes acute lung injury that is mediated by nitric oxide synthesis and prevented by vasoactive intestinal peptide. *Neuroscience* 1995; 65: 943-946.
- Said SI, Berisha HI, Pakbaz H. Excitotoxicity in the lung: N-methyl-D-aspartate-induced, nitric oxide-dependent, pulmonary edema is attenuated by vasoactive intestinal peptide and by inhibitors of poly(ADP-ribose) polymerase. *Proc Natl Acad Sci USA* 1996; 93: 4688-4692.
- Said SI, Pakbaz H, Berisha HI, Raza SI. NMDA receptor activation: critical role in oxidant tissue injury. *Free Radic Biol Med* 2000; 28: 1300-1302.
- Said SI. Glutamate receptors and asthmatic airway disease. *Trends Pharmacol Sci* 1999; 20: 132-134.
- Said SI, Dickman KG. Pathways of inflammation and cell death in the lung: modulation by vasoactive intestinal peptide. *Regulat Pept* 2000; 93: 21-29.
- Aas P, Tanso R, Fonnum F. Stimulation of peripheral cholinergic nerves by glutamate indicates a new peripheral glutamate receptor. *Eur J Pharmacol* 1989; 164: 93-102.
- Loenders B, Jorens PG, Herman AG. Epithelial modulation of cholinergic responses in rabbit trachea is partly due to neutral endopeptidase activity. *Eur J Pharmacol* 1996; 296: 89-96.
- Martin W, Drazan KM, Newby AC. Methylene blue but not changes in cyclic GMP inhibits resting and bradykinin-stimulated production of prostacyclin by pig aortic endothelial cells. *Br J Pharmacol* 1989; 97: 51-56.
- Okamura T, Yoshida K, Toda N. Suppression by methylene blue of prostaglandin I<sub>2</sub> synthesis in isolated dog renal arteries. *J Pharmacol Exp Ther* 1990; 254: 198-203.
- Gao Y, Vanhoutte PM. Effects of hydrogen peroxide on the responsiveness of isolated canine bronchi: role of prostaglandin E<sub>2</sub> and I<sub>2</sub>. *Am J Physiol* 1992; 263: L402-L408.
- Nasstrom J, Boo E, Stahlberg M, Berge OG. Tissue distribution of two NMDA receptor antagonists, [<sup>3</sup>H]CGS 19755 and [<sup>3</sup>H]MK-801, after intrathecal injection in mice. *Pharmacol Biochem Behav* 1993; 44: 9-15.
- Gill SS, Mueller RW, McGuire PF, Pulido OM. Potential target sites in peripheral tissues for excitatory neurotransmission and excitotoxicity. *Toxicol Pathol* 2000; 28: 277-284.
- Cincotta M, Beart PM, Summers RJ, Lodge D. Bidirectional transport of NMDA receptor and ionophore in the vagus nerve. *Eur J Pharmacol* 1989; 160: 167-171.
- Lewis SJ, Cincotta M, Verberne AJM, Jarrott B, Lodge D, Beart PM. Receptor autoradiography with [<sup>3</sup>H]-L-glutamate reveals the presence and axonal transport of glutamate receptors in vagal afferent neurones of the rat. *Eur J Pharmacol* 1987; 144: 413-415.
- Joos GF, Pauwels RA, van der Straeten ME. The mechanism of tachykinin-induced bronchoconstriction in the rat. *Am Rev Respir Dis* 1988; 137: 1038-1044.
- Stretton D, Belvisi MG, Barnes PJ. The effect of sensory nerve depletion on cholinergic neurotransmission in guinea pig airways. *J Pharmacol Exp Ther* 1992; 260: 1073-1080.
- John C, Brunner S, Tanaka DT. Neuromodulation mediated by neurokinin-1 subtype receptors in adult rabbit airways. *Am J Physiol* 1993; 265: L228-L233.
- Nicholson A, Phillips CL, Allen DH, Ward HE, Berend N. The effect of L-glutamic acid on airway function and reactivity in the rabbit. *Agents Actions* 1988; 25: 267-272.
- Spina D, Matera GM, Riccio MM, Page CP. A comparison of sensory nerve function in human, guinea-pig, rabbit and marmoset airways. *Life Sci* 1998; 63: 1629-1642.
- Yuan L, Burcher E, Nail BS. Characterization of tachykinin receptors mediating bronchomotor and vasodepressor responses to neuropeptide  $\gamma$  and substance P in the anaesthetized rabbit. *Pulmon Pharmacol Ther* 1998; 11: 31-39.
- Tousignant C, Chan CC, Young D, Guevremont D, Rodger IW. Neurokinin receptor mediated plasma extravasation in guinea pig and rat airways: comparison of <sup>125</sup>I-labelled human fibrinogen and 99mTc-labelled human serum albumin as markers of leakage. *Can J Physiol Pharmacol* 1993; 71: 506-511.
- Evangelista S, Paoli S, Giachetti A, Manzini S. Involvement of tachykinin NK1 receptors in plasma protein extravasation induced by tachykinins in the guinea pig upper airways. *Neuropeptides* 1997; 31: 65-70.
- Auberson S, Lundberg JM. Lactic acid-induced plasma protein extravasation in rat airways by stimulation of sensory nerves and NK1 receptor activation. *Pharmacol Toxicol* 1993; 73: 305-310.
- Matheson MJ, Rynell AC, McClean MA, Berend N. Tachykinins do not cause plasma leakage in the rabbit trachea. *Respir Physiol* 1997; 108: 165-170.
- Black J, Diment L, Armour C, Alouan L, Johnson P. Distribution of substance P receptors in rabbit airways, functional and autoradiographic studies. *J Pharmacol Exp Ther* 1990; 253: 381-386.
- Cook JA, Brunner SL, Tanaka DT. Neurokinin receptors mediating substance P-induced contraction in adult rabbit airways. *Am J Physiol* 1990; 258: L99-L106.
- Robertson BS, Satterfield BE, Said SI, Dey RD. N-Methyl-D-aspartate receptors are expressed by intrinsic neurons of rat larynx and esophagus. *Neurosci Lett* 1998; 244: 77-80.
- Canning BJ, Undem BJ, Karakousis PC, Dey RD. Effects of organotypic culture on parasympathetic innervation of guinea pig trachealis. *Am J Physiol* 1996; 271: L698-L706.
- Colasurdo GN, Loader JE, Graves JP, Larsen GL. Maturation of non-adrenergic non-cholinergic inhibitory system in normal and allergen-sensitized rabbits. *Am J Physiol* 1994; 267: L739-L744.
- Fame TM, Colasurdo GN, Loader JE, Graves JP, Larsen GL. Decrease in the airways nonadrenergic noncholinergic inhibitory system in allergen sensitized rabbits. *Pediatr Pulmonol* 1994; 17: 296-303.
- Fame TM, Loader JE, Graves JP, Colasurdo GN, Larsen GL. Decrease in the airways non-adrenergic non-cholinergic inhibitory system in allergen sensitized rabbits. *Am Rev Respir Dis* 1993; 147: A285.
- Uddman R, Alumets J, Densert O, Hakanson R, Sundler F. Occurrence and distribution of VIP nerves in the nasal mucosa and tracheobronchial wall. *Acta Otolaryngol* 1978; 86: 443-448.
- Sakai N, Tamaoki J, Kobayashi K, Kanemura T, Isono K, Takeyama K, Takeuchi S, Takizawa T. Vasoactive intestinal peptide stimulates ciliary motility in rabbit tracheal epithelium: modulation by neutral endopeptidase. *Regulat Pept* 1991; 34: 33-41.
- Said SI. Vasoactive intestinal peptide. In: Raeburn D, Giembycz MA, eds. *Airways Smooth Muscle: Peptide Receptors, Ion Channels and Signal Transduction*. Basel: Birkhäuser Verlag, 1995: 87-113.
- Regal JF, Johnson DE. Indomethacin alters the effects of substance-P and VIP on isolated airway smooth muscle. *Peptides* 1983; 4: 581-584.
- Farmer SG, Togo J. Effects of epithelium removal on relaxation of airway smooth muscle induced by vasoactive intestinal peptide and electrical field stimulation. *Br J Pharmacol* 1990; 100: 73-78.
- Uzuner K, Tunçel N, Aydin Y. Relaxing effects of vasoactive intestinal peptide (VIP) on the contractile actions of endothelin-3, histamine, and acetylcholine in isolated guinea pig tracheal smooth muscle with or without epithelium. *Peptides* 1996; 17: 299-303.

39. Colasurdo GN, Loader JE, Graves JP, Larsen GL. Modulation of acetylcholine release in rabbit airways in vitro. *Am J Physiol* 1995; 268: L432-L437.
40. Yoneda Y, Ogita K. Localization of [<sup>3</sup>H]glutamate binding sites in rat adrenal medulla. *Brain Res* 1986; 383: 387-391.
41. Yoneda Y, Ogita K. Enhancement of [<sup>3</sup>H]glutamate binding by N-methyl-D-aspartic acid in rat adrenal. *Brain Res* 1987; 406: 24-31.
42. Winter CR, Baker RC. L-Glutamate-induced changes in intracellular calcium oscillation frequency through non-classical glutamate receptor binding in cultured rat myocardial cells. *Life Sci* 1995; 57: 1925-1934.
43. Barnard EA. Ionotropic glutamate receptors: new types and new concepts. *Trends Pharmacol Sci* 1997; 18: 141-147.
44. Marin P, Quignard JF, Lafon-Cazal M, Bockaert J. Non-classical glutamate receptors, blocked by both NMDA and non-NMDA antagonists, stimulate nitric oxide production in neurons. *Neuropharmacology* 1993; 32: 29-36.

**Received 24 July 2001**

**Accepted 14 November 2001**



The Scientific  
World Journal

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Gastroenterology  
Research and Practice

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



MEDIATORS  
of  
INFLAMMATION

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Journal of  
Diabetes Research

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Disease Markers

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Journal of  
Immunology Research

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



PPAR Research

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>



International Journal of  
Endocrinology

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



BioMed  
Research International

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Journal of  
Ophthalmology

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Stem Cells  
International

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



eCAM  
Evidence-Based  
Complementary and  
Alternative Medicine

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Journal of  
Obesity

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Journal of  
Oncology

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Computational and  
Mathematical Methods  
in Medicine

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Behavioural  
Neurology

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Parkinson's  
Disease

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



AIDS  
Research and Treatment

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014



Oxidative Medicine and  
Cellular Longevity

Hindawi Publishing Corporation  
<http://www.hindawi.com>  
Volume 2014