

**Supplemental Table 1. The constituent ratio of different cirrhotic patients from
2002 to 2013**

	Hepatitis B cirrhosis	Hepatitis C cirrhosis	Alcoholic cirrhosis	Autoimmune cirrhosis
2002	81.53	6.39	3.34	2.77
2003	81.85	5.47	3.56	3.29
2004	81.73	6.15	4.45	2.83
2005	81.55	7.35	4.43	2.91
2006	79.90	8.71	4.13	3.08
2007	76.00	11.00	5.74	3.39
2008	73.62	11.77	5.77	4.50
2009	72.61	11.76	5.82	5.06
2010	70.01	13.14	7.07	4.19
2011	66.84	13.91	7.74	4.87
2012	65.00	13.90	8.00	6.50
2013	66.00	12.60	8.40	6.80

**Supplemental table 2. The constituent ratio of native place for different cirrhotic
patients**

	Central China	North China	Northeast China	Northwest China
Hepatitis B cirrhosis	8482 (14.44%)	22469 (38.25%)	9998 (17.02%)	2802 (4.77%)
Hepatitis C cirrhosis	1608 (16.70%)	3129 (32.50%)	2786 (28.94%)	408 (4.24%)
Alcoholic cirrhosis	535 (9.69%)	2324 (42.13%)	849 (15.39%)	156 (2.83%)
Autoimmune liver cirrhosis	515 (12.63%)	1626 (39.86%)	559 (13.70%)	167 (4.09%)