Erratum

Erratum to “Anti-Interleukin-22-Neutralizing Antibody Attenuates Angiotensin II-Induced Cardiac Hypertrophy in Mice”

Jing Ye,^{1,2} Ling Liu,¹ Qingwei Ji,^{1,3} Ying Huang,¹ Ying Shi,¹ Lei Shi,¹ Jianfang Liu,² Menglong Wang,² Yao Xu,² Huimin Jiang,² Zhen Wang,² Yingzhong Lin ^{1,2} and Jun Wan ²

¹Department of Cardiology, The People's Hospital of Guangxi Zhuang Autonomous Region, Nanning, China

²Department of Cardiology, Hubei Key Laboratory of Cardiology, Renmin Hospital of Wuhan University, Cardiovascular Research Institute, Wuhan University, Wuhan 430060, China

³Emergency & Critical Care Center, Beijing Institute of Heart, Lung, and Blood Vessel Diseases and Beijing Anzhen Hospital, Capital Medical University, Beijing 100029, China

Correspondence should be addressed to Yingzhong Lin; yingzhonglin@126.com and Jun Wan; wanjun1963@126.com

Received 11 June 2018; Accepted 12 July 2018; Published 26 August 2018

Copyright © 2018 Jing Ye et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article “Anti-Interleukin-22-Neutralizing Antibody Attenuates Angiotensin II-Induced Cardiac Hypertrophy in Mice” [1], there was an error in Figure 1(a), where the IL-22, β -MHC, and GAPDH bands were triplicated, due to a production error. The corrected figure is shown below.

FIGURE 1: Effect of chronic angiotensin II infusion on IL-22/IL-22R1 protein levels. (a) Representative and (b) quantitative expression of IL-22, IL-22R1, ANP, BNP, and β -MHC in LV tissue of baseline and angiotensin II-infused mice. $n = 4$ for each group. * $p < 0.05$ versus baseline level.

References

[1] J. Ye, L. Liu, Q. Ji et al., "Anti-Interleukin-22-neutralizing antibody attenuates angiotensin II-induced cardiac hypertrophy in mice," *Mediators of Inflammation*, vol. 2017, Article ID 5635929, 10 pages, 2017.