

## Review Article

# Immunological Perspective: *Helicobacter pylori* Infection and Gastritis

Hang Yang and Bing Hu 

Department of Gastroenterology, West China Hospital, Sichuan University, Chengdu, Sichuan, China

Correspondence should be addressed to Bing Hu; hubingnj@163.com

Received 5 February 2022; Accepted 2 March 2022; Published 8 March 2022

Academic Editor: Shin-ichi Yokota

Copyright © 2022 Hang Yang and Bing Hu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Helicobacter pylori* is a spiral-shaped gram-negative bacterium. Its infection is mainly transmitted via oral-oral and fecal-oral routes usually during early childhood. It can achieve persistent colonization by manipulating the host immune responses, which also causes mucosal damage and inflammation. *H. pylori* gastritis is an infectious disease and results in chronic gastritis of different severity in near all patients with infection. It may develop from acute/chronic inflammation, chronic atrophic gastritis, intestinal metaplasia, dysplasia, and intraepithelial neoplasia, eventually to gastric cancer. This review attempts to cover recent studies which provide important insights into how *H. pylori* causes chronic inflammation and what the characteristic is, which will immunologically explain *H. pylori* gastritis.

## 1. Introduction

*Helicobacter pylori* (*H. pylori*) has coevolved with its human host for no less than 30,000 years. It can safely colonize around the epithelium of gastric gland via their specific microstructure and self-synthesizing proteins such as chemoreceptors [1], flagella [2], and urease [3]. Chemotaxis assists *H. pylori* to find nutrients (urea and arginine) and avoid toxic substances like reactive oxygen species (ROS). *H. pylori* obtains nutrients from blood [4] and even extract nutrients from the host cells such as lipid, cobalt, iron, and nickel [5, 6]. Arginine is required for *H. pylori* growth and is sensed by TlpA [7]. Urea and gastric mucus pH gradient are sensed by TlpB for self-protection and chemotactic orientation [8, 9]. Additionally, *H. pylori* uses TlpB to sense injured sites and preferentially colonizes injured sites in the mouse stomach, independent on urea [10], which may indicate the inflammatory condition benefits the growth of *H.*

*pylori*. Spiral shape is necessary to fast move and penetrate in the mucus layer via corkscrew-like movement [11]. Two flagellins (FlaA and FlaB) are indispensable for bacteria motility [12, 13]. FlaA mutants show a greater decline of motility than that of FlaB mutants [14]. *H. pylori* lacking MotB is nonmotile and retains only a nonfunctioning flagellar structure [13]. Urease is a nickel-dependent metalloenzyme [15]. Both insufficient cytoplasmic nickel availability or excessive nickel entry impair the activation of urease and the survival of *H. pylori* [16, 17]. Urea hydrolyzed by urease finally produces ammonia and CO<sub>2</sub> to participate in lowering pH and the regulation of mucus viscoelasticity [18]. *H. pylori* enters the mucus layer after a short duration within the lumen and persistently distributes within approximately 25–30 μm of mucosal epithelial cells where the pH ranges from 4.5 to 6.5, or directly on the epithelium or deep inside the glands [3, 19, 20]. After BabA/SabA-mediated adhesion [21], *H. pylori*, gastric epithelial cells, and leukocytes interact with

each other to achieve the balance. During *H. pylori* infection, immune cells infiltrate to the lamina propria and submucosa, aiming to clear *H. pylori* [22]. *H. pylori* skews host immune response to avoid clearance and achieve persistence, such as cholesterol glycosylation, escaping the Toll-like receptor (TLR) recognition, tolerating dendritic cells (DCs), blocking T cell proliferation, inducing Treg skewing, and upregulating PD-L1 [23–26].

Inflammation is triggered when innate immune cells detect infection or tissue injury [27]. Although *H. pylori* manipulates the host immune system, inadequate immune response and inflammation are still initiated, leading to chronic active gastritis. Inflammation is an essential and a complex biological process that protects the body from potential harm caused by infection or injury [28] and develops in response to pathogen-associated molecular patterns (PAMPs) from *H. pylori* and damage-associated molecular patterns (DAMPs) from damaged epithelial cells [29]. Pattern recognition receptors (PRRs) are membrane associated or soluble, which are owned by both immune and nonimmune cells, and respond to PAMPs and DAMPs, initiating downstream signaling cascades, including the production and secretion of pro- and anti-inflammatory cytokines to further modulate immune response [30].

## 2. PAMPs Derived from *H. pylori* Inducing Inflammation: NF- $\kappa$ B and Type I IFNs

**2.1. Antigen-Presenting Cells (APCs).** PRRs expressed on APCs (macrophages, DCs, and B cells) including TLRs, nucleotide-binding oligomerization- (NOD-) like receptors (NLRs), and C-type lectin receptors (CLRs) can detect PAMPs derived from *H. pylori* [31] such as lipopolysaccharide (LPS), lipoproteins and peptidoglycan (TLR2), dsRNA and polycytidylic acid (TLR3), LPS and heat shock proteins (TLR4), flagellin (TLR5 and 11), and unmethylated CpG containing ssDNA (TLR9) [32–34]. TLR activation increases the activity of NF- $\kappa$ B and transcription of type I IFNs [35]. Both TLR2 and TLR4 on DCs and macrophages can recognize LPS and shape the *H. pylori*-induced pro- and anti-inflammatory cytokines and chemokine milieu [36, 37]. Receptor complex combining TLR2 with TLR1, 6 or 10, not TLR4 is also reported [38–40], which further indicates both pro- and anti-inflammation roles of TLR2. TLR9 is expressed exclusively in intracellular vesicles and recognizes unmethylated CpG DNA motifs to induce the transcription of NF- $\kappa$ B and IRF7 via MyD88 and subsequent increase of inflammatory cytokines and type I IFNs in APCs [41, 42]. It has also demonstrated that TLR9 is involved in both suppressing and promoting inflammation after recognizing *H. pylori* DNA [41]. For example, increasing Th1 cells (IFN $\gamma$ ) and Th17 cells (Th17) is found in *Tlr9*<sup>-/-</sup> mice [43, 44], and the roles of activating NF- $\kappa$ B, upregulating the expression of COX-2/prostaglandin E2, and activating neutrophils are also found after activating TLR9 [45, 46]. CLRs expressed by DCs are pivotal for both antigen presentation and Th cell differentiation [47]. CLRs on DCs and macrophages can be activated by *H. pylori* metabolites modified from host cholesterol (cholesteryl acyl  $\alpha$ -glucoside and cho-

lesteryl phosphatidyl  $\alpha$ -glucoside), exacerbating gastritis [48, 49]. C-type lectin DC-specific intercellular adhesion molecule-grabbing nonintegrin (DC-SIGN) can recognize LPS Le antigens [50], which can be misused by distinct mechanisms that either circumvent antigen processing or alter TLR-mediated signaling to decrease Th1 cells and increase Treg cells [51, 52]. For example, *H. pylori* modulates the Th1/Th2 balance through the phase-variable interaction between LPS and DC-SIGN and the variation of O-antigen decorated by fucose residues binding to DC-SIGN to block Th1 development [53, 54]. B cells also express TLR and MyD88. CpG (TLR9), LPS (TLR4), and peptidoglycan (TLR2) are found to induce B cell-derived IL-6, IL-12, and IL-10 [55, 56]. TLR signaling in B cells inhibits inflammatory T cell response (both Th1 and Th17 cells), which can be controlled by TLR agonists [57]. Further B cells produce IL-10 after being activated by *H. pylori* and suppress the differentiation of DCs activated by *H. pylori* [58]. And activated TLR2 by *H. pylori* on B cells also induces Treg (IL-10) [59].

**2.2. Gastric Epithelial Cells.** TLR2, TLR4, TLR5, TLR9, and NOD1 are expressed by epithelial cells and induce NF- $\kappa$ B activation and IFN production in these cells to defense against *H. pylori*, as a central principle of mucosal immunity [60–63]. Increasing expression of TLR2, TLR4, TLR5, and TLR9 in gastric epithelia of children's gastritis [64]; TLR4, TLR5, and TLR9 in adults' gastritis [60]; TLR2, TLR4, and TLR5 in gastric dysplasia [65]; and TLR4, TLR5, and TLR9 in gastric cancer (GC) [61] is found. Expression of TLR2 and TLR4 in chronic gastritis caused by *H. pylori* remains increased after eradication therapy in 3 months [66]. TLR2 is the most extensively expressed receptor among all the TLRs in gastric mucosa infected by *H. pylori* [67, 68]. In gastric epithelial cells, TLR2 cooperates with TLR4 to strengthen the innate immune response to LPS and activate NF- $\kappa$ B and inducible nitric oxide synthase (iNOS) [68]. Significantly increasing TLR9 in gastric epithelial cells are demonstrated in patients residing in the region with a high GC risk, and *H. pylori* isolated from them can cause increasing activation of TLR9 [69]. Regarding on CLRs, besides of the expression on the surface of DCs, DC-SIGN is overexpressed in gastric epithelial cells, when facing LPS stimulation [70, 71] and induces a Th1 dominating cytokine response [71]. Furthermore, DC-SIGN stimulated by LPS interacts with TLR4, promotes NLRP3, and regulates the production of IL-1 $\beta$  and IL-18 in gastric epithelial cells [70]. *H. pylori* peptidoglycan delivered into host cells by the T4SS is recognized by epithelial cells via Nod1 [72], leading to NF- $\kappa$ B activation and the production of  $\beta$ -defensin and type I IFNs from Nod1-activated gastric epithelial cells [73–75]. *H. pylori* can secrete outer membrane vesicles (OMVs) separated from the bacterial outer membrane [76]. OMVs containing peptidoglycan enter epithelial cells at cholesterol-rich lipid rafts and induce NOD1-dependent response [77]. Apart from PRRs, D-glycero- $\beta$ -D-manno-heptose 1,7-bisphosphate ( $\beta$ -HBP) is a T4SS-dependent effector of NF- $\kappa$ B activation via alpha-protein kinase 1- (ALPK1-) TRAF-interacting protein with forkhead-associated domain (TIFA) in gastric epithelial cells [78]. ADP-glycero- $\beta$ -D-manno-

heptose ( $\beta$ -ADP heptose), a derivative of  $\beta$ -HBP, is more active than  $\beta$ -HBP [79].  $\beta$ -ADP heptose mediates NF- $\kappa$ B activation and cytokine expression after directly binding the N-terminal domain of ALPK1 [80]. Additionally, CagA also contributes to the NF- $\kappa$ B signaling after TIFA and NOD1 activation [81].

### 3. DAMPs from Gastric Epithelium Inducing Inflammation

Acute and chronic inflammation caused by *H. pylori* can damage epithelial cells and induce inflammatory edema, atrophy, and necrosis/apoptosis. DAMPs are released during tissue damage and typically derived from intracellular and extracellular continents, which are recognized by PRRs (TLRs and NLRs) and by non-PRRs (RAGE, CD44, integrins, and CD91) to recruit neutrophils and monocytes and activate inflammation and tissue repair [82]. Interleukin-1 (IL-1) and tumor necrosis factor (TNF) are the notable proinflammatory cytokines in this process.

IL-1 (IL-1 $\alpha$  and IL-1 $\beta$ ) is the proinflammatory cytokine mainly produced by macrophages and acts through IL-1 receptor (IL-1R), which is important to recruit neutrophils and monocytes and to induce additional proinflammatory cytokines [83]. Caspase-1 is responsible to cleave IL-1 $\beta$  into the biologically active form [84]. Caspase-1 is activated by inflammasomes composed of a PRR of the NLR family such as NLRP3, NLRC4, and AIM2 inflammasome [85]. Research has identified that *H. pylori* can activate inflammasomes. For example, caspase-1 activation and production of IL-1 $\beta$  and IL-18 in mice are the consequence of *H. pylori* infection, in which IL-1 $\beta$  is produced for clearance and IL-18 is for persistence [86]. AIM2 inflammasome is demonstrated to recognize cytoplasmic DNA [87], which is also induced by the OMVs of gram-negative pathogens [88]. NLRC4 expression is upregulated by proinflammatory stimuli like TNF $\alpha$  [89]. *H. pylori* exploits the NLRC4 inflammasome to enhance neutrophil infiltration and induce IL-18 production in gastric epithelial cells to block  $\beta$ -defensin expression via NF- $\kappa$ B activation [90]. NLRP3 can recognize nucleic acids, bacterial proteins, and metabolites [91]. The proinflammatory cytokines released upon NLRP3 activation are IL-1 $\beta$ , IL-18, HMGB1, leukotrienes, and prostaglandins [92]. *H. pylori* infection activates the NLRP3 inflammasome and IL-1 $\beta$  production in neutrophils [93, 94], differentiated macrophages [95], and DCs [96]. Furthermore, the activation of NLRP3 inflammasome by *H. pylori* via ROS signaling pathway also leads to the production of IL-1 $\beta$  and IL-18 in human monocytes [97]. These results suggest a dual role of the inflammasome in *H. pylori* infection. Additionally, Treg cells can be activated by the activation of axis of urease enzyme of *H. pylori*/TLR2/NLRP3/caspase-1/IL-18 [98]. Furthermore, the application of exogenous activators induces NLRP3 inflammasome formation, and the secretion of high amounts of IL-1 $\beta$  in infected cells, which indicates cellular injury regardless of causes (e.g., bile, smoking, alcohol, drugs, and other gastric microbiota), may have synergistic effect with *H. pylori* infection to exacerbate damage to gastric epithelium [99].

TNF has been participated in regulating immunity and inflammation. It is widely and constitutively expressed by activated immune cells, as well as by fibroblasts and endothelial and epithelial cells responding to proinflammatory cytokines including TNF itself [100, 101]. It is cleaved by TNF $\alpha$ -converting enzyme to release soluble TNF $\alpha$  [102]. TNF mediates inflammatory pathology through binding to TNFR1 and TNFR2 [103]. TNFR1 interacts strongly with both membrane and soluble TNF $\alpha$ , whereas TNFR2 binds to membrane TNF $\alpha$  with much higher affinity. TNFR1 is expressed by near all cells. TNFR2 expression is limited to cells of immune and endothelial origin [101]. It means *H. pylori* infection can induce TNF production and activate its signal pathways in many aspects. TNF downstream pathways mainly involve NF- $\kappa$ B, MAPKs, caspases, and ROS/RNS. TNFR1 and 2 induce the activation of MAPK and NF- $\kappa$ B. TNFR1 can also stimulate apoptosis and necroptosis, as it harbors a death domain in the cytoplasmic part [101, 104]. As a proinflammatory cytokine, TNF $\alpha$  causes vasodilatation and edema, leukocyte adhesion to epithelium, and oxidative stress in inflammatory sites, mediated by the induction of NO, prostanoids, and ROS [105, 106]. TNF $\alpha$  may also regulating the production of ROS and RNS [107]. Macrophages, DCs, and gastric epithelial cells produce TNF and IL-1 $\beta$  in a dose- and time-dependent manner after exposure to *H. pylori* [108–110]. TNF $\alpha$  induces the apoptosis of parietal cells in *H. pylori* infected rat [111]. Increasing TNF $\alpha$  is found in *H. pylori* infected chronic atrophic gastritis (CAG) and associated with chronic inflammation degree [112]. Soluble TNF receptors (sTNFRs) are shown actively produced in *H. pylori* infected gastric mucosa, and anti-sTNFR monoclonal antibodies increase TNF-induced gastric epithelial cell apoptosis, which suggests that sTNFR has a protective effect [113]. In addition, TNFR1 increase is seemingly related to the aggressiveness of gastric lesions [114]. These may indicate severe infection cause increasing TNF $\alpha$  accompanied with increasing sTNFRs to avoid more loss of gastric epithelial cells.

NF- $\kappa$ B activation may prevent cell death because it controls the transcription of a number of genes involved in cell survival, proliferation, and inflammation [107]. Oxidative stress has a crosstalk with both TNF [107] and NF- $\kappa$ B [115] via bidirectional effects, as they have complex interaction with each other during production process. Briefly, it means oxidative stress can both activate and inhibit NF- $\kappa$ B pathway, and NF- $\kappa$ B pathway has anti- and prooxidant role in oxidative stress [116]. TNF promotes the production of ROS and NF- $\kappa$ B. TNF-induced ROS inhibits NF- $\kappa$ B activation, reduces NF- $\kappa$ B-mediated survival signaling, and explains the cell death associated with high ROS levels [116]. Mitochondrial ROS can facilitate TNF-mediated NF- $\kappa$ B activation [115]. Furthermore, these effects are further complicated by TNF-induced NO production. TNF-induced NF- $\kappa$ B promotes the transcription of the gene that encodes iNOS for producing NO with anti- and prooxidant roles [117]. There may be also a subtle NO/ROS/RNS balance in TNF signaling [118]. *H. pylori* increases ROS and RNS mainly from immune cells such as neutrophils and gastric epithelial cells, correlating with the severity of mucosal

inflammatory damage and genetic instability [119, 120]. In addition, iNOS expression is highly induced in the epithelium of atrophic gastritis, as well as metaplasia and dysplasia [121]. NF- $\kappa$ B can stimulate iNOS in gastric epithelial cells maybe further to prevent cell death [68], which depends on DNA damage caused by ROS/RNS in gastric epithelial cells [122, 123]. Furthermore, the long-lasting state with inflammation, oxidative stress, and DNA damage may also lead in GC [124].

#### 4. Neutrophil Recruitment and Inflammation

Neutrophils are the first leukocytes recruited to the inflammatory site during acute inflammation, aiming at eliminating pathogens by phagocytosis, deregulation, and neutrophil extracellular traps. Neutrophils and macrophages produce ROS, proteases, and growth factors, leading to tissue destruction, fibroblast proliferation, abnormal accumulation of collagen, and fibrosis [125]. For example, ROS produced by neutrophils and NO produced by macrophages fail killing *H. pylori*, but do damage on gastric epithelial cells (e.g., nuclear DNA and mitochondria, even cell death or gastric carcinogenesis) [119, 124, 126]. Neutrophil recruitment is started by changes on endothelial surface induced by inflammatory cytokines released from tissue-resident sentinel leukocytes exposed to pathogens [127]. Macrophages and mast cells reside in tissues are sentinel cells to initiate neutrophil recruitment via increasing the permeability of local blood vessels and chemokine secretion after the activation of PRRs such as TLRs, NLRs, and CLRs [128, 129]. For instance, TNF $\alpha$  produced from activated macrophages and mast cells acts as a crucial role to recruit neutrophils [130]. *H. pylori* infection is characterized by rapid and continuous recruitment of neutrophils followed by T and B cells, plasma cells, and macrophages [131]. Apart from the activation of PRRs causing neutrophil recruitment (e.g., both TNF $\alpha$  and IL-1 $\beta$  can induce neutrophil recruitment), *H. pylori* neutrophil-activating protein (HP-NAP) can induce trans-endothelial migration of neutrophils and activate neutrophils such as the release of myeloperoxidase and production of ROS/RNS, IL-8, and CCL4 [132–135], which is attenuated when lacking SabA [136] or ablating hepatoma-derived growth factor [137]. HP-NAP is shown to determine the host risk of dyspepsia by ROS exposure and chronic inflammation [138]. Additionally, HP-NAP has immune modulating roles and induces cytokines from other immune cells [139]. After entering into the inflammatory tissue site, neutrophils express many cell surface receptors [140] and recognize PAMPs and DAMPs by neutrophil PRRs [141], opsonins by opsonic receptors and bacterial products [142], and endogenous molecules released during inflammation by G protein coupled receptors [143]. Neutrophils are involved in the complex bilateral interactions with aforementioned immune cells [144, 145]. HP-NAP can regulate immunity, which may be attributed to the regulation from neutrophil itself, which means immunity is actually regulated by neutrophils, and HP-NAP is just responsible to recruit neutrophils.

#### 5. Adaptive Immune and Inflammation

Adaptive immunity is responsible for the production of antibodies and the activation of cytotoxic lymphocytes after recognizing antigen peptides presented by APCs [146]. Naïve CD4+ T cells are induced to differentiate towards Th1 (IL-2 and IFN $\gamma$ ), Th2 (IL-4, IL-5, IL-3, and IL-13), Th17 (IL-17 and IL-22), and Treg (IL-10 and TGF- $\beta$ ) phenotypes according to the local cytokine milieu [147, 148]. *H. pylori* infection can induce these cells differentiation with different proportions and a balanced cytokine network. As a pathogen, innate and subsequent adaptive immune responses are evoked to eliminate *H. pylori*. Th1 and Th17 and their corresponding cytokines are required for infection control [149, 150]. As a persistent colonizer coevolving with human, it can skew adaptive immune response. For example, both VacA and  $\gamma$ -glutamyltransferase (GGT) possess pro- and anti-inflammatory effects. They block the proliferation of T cells via blocking the cell cycle [151, 152] and induce cell death and proinflammatory cytokine production (TNF- $\alpha$  and IL-1 $\beta$  induced by VacA, and cyclooxygenase-2, prostaglandin E2, NF- $\kappa$ B, and IL-8 induced by GGT) [153–157]. *H. pylori* can induce the Treg cell differentiation and increase IL-10 and TGF- $\beta$ 1 in infected patients, in particular in children [158–160]. Treg cell differentiation requires the direct interaction between naïve T cells and tolerogenic DCs exposed to *H. pylori* [161, 162]. GGT and VacA contribute to *H. pylori*'s tolerance-promoting effects on DCs [163]. The cytokine network is concluded in detail.

IL-12 cytokine family contains IL-12, IL-23, IL-27, and IL-35, which serve as a vital bridge between innate and adaptive response [164]. IL-12 expression regulates innate response and controls the differentiation of Th cell type. IL-12 is produced by DCs, monocytes/macrophages, and neutrophils and triggers the differentiation of Th1 cells (IL-2 and IFN $\gamma$ ) [165]. IL-12 can also activate NK cells, other T cells, and DCs/macrophages to produce IFN $\gamma$  [166, 167]. IFN $\gamma$  stimulates the bactericidal activity of phagocytic cells to boost innate immune response, which is modulated by IL-4, IL-10, and IL-18 [168]. IL-23 is produced by myeloid cells, especially DCs and monocyte/macrophage lineage cells [169]. IL-23 induces IL-1 $\beta$  and TNF $\alpha$  produced by myeloid cells and NK and T cells and drives IL-12 and IFN $\gamma$  production [168]. IL-23 combined with TGF- $\beta$  and IL-6 determines the differentiation of Th17, in which TGF- $\beta$  plus IL-6 (mainly produced by monocytes/macrophages) act as the differentiation factor and IL-23 acts as the growth and stabilization factor [170]. Rapid IL-23/IL-17 immune response can promote chronic inflammation with cytokines such as IL-17, IL-6, IL-8, and TNF [171]. For example, activated sentinel DCs and macrophages produce IL-23 to trigger the release of IL-17 from tissue-resident T cells and NK cells. IL-17 increases the secretion of IL-1, IL-6, IL-8, CXCL1, and TNF in stromal, epithelial, and endothelial cells and monocyte subpopulations and further recruits neutrophils [172]. IL-27 produced by the myeloid lineage (mainly monocytes and DCs) [173] limits the production of IL-2 and GM-CSF; reverses the IL-23 mediated lineage commitment of Th17 cells; induces IL-10 produced from Th1,

Th2, Th17, and Treg cells; and promotes Treg cells specialized to limit Th1 cells [174]. Further, both IFN $\gamma$  and IL-27 promote a population of Treg cells, which restrict Th1 cell-mediated pathology [175].

IL-35 is produced by Treg cells, CD8<sup>+</sup> Treg cells, DCs, and B cells. It inhibits Th17 cell differentiation and promotes both Treg cell proliferation and corresponding functions [176, 177]. IL-10 is produced by APCs, mast cells, eosinophils, neutrophils, NK cells, and T cells [178]. IL-10 has profound anti-inflammatory functions. IL-10 inhibits the release of proinflammatory cytokines (e.g., TNF $\alpha$ , IL-1 $\beta$ , IL-6, IL-8, and GM-CSF) and chemokines (e.g., MCP1, IL-8, and IP-10) from DCs and monocytes/macrophages. IL-10 suppresses both IL-12 and IL-23 to limit CD4<sup>+</sup> T cell differentiation and proliferation. IL-10 attenuates neutrophil recruitment by decreasing inflammatory cytokines [179]. TGF- $\beta$ 1 is widely expressed in leukocytes and stromal cells and responsible for wound healing, immune tolerance, and the modulation of cell growth and differentiation [180]. TGF- $\beta$ 1 combined with IL-2 is critical for the differentiation of Treg cells from naïve T cells [181]. TGF- $\beta$ 1 produced by Treg cells is required to inhibit Th1 cell differentiation and promote immune tolerance [182, 183].

Type I IFNs (IFN $\alpha$  and IFN $\beta$ ) and type II IFN (IFN $\gamma$ ) orchestrate innate and adaptive immunity via multiple mechanisms. Type I IFNs are produced by almost every cell type with DCs, macrophages, and epithelial cells mainly through TLR-dependent pathways [184, 185]. Type I IFNs activate NK cells, macrophages, and DCs to boost innate immunity [186] and enhance adaptive immunity through promoting the differentiation of T and B cells and their activation. Type I IFNs can also inhibit Th17 cells and induce IL-10 and PD-1 ligand expression by immune cells like DCs and macrophages in chronic infection [187, 188]. IFN $\gamma$  is the sole type II IFN, exclusively produced by Th1 cells, CD8<sup>+</sup> cytotoxic T lymphocytes (CTLs), NK cells, innate lymphoid cells (ILCs), and DCs [189]. An early production of IFN $\gamma$  is from ILCs, and abundant and sustained IFN $\gamma$  are produced by Th1 cells or CTLs after recognizing microbial peptides from APCs [190]. IFN $\gamma$  is positively regulated by IL-12 and IL-18 and negatively regulated by IL-4, IL-10, TGF- $\beta$ , and glucocorticoids [189]. The effects of IFN $\gamma$  on adaptive immunity have been reviewed on facilitating Th1 cell differentiation, inhibiting Th2 and Th17 cells, activating Treg cells, and promoting B cell class switching [191]. IFN $\gamma$  can also strongly promote innate immune. IFN $\gamma$ -polarized macrophages are highly responsive to a variety of inflammatory stimuli such as TNF, type I IFNs, microbial products, and ligands for TLRs [192]. IFN $\gamma$  orchestrates the differentiation of monocytes into DCs and macrophages, which is the primary sources of IL-12 at the infection site [193].

## 6. Conclusions

*H. pylori* gastritis is ultimately attributed to the activation of PRRs on APCs, gastric epithelial cells, and neutrophils. And complex multilateral crosstalk between gastric epithelium, innate immune response, and adaptive immune response permit *H. pylori* persistent colonization. From the aforemen-

tioned immune response and cytokines, there is a dynamic balance between inflammation and immunity, which processes to a newer and severer balance gradually, as the persistence of *H. pylori* provides sustained stimuli. The balance is achieved by pro- and anti-inflammatory cytokine network from these leukocytes. It further indicates that relieving or aggravating the inflammation by increasing/decreasing one specific cytokine to break this balance may facilitate the eradication of *H. pylori*, which also explains the reason why addition of vitamin C (an antioxidant and -inflammation role to break the balance) to *H. pylori* eradication treatment may increase the eradication rate. Further research is needed.

## Conflicts of Interest

The authors declared no potential conflicts of interest.

## References

- [1] K. S. Johnson and K. M. Ottemann, "Colonization, localization, and inflammation: the roles of *H. pylori* chemotaxis *in vivo*," *Current Opinion in Microbiology*, vol. 41, pp. 51–57, 2018.
- [2] H. Gu, "Role of flagella in the pathogenesis of *Helicobacter pylori*," *Current Microbiology*, vol. 74, no. 7, pp. 863–869, 2017.
- [3] S. Ansari and Y. Yamaoka, "Survival of *Helicobacter pylori* in gastric acidic territory," *Helicobacter*, vol. 22, no. 4, article e12386, 2017.
- [4] G. L. Mendz and B. P. Burns, "Characterization of arginine transport in *Helicobacter pylori*," *Helicobacter*, vol. 8, no. 4, pp. 245–251, 2003.
- [5] S. Tan, J. M. Noto, J. Romero-Gallo, R. M. Peek Jr., and M. R. Amieva, "*Helicobacter pylori* perturbs iron trafficking in the epithelium to grow on the cell surface," *PLoS Pathogens*, vol. 7, no. 5, article e1002050, 2011.
- [6] B. Kalali, R. Mejías-Luque, A. Javaheri, and M. Gerhard, "*H. pylori* virulence factors: influence on immune system and pathology," *Mediators of Inflammation*, vol. 2014, Article ID 426309, 9 pages, 2014.
- [7] O. A. Cerda, F. Núñez-Villena, S. E. Soto, J. M. Ugalde, R. López-Solís, and H. Toledo, "tlpA gene expression is required for arginine and bicarbonate chemotaxis in *Helicobacter pylori*," *Biological Research*, vol. 44, no. 3, pp. 277–282, 2011.
- [8] E. Goers Sweeney, J. N. Henderson, J. Goers et al., "Structure and Proposed Mechanism for the pH-Sensing *Helicobacter pylori* Chemoreceptor TlpB," *Structure*, vol. 20, no. 7, pp. 1177–1188, 2012.
- [9] M. A. Croxen, G. Sisson, R. Melano, and P. S. Hoffman, "The *Helicobacter pylori* chemotaxis receptor TlpB HP0103 is required for pH taxis and for colonization of the gastric mucosa," *Journal of Bacteriology*, vol. 188, no. 7, pp. 2656–2665, 2006.
- [10] H. Hanyu, K. A. Engevik, A. L. Matthis, K. M. Ottemann, M. H. Montrose, and E. Aihara, "*Helicobacter pylori* uses the TlpB receptor to sense sites of gastric injury," *Infection and Immunity*, vol. 87, no. 89, 2019.
- [11] L. K. Sycuro, T. J. Wyckoff, J. Biboy et al., "Multiple peptidoglycan modification networks modulate *Helicobacter pylori*'s

- cell shape, motility, and colonization potential," *PLoS Pathogens*, vol. 8, no. 3, article e1002603, 2012.
- [12] C. Y. Kao, B. S. Sheu, and J. J. Wu, "\_Helicobacter pylori\_ infection: An overview of bacterial virulence factors and pathogenesis," *Biomedical Journal*, vol. 39, no. 1, pp. 14–23, 2016.
- [13] K. M. Ottemann and A. C. Lowenthal, "Helicobacter pylori uses motility for initial colonization and to attain robust infection," *Infection and Immunity*, vol. 70, no. 4, pp. 1984–1990, 2002.
- [14] C. Dunne, B. Dolan, and M. Clyne, "Factors that mediate colonization of the human stomach by Helicobacter pylori," *World Journal of Gastroenterology*, vol. 20, no. 19, pp. 5610–5624, 2014.
- [15] N. C. Ha, S. T. Oh, J. Y. Sung, K. A. Cha, M. H. Lee, and B. H. Oh, "Supramolecular assembly and acid resistance of Helicobacter pylori urease," *Nature Structural Biology*, vol. 8, no. 6, pp. 505–509, 2001.
- [16] A. H. van Vliet, F. D. Ernst, and J. G. Kusters, "NikR-mediated regulation of Helicobacter pylori acid adaptation," *Trends in Microbiology*, vol. 12, no. 11, pp. 489–494, 2004.
- [17] S. B. Mulrooney and R. P. Hausinger, "Nickel uptake and utilization by microorganisms," *FEMS Microbiology Reviews*, vol. 27, no. 2-3, pp. 239–261, 2003.
- [18] B. E. Dunn and S. H. Phadnis, "Structure, function and localization of Helicobacter pylori urease," *The Yale Journal of Biology and Medicine*, vol. 71, no. 2, pp. 63–73, 1998.
- [19] A. T. B. Abadi, "Strategies used by Helicobacter pylori to establish persistent infection," *World Journal of Gastroenterology*, vol. 23, no. 16, pp. 2870–2882, 2017.
- [20] M. R. Howitt, J. Y. Lee, P. Lertsethtakarn et al., "ChePep controls Helicobacter pylori infection of the gastric glands and chemotaxis in the Epsilonproteobacteria," *mBio*, vol. 2, no. 4, article e00098, 2011.
- [21] S. K. Lindén, C. Wickström, G. Lindell, K. Gilshenan, and I. Carlstedt, "Four modes of adhesion are used during Helicobacter pylori binding to human mucins in the oral and gastric niches," *Helicobacter*, vol. 13, no. 2, pp. 81–93, 2008.
- [22] K. Robinson, R. H. Argent, and J. C. Atherton, "The inflammatory and immune response to Helicobacter pylori infection," *Best Practice & Research Clinical Gastroenterology*, vol. 21, no. 2, pp. 237–259, 2007.
- [23] N. R. Salama, M. L. Hartung, and A. Müller, "Life in the human stomach: persistence strategies of the bacterial pathogen Helicobacter pylori," *Nature Reviews. Microbiology*, vol. 11, no. 6, pp. 385–399, 2013.
- [24] X. Zhang, I. C. Arnold, and A. Müller, "Mechanisms of persistence, innate immune activation and immunomodulation by the gastric pathogen Helicobacter pylori," *Current Opinion in Microbiology*, vol. 54, pp. 1–10, 2020.
- [25] L. Holokai, J. Chakrabarti, T. Broda et al., "Increased programmed death-ligand 1 is an early epithelial cell response to Helicobacter pylori infection," *PLoS Pathogens*, vol. 15, no. 1, article e1007468, 2019.
- [26] M. A. Qaria, S. Kumar, L. P. Sepe, and N. Ahmed, "Cholesterol glucosylation-based survival strategy in Helicobacter pylori," *Helicobacter*, vol. 26, no. 2, article e12777, 2021.
- [27] K. Newton and V. M. Dixit, "Signaling in innate immunity and inflammation," *Cold Spring Harbor Perspectives in Biology*, vol. 4, no. 3, 2012.
- [28] L. S. Taams, "Inflammation and immune resolution," *Clinical and Experimental Immunology*, vol. 193, no. 1, pp. 1–2, 2018.
- [29] E. Kolaczowska and P. Kubers, "Neutrophil recruitment and function in health and inflammation," *Nature Reviews. Immunology*, vol. 13, no. 3, pp. 159–175, 2013.
- [30] P. I. Seoane, B. Lee, C. Hoyle et al., "The NLRP3-inflammasome as a sensor of organelle dysfunction," *The Journal of Cell Biology*, vol. 219, no. 12, 2020.
- [31] R. Medzhitov and C. Janeway Jr., "Innate immunity," *The New England Journal of Medicine*, vol. 343, no. 5, pp. 338–344, 2000.
- [32] K. Takeda and S. Akira, "Toll-like receptors," *Current Protocols in Immunology*, vol. 109, no. 1, pp. 335–376, 2015.
- [33] T. Satoh and S. Akira, "Toll-like receptor signaling and its inducible proteins," *Microbiology spectrum*, vol. 4, no. 6, 2016.
- [34] K. Vijay, "Toll-like receptors in immunity and inflammatory diseases: past, present, and future," *International Immunopharmacology*, vol. 59, pp. 391–412, 2018.
- [35] K. A. Fitzgerald and J. C. Kagan, "Toll-like receptors and the control of immunity," *Cell*, vol. 180, no. 6, pp. 1044–1066, 2020.
- [36] T. Neuper, T. Frauenlob, M. Sarajlic, G. Posselt, S. Wessler, and J. Horejs-Hoeck, "TLR2, TLR4 and TLR10 shape the cytokine and chemokine release of H. pylori-infected human DCs," *International Journal of Molecular Sciences*, vol. 21, no. 11, p. 3897, 2020.
- [37] M. Obonyo, M. Sabet, S. P. Cole et al., "Deficiencies of myeloid differentiation factor 88, Toll-like receptor 2 TLR2, or TLR4 produce specific defects in macrophage cytokine secretion induced by Helicobacter pylori," *Infection and Immunity*, vol. 75, no. 5, pp. 2408–2414, 2007.
- [38] H. Nagashima, S. Iwatani, M. Cruz et al., "Toll-like receptor 10 in Helicobacter pylori infection," *The Journal of Infectious Diseases*, vol. 212, no. 10, pp. 1666–1676, 2015.
- [39] S. Yokota, T. Ohnishi, M. Muroi, K. Tanamoto, N. Fujii, and K. Amano, "Highly-purified Helicobacter pylori LPS preparations induce weak inflammatory reactions and utilize toll-like receptor 2 complex but not Toll-like receptor 4 complex," *FEMS Immunology and Medical Microbiology*, vol. 51, no. 1, pp. 140–148, 2007.
- [40] J. Li, D. S. Lee, and J. Madrenas, "Evolving bacterial envelopes and plasticity of TLR2-dependent responses: basic research and translational opportunities," *Frontiers in Immunology*, vol. 4, p. 347, 2013.
- [41] M. G. Varga and R. M. Peek, "DNA transfer and toll-like receptor modulation by Helicobacter pylori," *Current Topics in Microbiology and Immunology*, vol. 400, pp. 169–193, 2017.
- [42] R. Rad, W. Ballhorn, P. Volland et al., "Extracellular and intracellular pattern recognition receptors cooperate in the recognition of Helicobacter pylori," *Gastroenterology*, vol. 136, no. 7, pp. 2247–2257, 2009.
- [43] M. G. Varga, M. B. Piazuelo, J. Romero-Gallo et al., "TLR9 activation suppresses inflammation in response to Helicobacter pylori infection," *Gastrointestinal and Liver Physiology*, vol. 311, no. 5, pp. G852–G858, 2016.
- [44] K. Otani, T. Tanigawa, T. Watanabe et al., "Toll-like receptor 9 signaling has anti-inflammatory effects on the early phase of Helicobacter pylori-induced gastritis," *Biochemical and*

- Biophysical Research Communications*, vol. 426, no. 3, pp. 342–349, 2012.
- [45] L. Alvarez-Arellano, P. Cortés-Reynosa, N. Sánchez-Zauco, E. Salazar, J. Torres, and C. Maldonado-Bernal, “TLR9 and NF- $\kappa$ B are partially involved in activation of human neutrophils by *Helicobacter pylori* and its purified DNA,” *PLoS One*, vol. 9, no. 7, article e101342, 2014.
- [46] Y. J. Chang, M. S. Wu, J. T. Lin et al., “Induction of cyclooxygenase-2 overexpression in human gastric epithelial cells by *Helicobacter pylori* involves TLR2/TLR9 and c-Src-dependent nuclear factor-kappaB activation,” *Molecular Pharmacology*, vol. 66, no. 6, pp. 1465–1477, 2004.
- [47] T. B. Geijtenbeek and S. I. Gringhuis, “C-type lectin receptors in the control of T helper cell differentiation,” *Nature Reviews Immunology*, vol. 16, no. 7, pp. 433–448, 2016.
- [48] M. Nagata, K. Toyonaga, E. Ishikawa et al., “*Helicobacter pylori* metabolites exacerbate gastritis through C-type lectin receptors,” *The Journal of Experimental Medicine*, vol. 218, no. 1, article 20200815, 2021.
- [49] M. S. M. Timmer, T. J. Teunissen, K. Kodar, A. J. Foster, S. Yamasaki, and B. L. Stocker, “Cholesteryl glucosides signal through the carbohydrate recognition domain of the macrophage inducible C-type lectin mincle,” *Organic & Biomolecular Chemistry*, vol. 19, no. 10, pp. 2198–2202, 2021.
- [50] B. J. Appelmelk, I. van Die, S. J. van Vliet, C. M. Vandembroucke-Grauls, T. B. Geijtenbeek, and Y. van Kooyk, “Cutting edge: carbohydrate profiling identifies new pathogens that interact with dendritic cell-specific ICAM-3-grabbing nonintegrin on dendritic cells,” *Journal of Immunology*, vol. 170, no. 4, pp. 1635–1639, 2003.
- [51] Y. van Kooyk and T. B. Geijtenbeek, “DC-SIGN: escape mechanism for pathogens,” *Nature Reviews Immunology*, vol. 3, no. 9, pp. 697–709, 2003.
- [52] P. Conde, M. Rodriguez, W. van der Touw et al., “DC-SIGN<sup>+</sup> Macrophages Control the Induction of Transplantation Tolerance,” *Immunity*, vol. 42, no. 6, pp. 1143–1158, 2015.
- [53] M. P. Bergman, A. Engering, H. H. Smits et al., “*Helicobacter pylori* modulates the T helper cell 1/T helper cell 2 balance through phase-variable interaction between lipopolysaccharide and DC-SIGN,” *The Journal of Experimental Medicine*, vol. 200, no. 8, pp. 979–990, 2004.
- [54] S. I. Gringhuis, J. den Dunnen, M. Litjens, M. van der Vliet, and T. B. Geijtenbeek, “Carbohydrate-specific signaling through the DC-SIGN signalosome tailors immunity to *Mycobacterium tuberculosis*, HIV-1 and *Helicobacter pylori*,” *Nature Immunology*, vol. 10, no. 10, pp. 1081–1088, 2009.
- [55] T. A. Barr, S. Brown, G. Ryan, J. Zhao, and D. Gray, “TLR-mediated stimulation of APC: distinct cytokine responses of B cells and dendritic cells,” *European Journal of Immunology*, vol. 37, no. 11, pp. 3040–3053, 2007.
- [56] R. Rad, L. Brenner, A. Krug et al., “Toll-Like Receptor-Dependent Activation of Antigen-Presenting Cells Affects Adaptive Immunity to *Helicobacter pylori*,” *Gastroenterology*, vol. 133, no. 1, pp. 150–163.e3, 2007.
- [57] V. Lampropoulou, K. Hoehlig, T. Roch et al., “TLR-activated B cells suppress T cell-mediated autoimmunity,” *Journal of Immunology*, vol. 180, no. 7, pp. 4763–4773, 2008.
- [58] D. Altunöz and Y. A. Sayi, “*Helicobacter*-stimulated IL-10-producing B cells suppress differentiation of lipopolysaccharide/*Helicobacter felis*-activated stimulatory dendritic cells,” *Turkish journal of biology = Turk biyoloji dergisi*, vol. 45, no. 2, pp. 214–224, 2021.
- [59] A. Sayi, E. Kohler, I. M. Toller et al., “TLR-2-activated B cells suppress *Helicobacter*-induced preneoplastic gastric immunopathology by inducing T regulatory-1 cells. Journal of immunology,” *Journal of Immunology*, vol. 186, pp. 878–890, 2011.
- [60] B. Schmausser, M. Andrulis, S. Endrich et al., “Expression and subcellular distribution of toll-like receptors TLR4, TLR5 and TLR9 on the gastric epithelium in *Helicobacter pylori* infection,” *Clinical and Experimental Immunology*, vol. 136, no. 3, pp. 521–526, 2004.
- [61] B. Schmausser, M. Andrulis, S. Endrich, H. K. Müller-Hermelink, and M. Eck, “Toll-like receptors TLR4, TLR5 and TLR9 on gastric carcinoma cells: An implication for interaction with *Helicobacter pylori*,” *International Journal Of Medical Microbiology*, vol. 295, no. 3, pp. 179–185, 2005.
- [62] C. C. Allison, J. Ferrand, L. McLeod et al., “Nucleotide oligomerization domain 1 enhances IFN- $\gamma$  signaling in gastric epithelial cells during *Helicobacter pylori* infection and exacerbates disease severity,” *Journal of Immunology*, vol. 190, no. 7, pp. 3706–3715, 2013.
- [63] L. Ying and R. L. Ferrero, “Role of NOD1 and ALPK1/TIFA signalling in innate immunity against *Helicobacter pylori* infection,” *Current Topics in Microbiology and Immunology*, vol. 421, pp. 159–177, 2019.
- [64] H. Lagunes-Servin, J. Torres, C. Maldonado-Bernal et al., “Toll-like receptors and cytokines are upregulated during *Helicobacter pylori* infection in children,” *Helicobacter*, vol. 18, no. 6, pp. 423–432, 2013.
- [65] P. Pimentel-Nunes, L. Afonso, P. Lopes et al., “Increased expression of toll-like receptors (TLR) 2, 4 and 5 in gastric dysplasia,” *Pathology oncology research: POR*, vol. 17, no. 3, pp. 677–683, 2011.
- [66] A. C. Cadamuro, A. F. Rossi, J. Matos Biselli-Périco et al., “Effect of *Helicobacter pylori* eradication on TLR2 and TLR4 expression in patients with gastric lesions,” *Mediators of Inflammation*, vol. 2015, 9 pages, 2015.
- [67] P. Pimentel-Nunes, N. Gonçalves, I. Boal-Carvalho et al., “*Helicobacter pylori* induces increased expression of Toll-like receptors and decreased Toll-interacting protein in gastric mucosa that persists throughout gastric carcinogenesis,” *Helicobacter*, vol. 18, no. 1, pp. 22–32, 2013.
- [68] K. Uno, K. Kato, T. Atsumi et al., “Toll-like receptor (TLR) 2 induced through TLR4 signaling initiated by *Helicobacter pylori* cooperatively amplifies iNOS induction in gastric epithelial cells,” *American Journal of Physiology Gastrointestinal and Liver Physiology*, vol. 293, no. 5, pp. G1004–G1012, 2007.
- [69] M. G. Varga, C. L. Shaffer, J. C. Sierra et al., “Pathogenic *Helicobacter pylori* strains translocate DNA and activate TLR9 via the cancer-associated *cag* type IV secretion system,” *Oncogene*, vol. 35, no. 48, pp. 6262–6269, 2016.
- [70] Y. Chen, J. Huang, and C. Xu, “Lipopolysaccharide-induced DC-SIGN/TLR4 crosstalk activates NLRP3 inflammasomes via MyD88-independent signaling in gastric epithelial cells,” *Experimental Cell Research*, vol. 396, no. 1, article 112292, 2020.
- [71] J. Wu, K. Lin, J. Zeng et al., “Role of DC-SIGN in *Helicobacter pylori* infection of gastrointestinal cells,” *Frontiers in Bioscience*, vol. 19, no. 5, pp. 825–834, 2014.

- [72] J. Viala, C. Chaput, I. G. Boneca et al., “Nod 1 responds to peptidoglycan delivered by the *Helicobacter pylori* cag pathogenicity island,” *Nature Immunology*, vol. 5, no. 11, pp. 1166–1174, 2004.
- [73] M. L. Hutton, M. Kaparakis-Liaskos, L. Turner, A. Cardona, T. Kwok, and R. L. Ferrero, “*Helicobacter pylori* exploits cholesterol-rich microdomains for induction of NF-kappaB-dependent responses and peptidoglycan delivery in epithelial cells,” *Infection and Immunity*, vol. 78, no. 11, pp. 4523–4531, 2010.
- [74] P. K. Boughan, R. H. Argent, M. Body-Malapel et al., “Nucleotide-binding Oligomerization Domain-1 and Epidermal Growth Factor Receptor:,” *The Journal of Biological Chemistry*, vol. 281, no. 17, pp. 11637–11648, 2006.
- [75] T. Watanabe, N. Asano, A. Kitani, I. J. Fuss, T. Chiba, and W. Strober, “Activation of type I IFN signaling by NOD1 mediates mucosal host defense against *Helicobacter pylori* infection,” *Gut Microbes*, vol. 2, no. 1, pp. 61–65, 2011.
- [76] M. F. González, P. Díaz, A. Sandoval-Bórquez, D. Herrera, and A. F. G. Quest, “*Helicobacter pylori* outer membrane vesicles and extracellular vesicles from *Helicobacter pylori*-infected cells in gastric disease development,” *International Journal of Molecular Sciences*, vol. 22, no. 9, p. 4823, 2021.
- [77] M. Kaparakis, L. Turnbull, L. Carneiro et al., “Bacterial membrane vesicles deliver peptidoglycan to NOD1 in epithelial cells,” *Cellular Microbiology*, vol. 12, no. 3, pp. 372–385, 2010.
- [78] S. Zimmermann, L. Pfannkuch, M. A. Al-Zeer et al., “ALPK1- and TIFA-Dependent Innate Immune Response Triggered by the *Helicobacter pylori* Type IV Secretion System,” *Cell Reports*, vol. 20, no. 10, pp. 2384–2395, 2017.
- [79] L. Pfannkuch, R. Hurwitz, J. Traulsen et al., “ADP heptose, a novel pathogen-associated molecular pattern identified in *Helicobacter pylori*,” *FASEB Journal*, vol. 33, no. 8, pp. 9087–9099, 2019.
- [80] P. Zhou, Y. She, N. Dong et al., “Alpha-kinase 1 is a cytosolic innate immune receptor for bacterial ADP- heptose,” *Nature*, vol. 561, no. 7721, pp. 122–126, 2018.
- [81] A. Gall, R. G. Gaudet, S. D. Gray-Owen, and N. R. Salama, “TIFA signaling in gastric epithelial cells initiates the cag type 4 secretion system-dependent innate immune response to *Helicobacter pylori* infection,” *mBio*, vol. 8, no. 4, article e01168-17, 2017.
- [82] J. Zindel and P. Kubers, “DAMPs, PAMPs, and LAMPs in immunity and sterile inflammation,” *Annual Review of Pathology*, vol. 15, no. 1, pp. 493–518, 2020.
- [83] C. Gabay, C. Lamacchia, and G. Palmer, “IL-1 pathways in inflammation and human diseases,” *Nature Reviews Rheumatology*, vol. 6, no. 4, pp. 232–241, 2010.
- [84] A. Denes, G. Lopez-Castejon, and D. Brough, “Caspase-1: is IL-1 just the tip of the ICEberg?,” *Cell Death & Disease*, vol. 3, no. 7, article e338, 2012.
- [85] T. Strowig, J. Henao-Mejia, E. Elinav, and R. Flavell, “Inflammasomes in health and disease,” *Nature*, vol. 481, no. 7381, pp. 278–286, 2012.
- [86] I. Hitzler, A. Sayi, E. Kohler et al., “Caspase-1 has both proinflammatory and regulatory properties in *Helicobacter* infections, which are differentially mediated by its substrates IL-1 $\beta$  and IL-18,” *Journal of Immunology*, vol. 188, no. 8, pp. 3594–3602, 2012.
- [87] T. Fernandes-Alnemri, J. W. Yu, P. Datta, J. Wu, and E. S. Alnemri, “AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA,” *Nature*, vol. 458, no. 7237, pp. 509–513, 2009.
- [88] N. J. Bitto, P. J. Baker, J. K. Dowling et al., “Membrane vesicles from *Pseudomonas aeruginosa* activate the noncanonical inflammasome through caspase-5 in human monocytes,” *Immunology and Cell Biology*, vol. 96, no. 10, pp. 1120–1130, 2018.
- [89] O. Gutierrez, C. Pipaon, and J. L. Fernandez-Luna, “Ipaf is upregulated by tumor necrosis factor-alpha in human leukemia cells,” *FEBS Letters*, vol. 568, no. 1-3, pp. 79–82, 2004.
- [90] R. P. Semper, M. Vieth, M. Gerhard, and R. Mejías-Luque, “*Helicobacter pylori* Exploits the NLRC4 inflammasome to dampen host defenses,” *Journal of Immunology*, vol. 203, no. 8, pp. 2183–2193, 2019.
- [91] D. Sharma and T. D. Kanneganti, “The cell biology of inflammasomes: mechanisms of inflammasome activation and regulation,” *The Journal of Cell Biology*, vol. 213, no. 6, pp. 617–629, 2016.
- [92] M. S. J. Mangan, E. J. Olhava, W. R. Roush, H. M. Seidel, G. D. Glick, and E. Latz, “Targeting the NLRP3 inflammasome in inflammatory diseases,” *Drug discovery*, vol. 17, no. 8, pp. 588–606, 2018.
- [93] A. R. Jang, M. J. Kang, J. I. Shin et al., “Unveiling the crucial role of type IV secretion system and motility of *Helicobacter pylori* in IL-1 $\beta$  production via NLRP3 inflammasome activation in neutrophils,” *Frontiers in Immunology*, vol. 11, p. 1121, 2020.
- [94] E. Pérez-Figueroa, J. Torres, N. Sánchez-Zauco, A. Contreras-Ramos, L. Alvarez-Arellano, and C. Maldonado-Bernal, “Activation of NLRP3 inflammasome in human neutrophils by *Helicobacter pylori* infection,” *Innate Immunity*, vol. 22, no. 2, pp. 103–112, 2016.
- [95] S. Kameoka, T. Kameyama, T. Hayashi et al., “*Helicobacter pylori* induces IL-1 $\beta$  protein through the inflammasome activation in differentiated macrophagic cells,” *Biomedical Research*, vol. 37, no. 1, pp. 21–27, 2016.
- [96] D. J. Kim, J. H. Park, L. Franchi, S. Backert, and G. Núñez, “The Cag pathogenicity island and interaction between TLR2/NOD2 and NLRP3 regulate IL-1 $\beta$  production in *Helicobacter pylori* infected dendritic cells,” *European Journal of Immunology*, vol. 43, no. 10, pp. 2650–2658, 2013.
- [97] X. Li, S. Liu, J. Luo et al., “*Helicobacter pylori* induces IL-1 $\beta$  and IL-18 production in human monocytic cell line through activation of NLRP3 inflammasome via ROS signaling pathway,” *Pathogens and Disease*, vol. 73, no. 4, 2015.
- [98] K. N. Koch and A. Müller, “*Helicobacter pylori* activates the TLR2/NLRP3/caspase-1/IL-18 axis to induce regulatory T-cells, establish persistent infection and promote tolerance to allergens,” *Gut Microbes*, vol. 6, no. 6, pp. 382–387, 2015.
- [99] S. K. Pachathundikandi, N. Blaser, H. Bruns, and S. Backert, “*Helicobacter pylori* avoids the critical activation of NLRP3 inflammasome-mediated production of oncogenic mature IL-1 $\beta$  in human immune cells,” *SSCancers*, vol. 12, no. 4, p. 803, 2020.
- [100] S. I. Grivennikov, A. V. Tumanov, D. J. Liepinsh et al., “Distinct and nonredundant in vivo functions of TNF produced by t cells and macrophages/neutrophils: protective and deleterious effects,” *Immunity*, vol. 22, no. 1, pp. 93–104, 2005.
- [101] H. Wajant and D. Siegmund, “TNFR1 and TNFR2 in the control of the life and death balance of macrophages,” *Frontiers in cell and developmental biology*, vol. 7, p. 91, 2019.



- [102] P. D. Issuree, T. Marezky, D. R. McIlwain et al., “iRHOM2 is a critical pathogenic mediator of inflammatory arthritis,” *The Journal of Clinical Investigation*, vol. 123, no. 2, pp. 928–932, 2013.
- [103] K. Horiuchi, T. Kimura, T. Miyamoto et al., “Cutting edge: TNF- $\alpha$ -converting enzyme (TACE/ADAM17) inactivation in mouse myeloid cells prevents lethality from endotoxin shock,” *Journal of Immunology*, vol. 179, no. 5, pp. 2686–2689, 2007.
- [104] E. Varfolomeev and D. Vucic, “Intracellular regulation of TNF activity in health and disease,” *Cytokine*, vol. 101, pp. 26–32, 2018.
- [105] H. Zelová and J. Hošek, “TNF- $\alpha$  signalling and inflammation: interactions between old acquaintances,” *Inflammation Research*, vol. 62, no. 7, pp. 641–651, 2013.
- [106] J. R. Bradley, “TNF-mediated inflammatory disease,” *The Journal of Pathology*, vol. 214, no. 2, pp. 149–160, 2008.
- [107] H. Blaser, C. Dostert, T. W. Mak, and D. Brenner, “TNF and ROS crosstalk in inflammation,” *Trends in Cell Biology*, vol. 26, no. 4, pp. 249–261, 2016.
- [108] R. Tavares and S. K. Pathak, “Induction of TNF, CXCL8 and IL-1 $\beta$  in macrophages by *Helicobacter pylori* secreted protein HP1173 occurs via MAP-kinases, NF- $\kappa$ B and AP-1 signaling pathways,” *Microbial Pathogenesis*, vol. 125, pp. 295–305, 2018.
- [109] K. Kranzer, L. Söllner, M. Aigner et al., “Impact of *Helicobacter pylori* virulence factors and compounds on activation and maturation of human dendritic cells,” *Infection and Immunity*, vol. 73, no. 7, pp. 4180–4189, 2005.
- [110] I. L. Beales and J. Calam, “Stimulation of il-8 production in human gastric epithelial cells by *helicobacter pylori*, IL-1 $\beta$  and TNF- $\alpha$  requires tyrosine kinase activity, but not protein kinase C,” *Cytokine*, vol. 9, no. 7, pp. 514–520, 1997.
- [111] B. Neu, A. J. Puschmann, A. Mayerhofer et al., “TNF- $\alpha$  induces apoptosis of parietal cells,” *Biochemical Pharmacology*, vol. 65, no. 10, pp. 1755–1760, 2003.
- [112] G. A. Siregar, S. Halim, and V. R. Sitepu, “Serum TNF- $\alpha$ , IL-8, VEGF levels in *Helicobacter pylori* infection and their association with degree of gastritis,” *Acta Medica Indonesiana*, vol. 47, no. 2, pp. 120–126, 2015.
- [113] J. Shibata, H. Goto, T. Arisawa et al., “Regulation of tumour necrosis factor (TNF) induced apoptosis by soluble TNF receptors in *Helicobacter pylori* infection,” *Gut*, vol. 45, no. 1, pp. 24–31, 1999.
- [114] G. Bounder, “Association of tumor necrosis factor receptor 1 promoter gene polymorphisms (-580 A/G and -609 G/T) and TNFR1 serum levels with the susceptibility to gastric precancerous lesions and gastric cancer related to *H. pylori* infection in a Moroccan population,” *BioMed Research International*, vol. 2020, Article ID 2451854, 7 pages, 2020.
- [115] M. J. Morgan and Z. G. Liu, “Crosstalk of reactive oxygen species and NF- $\kappa$ B signaling,” *Cell Research*, vol. 21, no. 1, pp. 103–115, 2011.
- [116] S. Nakajima and M. Kitamura, “Bidirectional regulation of NF- $\kappa$ B by reactive oxygen species: a role of unfolded protein response,” *Free Radical Biology & Medicine*, vol. 65, pp. 162–174, 2013.
- [117] K. Turpaev, A. Glatigny, J. Bignon, H. Delacroix, and J. C. Drapier, “Variation in gene expression profiles of human monocytic U937 cells exposed to various fluxes of nitric oxide,” *Free Radical Biology & Medicine*, vol. 48, no. 2, pp. 298–305, 2010.
- [118] D. A. Wink and J. B. Mitchell, “Chemical biology of nitric oxide: insights into regulatory, cytotoxic, and cytoprotective mechanisms of nitric oxide,” *Free Radical Biology & Medicine*, vol. 25, no. 4–5, pp. 434–456, 1998.
- [119] T. Shimizu, T. Chiba, and H. Marusawa, “*Helicobacter pylori*-mediated genetic instability and gastric carcinogenesis,” *Current Topics in Microbiology and Immunology*, vol. 400, pp. 305–323, 2017.
- [120] M. Katsurahara, Y. Kobayashi, M. Iwasa et al., “Reactive nitrogen species mediate DNA damage in *Helicobacter pylori*-infected gastric mucosa,” *Helicobacter*, vol. 14, no. 6, pp. 552–558, 2009.
- [121] S. K. Tiwari, A. S. Shaik, A. P. Shaik et al., “Gene expression patterns of COX-1, COX-2 and iNOS in *H. Pylori* infected histopathological conditions,” *Microbial Pathogenesis*, vol. 135, p. 103634, 2019.
- [122] O. Handa, Y. Naito, and T. Yoshikawa, “*Helicobacter pylori*: a ROS-inducing bacterial species in the stomach,” *Inflammation Research*, vol. 59, no. 12, pp. 997–1003, 2010.
- [123] K. D. Molecular, “Mechanisms of *H. pylori*-induced DNA double-strand breaks,” *International Journal of Molecular Sciences*, p. 19, 2018.
- [124] L. D. Butcher, G. den Hartog, P. B. Ernst, and S. E. Crowe, “Oxidative Stress Resulting From *Helicobacter pylori* Infection Contributes to Gastric Carcinogenesis,” *Cellular and Molecular Gastroenterology and Hepatology*, vol. 3, no. 3, pp. 316–322, 2017.
- [125] G. Y. Chen and G. Nuñez, “Sterile inflammation: sensing and reacting to damage,” *Nature Reviews Immunology*, vol. 10, no. 12, pp. 826–837, 2010.
- [126] A. P. Gobert and K. T. Wilson, “The immune battle against *Helicobacter pylori* infection: NO offense,” *Trends in Microbiology*, vol. 24, no. 5, pp. 366–376, 2016.
- [127] T. N. Mayadas, X. Cullere, and C. A. Lowell, “The multifaceted functions of neutrophils,” *Annual Review of Pathology*, vol. 9, no. 1, pp. 181–218, 2014.
- [128] M. N. Ajuebor, A. M. Das, L. Virág, R. J. Flower, C. Szabó, and M. Perretti, “Role of resident peritoneal macrophages and mast cells in chemokine production and neutrophil migration in acute inflammation: evidence for an inhibitory loop involving endogenous IL-10,” *Journal of Immunology*, vol. 162, pp. 1685–1691, 1999.
- [129] N. D. Kim and A. D. Luster, “The role of tissue resident cells in neutrophil recruitment,” *Trends in Immunology*, vol. 36, no. 9, pp. 547–555, 2015.
- [130] S. M. Vieira, H. P. Lemos, R. Grespan et al., “A crucial role for TNF- $\alpha$  in mediating neutrophil influx induced by endogenously generated or exogenous chemokines, KC/CXCL1 and LIX/CXCL5,” *British Journal of Pharmacology*, vol. 158, no. 3, pp. 779–789, 2009.
- [131] S. Suerbaum and P. Michetti, “*Helicobacter pylori* Infection,” *The New England Journal of Medicine*, vol. 347, no. 15, pp. 1175–1186, 2002.
- [132] H. W. Fu, “*Helicobacter pylori* neutrophil-activating protein: from molecular pathogenesis to clinical applications,” *World Journal of Gastroenterology*, vol. 20, no. 18, pp. 5294–5301, 2014.
- [133] M. Brisslert, K. Enarsson, S. Lundin et al., “*Helicobacter pylori* induce neutrophil transendothelial migration: role of

- the bacterial HP-NAP," *FEMS Microbiology Letters*, vol. 249, no. 1, pp. 95–103, 2005.
- [134] C. A. Wang, Y. C. Liu, S. Y. Du, C. W. Lin, and H. W. Fu, "Helicobacter pylori neutrophil-activating protein promotes myeloperoxidase release from human neutrophils," *Biochemical and Biophysical Research Communications*, vol. 377, no. 1, pp. 52–56, 2008.
- [135] M. M. D'Elia, A. Amedei, A. Cappon, G. Del Prete, and M. de Bernard, "The neutrophil-activating protein of Helicobacter pylori (HP-NAP) as an immune modulating agent," *FEMS Immunology and Medical Microbiology*, vol. 50, no. 2, pp. 157–164, 2007.
- [136] M. Unemo, M. Aspholm-Hurtig, D. Ilver et al., "The Sialic Acid Binding SabA Adhesin of Helicobacter pylori Is essential for nonopsonic activation of human neutrophils," *The Journal of Biological Chemistry*, vol. 280, no. 15, pp. 15390–15397, 2005.
- [137] T. H. Chu, S. T. Huang, S. F. Yang et al., "Hepatoma-derived growth factor participates in Helicobacter Pylori-induced neutrophils recruitment, gastritis and gastric carcinogenesis," *Oncogene*, vol. 38, no. 37, pp. 6461–6477, 2019.
- [138] J. Matsuzaki, H. Tsugawa, Y. Kashiwazaki et al., "Neutrophil-activating protein polymorphism of Helicobacter pylori determines the host risk of dyspepsia," *Cellular and Molecular Gastroenterology and Hepatology*, vol. 8, no. 2, pp. 295–297.e6, 2019.
- [139] M. de Bernard and M. M. D'Elia, "The immune modulating activity of the Helicobacter pylori HP-NAP: Friend or foe?," *Toxicon*, vol. 56, no. 7, pp. 1186–1192, 2010.
- [140] K. Futosi, S. Fodor, and A. Mócsai, "Neutrophil cell surface receptors and their intracellular signal transduction pathways," *International Immunopharmacology*, vol. 17, no. 3, pp. 638–650, 2013.
- [141] C. J. Thomas and K. Schroder, "Pattern recognition receptor function in neutrophils," *Trends in Immunology*, vol. 34, no. 7, pp. 317–328, 2013.
- [142] Y. Wang and F. Jönsson, "Expression, role, and regulation of neutrophil Fcγ receptors," *Frontiers in Immunology*, vol. 10, p. 1958, 2019.
- [143] T. Lämmermann and W. Kastner, "Concepts of GPCR-controlled navigation in the immune system," *Immunological Reviews*, vol. 289, no. 1, pp. 205–231, 2019.
- [144] A. Mantovani, M. A. Cassatella, C. Costantini, and S. Jaillon, "Neutrophils in the activation and regulation of innate and adaptive immunity," *Nature Reviews Immunology*, vol. 11, no. 8, pp. 519–531, 2011.
- [145] B. L. Johnson III, J. W. Kuethe, and C. C. Caldwell, "Neutrophil derived microvesicles: emerging role of a key mediator to the immune response," *Endocrine, Metabolic & Immune Disorders Drug Targets*, vol. 14, no. 3, pp. 210–217, 2014.
- [146] F. A. Bonilla and H. C. Oettgen, "Adaptive immunity," *The Journal of Allergy and Clinical Immunology*, vol. 125, no. 2, pp. S33–S40, 2010.
- [147] S. Jiang and C. Dong, "A complex issue on CD4<sup>+</sup> T-cell subsets," *Immunological Reviews*, vol. 252, no. 1, pp. 5–11, 2013.
- [148] J. Zhu and W. E. Paul, "Peripheral CD4<sup>+</sup> T-cell differentiation regulated by networks of cytokines and transcription factors," *Immunological Reviews*, vol. 238, no. 1, pp. 247–262, 2010.
- [149] Y. Shi, X. F. Liu, Y. Zhuang et al., "Helicobacter pylori-induced Th17 responses modulate Th1 cell responses, benefit bacterial growth, and contribute to pathology in mice," *Journal of Immunology*, vol. 184, no. 9, pp. 5121–5129, 2010.
- [150] N. Bagheri, L. Salimzadeh, and H. Shirzad, "The role of T helper 1-cell response in Helicobacter pylori-infection," *Microbial Pathogenesis*, vol. 123, pp. 1–8, 2018.
- [151] B. Gebert, W. Fischer, E. Weiss, R. Hoffmann, and R. Haas, "Helicobacter pylori vacuolating cytotoxin inhibits T lymphocyte activation," *Science*, vol. 301, no. 5636, pp. 1099–1102, 2003.
- [152] C. Schmees, C. Prinz, T. Treptau et al., "Inhibition of T-Cell Proliferation by Helicobacter pylori γ-Glutamyl Transpeptidase," *Gastroenterology*, vol. 132, no. 5, pp. 1820–1833, 2007.
- [153] V. Supajatura, H. Ushio, A. Wada et al., "Cutting edge: VacA, a vacuolating cytotoxin of Helicobacter pylori, directly activates mast cells for migration and production of proinflammatory cytokines," *Journal of Immunology*, vol. 168, no. 6, pp. 2603–2607, 2002.
- [154] M. de Bernard, A. Cappon, L. Pancotto et al., "The Helicobacter pylori VacA cytotoxin activates RBL-2H3 cells by inducing cytosolic calcium oscillations," *Cellular Microbiology*, vol. 7, no. 2, pp. 191–198, 2005.
- [155] K. M. Kim, S. G. Lee, M. G. Park et al., "γ-Glutamyltranspeptidase of Helicobacter pylori induces mitochondria-mediated apoptosis in AGS cells," *Biochemical and Biophysical Research Communications*, vol. 355, no. 2, pp. 562–567, 2007.
- [156] M. Gong, S. S. Ling, S. Y. Lui, K. G. Yeoh, and B. Ho, "Helicobacter pylori γ-Glutamyl transpeptidase is a pathogenic factor in the development of peptic ulcer disease," *Gastroenterology*, vol. 139, no. 2, pp. 564–573, 2010.
- [157] E. Ricciotti and G. A. FitzGerald, "Prostaglandins and inflammation," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 31, no. 5, pp. 986–1000, 2011.
- [158] A. Lundgren, E. Strömberg, Å. Sjöling et al., "Mucosal FOXP3-expressing CD4<sup>+</sup> CD25<sup>high</sup> regulatory T cells in Helicobacter pylori-infected patients," *Infection and Immunity*, vol. 73, no. 1, pp. 523–531, 2005.
- [159] A. Kandulski, T. Wex, D. Kuester et al., "Naturally occurring regulatory T cells (CD4<sup>+</sup>, CD25<sup>high</sup>, FOXP3<sup>+</sup>) in the antrum and cardia are associated with higher H. pylori colonization and increased gene expression of TGF-β1," *Helicobacter*, vol. 13, no. 4, pp. 295–303, 2008.
- [160] P. R. Harris, S. W. Wright, C. Serrano et al., "Helicobacter pylori gastritis in children is associated with a regulatory T-cell response," *Gastroenterology*, vol. 134, no. 2, pp. 491–499, 2008.
- [161] J. Y. Kao, M. Zhang, M. J. Miller et al., "Helicobacter pylori immune escape is mediated by dendritic cell-induced treg skewing and Th17 suppression in mice," *Gastroenterology*, vol. 138, no. 3, pp. 1046–1054, 2010.
- [162] M. Oertli, M. Sundquist, I. Hitzler et al., "DC-derived IL-18 drives Treg differentiation, murine helicobacter pylori-specific immune tolerance, and asthma protection," *The Journal of Clinical Investigation*, vol. 122, no. 3, pp. 1082–1096, 2012.
- [163] M. Oertli, M. Noben, D. B. Engler et al., "Helicobacter pylori γ-glutamyl transpeptidase and vacuolating cytotoxin promote gastric persistence and immune tolerance," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 110, no. 8, pp. 3047–3052, 2013.

- [164] D. A. Vignali and V. K. Kuchroo, "IL-12 family cytokines: immunological playmakers," *Nature Immunology*, vol. 13, no. 8, pp. 722–728, 2012.
- [165] D. Ashour, P. Arampatzis, V. Pavlovic et al., "IL-12 from endogenous cDC1, and not vaccine DC, is required for Th1 induction," *JCI insight*, vol. 5, no. 10, 2020.
- [166] W. T. Watford, M. Moriguchi, A. Morinobu, and J. J. O'Shea, "The biology of IL-12: coordinating innate and adaptive immune responses," *Cytokine & Growth Factor Reviews*, vol. 14, no. 5, pp. 361–368, 2003.
- [167] E. D. Tait Wojno, C. A. Hunter, and J. S. Stumhofer, "The immunobiology of the interleukin-12 family: room for discovery," *Immunity*, vol. 50, no. 4, pp. 851–870, 2019.
- [168] C. L. Langrish, B. S. McKenzie, N. J. Wilson, M. R. de Waal, R. A. Kastelein, and D. J. Cua, "IL-12 and IL-23: master regulators of innate and adaptive immunity," *Immunological Reviews*, vol. 202, no. 1, pp. 96–105, 2004.
- [169] E. M. Gravallese and G. Schett, "Effects of the IL-23-IL-17 pathway on bone in spondyloarthritis," *Nature Reviews Rheumatology*, vol. 14, no. 11, pp. 631–640, 2018.
- [170] T. Korn, E. Bettelli, M. Oukka, and V. K. Kuchroo, "IL-17 and Th17 cells," *Annual Review of Immunology*, vol. 27, no. 1, pp. 485–517, 2009.
- [171] B. S. McKenzie, R. A. Kastelein, and D. J. Cua, "Understanding the IL-23-IL-17 immune pathway," *Trends in Immunology*, vol. 27, no. 1, pp. 17–23, 2006.
- [172] J. K. Kolls and A. Lindén, "Interleukin-17 family members and inflammation," *Immunity*, vol. 21, no. 4, pp. 467–476, 2004.
- [173] S. Pflanz, J. C. Timans, J. Cheung et al., "IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4<sup>+</sup> T cells," *Immunity*, vol. 16, no. 6, pp. 779–790, 2002.
- [174] H. Yoshida and C. A. Hunter, "The immunobiology of interleukin-27," *Annual Review of Immunology*, vol. 33, pp. 417–443, 2015.
- [175] A. O. Hall, D. P. Beiting, C. Tato et al., "The cytokines interleukin 27 and interferon- $\gamma$  promote distinct Treg cell populations required to limit infection-induced pathology," *Immunity*, vol. 37, no. 3, pp. 511–523, 2012.
- [176] J. Zhang, Y. Zhang, Q. Wang et al., "Interleukin-35 in immune-related diseases: protection or destruction," *Immunology*, vol. 157, no. 1, pp. 13–20, 2019.
- [177] L. W. Collison, C. J. Workman, T. T. Kuo et al., "The inhibitory cytokine IL-35 contributes to regulatory T-cell function," *Nature*, vol. 450, no. 7169, pp. 566–569, 2007.
- [178] W. Ouyang, S. Rutz, N. K. Crellin, P. A. Valdez, and S. G. Hymowitz, "Regulation and functions of the IL-10 family of cytokines in inflammation and disease," *Annual Review of Immunology*, vol. 29, no. 1, pp. 71–109, 2011.
- [179] X. Wang, K. Wong, W. Ouyang, and S. Rutz, "Targeting IL-10 family cytokines for the treatment of human diseases," *Cold Spring Harbor Perspectives in Biology*, vol. 11, no. 2, 2019.
- [180] M. Morikawa, R. Derynck, and K. Miyazono, "TGF- $\beta$  and the TGF- $\beta$  family: context-dependent roles in cell and tissue physiology," *Cold Spring Harbor Perspectives in Biology*, vol. 8, no. 5, 2016.
- [181] M. Kanamori, H. Nakatsukasa, M. Okada, Q. Lu, and A. Yoshimura, "Induced regulatory T cells: their development, stability, and applications," *Trends in Immunology*, vol. 37, no. 11, pp. 803–811, 2016.
- [182] D. Q. Tran, "TGF- $\beta$ : the sword, the wand, and the shield of FOXP3<sup>+</sup> regulatory T cells," *Journal of Molecular Cell Biology*, vol. 4, no. 1, pp. 29–37, 2012.
- [183] M. O. Li, Y. Y. Wan, and R. A. Flavell, "T Cell-Produced Transforming Growth Factor- $\beta$ 1 controls T cell tolerance and regulates TH1- and TH17-cell differentiation," *Immunity*, vol. 26, no. 5, pp. 579–591, 2007.
- [184] K. M. Monroe, S. M. McWhirter, and R. E. Vance, "Induction of type I interferons by bacteria," *Cellular Microbiology*, vol. 12, no. 7, pp. 881–890, 2010.
- [185] G. Trinchieri, "Type I interferon: friend or foe?," *The Journal of Experimental Medicine*, vol. 207, no. 10, pp. 2053–2063, 2010.
- [186] A. Takaoka and H. Yanai, "Interferon signalling network in innate defence," *Cellular Microbiology*, vol. 8, no. 6, pp. 907–922, 2006.
- [187] F. J. Barrat, M. K. Crow, and L. B. Ivashkiv, "Interferon target-gene expression and epigenomic signatures in health and disease," *Nature Immunology*, vol. 20, no. 12, pp. 1574–1583, 2019.
- [188] L. M. Snell, T. L. McGaha, and D. G. Brooks, "Type I Interferon in chronic virus infection and cancer," *Trends in Immunology*, vol. 38, no. 8, pp. 542–557, 2017.
- [189] K. Schroder, P. J. Hertzog, T. Ravasi, and D. A. Hume, "Interferon-gamma: an overview of signals, mechanisms and functions," *Journal of Leukocyte Biology*, vol. 75, no. 2, pp. 163–189, 2004.
- [190] S. M. Soudja, A. L. Ruiz, J. C. Marie, and G. Lauvau, "Inflammatory monocytes activate memory CD8<sup>+</sup> T and Innate NK lymphocytes independent of cognate antigen during microbial pathogen invasion," *Immunity*, vol. 37, no. 3, pp. 549–562, 2012.
- [191] L. B. Ivashkiv, "IFN $\gamma$ : signalling, epigenetics and roles in immunity, metabolism, disease and cancer immunotherapy," *Nature Reviews Immunology*, vol. 18, no. 9, pp. 545–558, 2018.
- [192] C. K. Glass and G. Natoli, "Molecular control of activation and priming in macrophages," *Nature Immunology*, vol. 17, no. 1, pp. 26–33, 2016.
- [193] R. S. Goldszmid, P. Caspar, A. Rivollier et al., "NK cell-derived interferon- $\gamma$  orchestrates cellular dynamics and the differentiation of monocytes into dendritic cells at the site of infection," *Immunity*, vol. 36, no. 6, pp. 1047–1059, 2012.