

Research Article

Solving Nonlinear Boundary Value Problems Using He's Polynomials and Padé Approximants

Syed Tauseef Mohyud-Din^{1,2} and Ahmet Yildirim³

¹ Institute of Aerospace & Avionics, Air University, Sector E-9, 44000 Islamabad, Pakistan

² COMSATS Institute of Information Technology, Lahore, Pakistan

³ Department of Mathematics, Science Faculty, Ege University, 35100 Bornova İzmir, Turkey

Correspondence should be addressed to Syed Tauseef Mohyud-Din, syedtauseefs@hotmail.com

Received 5 August 2009; Accepted 12 September 2009

Recommended by Ji Huan He

We apply He's polynomials coupled with the diagonal Padé approximants for solving various singular and nonsingular boundary value problems which arise in engineering and applied sciences. The diagonal Padé approximants prove to be very useful for the understanding of physical behavior of the solution. Numerical results reveal the complete reliability of the proposed combination.

Copyright © 2009 S. T. Mohyud-Din and A. Yildirim. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

With the rapid development of nonlinear sciences, many analytical and numerical techniques have been developed by various scientists for solving singular and nonsingular initial and boundary value problems which arise in the mathematical modeling of diversified physical problems related to engineering and applied sciences. The application of these problems involves physics, astrophysics, experimental and mathematical physics, nuclear charge in heavy atoms, thermal behavior of a spherical cloud of gas, thermodynamics, population models, chemical kinetics, and fluid mechanics see [1–68] and the references therein. Several techniques [1–68] including decomposition, variational iteration, finite difference, polynomial spline, differential transform, exp-function and homotopy perturbation have been developed for solving such problems. Most of these methods have their inbuilt deficiencies coupled with the major drawback of huge computational work. He [19–24] developed the homotopy perturbation method (HPM) for solving linear, nonlinear, initial and boundary value problems. The homotopy perturbation method was formulated by merging the standard homotopy with perturbation. Recently, Ghorbani and Saberi-Nadjafi [15, 16] introduced He's polynomials by splitting the nonlinear term and also proved that He's polynomials are fully compatible with Adomian's polynomials but are easier

to calculate and are more user friendly. The basic motivation of this paper is to apply He's polynomials coupled with the diagonal Padé approximants for solving singular and nonsingular boundary value problems. The Padé approximants are applied in order to make the work more concise and for the better understanding of the solution behavior. The use of Padé approximants shows real promise in solving boundary value problems in an infinite domain; see [42, 50, 56–59]. It is well known in the literature that polynomials are used to approximate the truncated power series. It was observed [42, 50, 56–59] that polynomials tend to exhibit oscillations that may give an approximation error bounds. Moreover, polynomials can never blow up in a finite plane and this makes the singularities not apparent. To overcome these difficulties, the obtained series is best manipulated by Padé approximants for numerical approximations. Using the power series, isolated from other concepts, is not always useful because the radius of convergence of the series may not contain the two boundaries. It is now well known that Padé approximants [42, 50, 56–59] have the advantage of manipulating the polynomial approximation into rational functions of polynomials. By this manipulation, we gain more information about the mathematical behavior of the solution. In addition, the power series are not useful for large values of x . It is an established fact that power series in isolation are not useful to handle boundary value problems. This can be attributed to the possibility that the radius of convergence may not be sufficiently large to contain the boundaries of the domain. It is therefore essential to combine the series solution with the Padé approximants to provide an effective tool to handle boundary value problems on an infinite or semi-infinite domain. We apply this powerful combination of series solution and Padé approximants for solving a variety of boundary value problems. Precisely the proposed combination is applied on boundary layer problem, unsteady flow of gas through a porous medium, Thomas-Fermi equation, Flierl-Petviashvili (FP) equation, and Blasius problem. It is worth mentioning that Flierl-Petviashvili equation has singularity behavior at $x = 0$ which is a difficult element in this type of equations. We transform the FP equation to a first-order initial value problem and He's polynomials are applied to the reformulated first-order initial value problem which leads the solution in terms of transformed variable. The desired series solution is obtained by implementing the inverse transformation. The fact that the proposed algorithm solves nonlinear problems without using Adomian's polynomials is a clear advantage of this technique over the decomposition method.

2. Homotopy Perturbation Method and He's Polynomials

To explain the He's homotopy perturbation method, we consider a general equation of the type

$$L(u) = 0, \quad (2.1)$$

where L is any integral or differential operator. We define a convex homotopy $H(u, p)$ by

$$H(u, p) = (1 - p)F(u) + pL(u), \quad (2.2)$$

where $F(u)$ is a functional operator with known solutions v_0 , which can be obtained easily. It is clear that, for

$$H(u, p) = 0, \quad (2.3)$$

we have

$$H(u, 0) = F(u), \quad H(u, 1) = L(u). \quad (2.4)$$

This shows that $H(u, p)$ continuously traces an implicitly defined curve from a starting point $H(v_0, 0)$ to a solution function $H(f, 1)$. The embedding parameter monotonically increases from zero to unit as the trivial problem $F(u) = 0$, continuously deforms the original problem $L(u) = 0$. The embedding parameter $p \in (0, 1]$ can be considered as an expanding parameter [15, 16, 19–24, 41–50, 60, 63–68]. The homotopy perturbation method uses the homotopy parameter p as an expanding parameter [19–24] to obtain

$$u = \sum_{i=0}^{\infty} p^i u_i = u_0 + pu_1 + p^2 u_2 + p^3 u_3 + \dots, \quad (2.5)$$

if $p \rightarrow 1$, then (2.5) corresponds to (2.2) and becomes the approximate solution of the form

$$f = \lim_{p \rightarrow 1} u = \sum_{i=0}^{\infty} u_i. \quad (2.6)$$

It is well known that series (2.6) is convergent for most of the cases and also the rate of convergence is dependent on $L(u)$; see [19–24]. We assume that (3.2) has a unique solution. The comparisons of like powers of p give solutions of various orders. In sum, according to [15, 16], He's HPM considers the nonlinear term $N(u)$ as

$$N(u) = \sum_{i=0}^{\infty} p^i H_i = H_0 + pH_1 + p^2 H_2 + \dots, \quad (2.7)$$

where H_n 's are the so-called He's polynomials [15, 16], which can be calculated by using the formula

$$H_n(u_0, \dots, u_n) = \frac{1}{n!} \frac{\partial^n}{\partial p^n} \left(N \left(\sum_{i=0}^n p^i u_i \right) \right)_{p=0}, \quad n = 0, 1, 2, \dots \quad (2.8)$$

of various orders.

3. Padé Approximants

A Padé approximant is the ratio of two polynomials constructed from the coefficients of the Taylor series expansion of a function $u(x)$. The $[L/M]$ Padé approximants to a function $y(x)$ are given by [42, 50, 56–59]

$$\left[\frac{L}{M} \right] = \frac{P_L(x)}{Q_M(x)}, \quad (3.1)$$

where $P_L(x)$ is polynomial of degree at most L and $Q_M(x)$ is a polynomial of degree at most M . The formal power series

$$y(x) = \sum_{i=1}^{\infty} a_i x^i, \quad (3.2)$$

$$y(x) - \frac{P_L(x)}{Q_M(x)} = O(x^{L+M+1}) \quad (3.3)$$

determine the coefficients of $P_L(x)$ and $Q_M(x)$ by the equation. Since we can clearly multiply the numerator and denominator by a constant and leave $[L/M]$ unchanged, we imposed the normalization condition

$$Q_M(0) = 1.0. \quad (3.4)$$

Finally, we require that $P_L(x)$ and $Q_M(x)$ have noncommon factors. If we write the coefficient of $P_L(x)$ and $Q_M(x)$ as

$$\begin{aligned} P_L(x) &= p_0 + p_1x + p_2x^2 + \cdots + p_Lx^L, \\ Q_M(x) &= q_0 + q_1x + q_2x^2 + \cdots + q_Mx^M, \end{aligned} \quad (3.5)$$

then by (3.6) and (3.7), we may multiply (3.3) by $Q_M(x)$, which linearizes the coefficient equations. We can write out (3.5) in more details as

$$\begin{aligned} a_{L+1} + a_L q_1 + \cdots + a_{L-M} q_M &= 0, \\ q_{L+2} + q_{L+1} q_1 + \cdots + a_{L-M+2} q_M &= 0, \\ &\vdots \end{aligned} \quad (3.6)$$

$$\begin{aligned} a_{L+M} + a_{L+M-1} q_1 + \cdots + a_L q_M &= 0, \\ a_0 &= p_0, \\ a_0 + a_0 q_1 + \cdots &= p_1, \\ &\vdots \end{aligned} \quad (3.7)$$

$$a_L + a_{L-1} q_1 + \cdots + a_0 q_L = p_L.$$

To solve these equations, we start with (3.6), which is a set of linear equations for all the unknown q 's. Once the q 's are known, then (3.7) gives an explicit formula for the unknown p 's, which complete the solution. If (3.6) and (3.7) are nonsingular, then we can solve them

Table 1: Numerical values for $\alpha = f''(0)$ for $0 < n < 1$ by using diagonal Padé approximants [51, 59].

n	[2/2]	[3/3]	[4/4]	[5/5]	[6/6]
0.2	-0.3872983347	-0.3821533832	-0.3819153845	-0.3819148088	-0.3819121854
1/3	-0.5773502692	-0.5615999244	-0.5614066588	-0.5614481405	-0.561441934
0.4	-0.6451506398	-0.6397000575	-0.6389732578	-0.6389892681	-0.6389734794
0.6	-0.8407967591	-0.8393603021	-0.8396060478	-0.8395875381	-0.8396056769
0.8	-1.007983207	-1.007796981	-1.007646828	-1.007646828	-1.007792100

directly and obtain (3.8) [42, 50, 56–59], where (3.8) holds, and if the lower index on a sum exceeds the upper, the sum is replaced by zero:

$$\left[\frac{L}{M} \right] = \frac{\det \begin{bmatrix} a_{L-M+1} & a_{L-M+2} & \dots & a_{L+1} \\ \vdots & & & \vdots \\ a_L & a_{L+1} & \dots & a_{L+M} \\ \sum_{j=M}^L a_{j-M}x^j & \sum_{j=M-1}^L a_{j-M+1}x^j & \dots & \sum_{j=0}^L a_jx^j \end{bmatrix}}{\det \begin{bmatrix} a_{L-M+1} & a_{L-M+2} & \dots & a_{L+1} \\ \vdots & \vdots & \ddots & \\ a_L & a_{L+1} & \dots & a_{L+M} \\ x^M & x^{M-1} & \dots & 1 \end{bmatrix}}. \tag{3.8}$$

To obtain diagonal Padé approximants of different order such as [2/2], [4/4], or [6/6], we can use the symbolic calculus software Maple.

4. Numerical Applications

In this section, we apply He’s polynomials for solving boundary layer problem, unsteady flow of gas through a porous medium, Thomas-Fermi equation, Flierl-Petviashvili equation, and Blasius problem. The powerful Padé approximants are applied for making the work more concise and to get the better understanding of solution behavior.

Example 4.1 (see [51, 59]). Consider the following nonlinear third-order boundary layer problem which appears mostly in the mathematical modeling of physical phenomena in fluid mechanics [51, 59]

$$f'''(x) + (n - 1)f(x)f''(x) - 2n(f'(x))^2 = 0, \quad n > 0, \tag{4.1}$$

with boundary conditions

$$f(0) = 0, \quad f'(0) = 1, \quad f'(\infty) = 0, \quad n > 0 \tag{4.2}$$

By applying the convex homotopy, we have

$$f_0 + pf_1 + \dots = f_0(x) - p \int \int_0^x \left((n-1)(f_0 + pf_1 + \dots)(f_0'' + pf_1'' + \dots) - 2n(f_0' + pf_1' + \dots)^2 \right) dx dx, \quad n > 0, \quad (4.3)$$

comparing the co-efficient of like powers of p , following approximants are made

$$\begin{aligned} p^{(0)} : f_0(x) &= x, \\ p^{(1)} : f_1(x) &= \frac{1}{2}\alpha x^2 + \frac{1}{3}x^3, \\ p^{(2)} : f_2(x) &= \frac{1}{24}\alpha(3n+1)x^4 + \frac{1}{30}n(n+1)x^5, \\ p^{(3)} : f_3(x) &= \frac{1}{120}\alpha^2(3n+1)x^5 + \frac{1}{720}\alpha(19n^2+18n+3)x^6 + \frac{1}{315}n(2n^2+2n+1)x^7, \\ p^{(4)} : f_4(x) &= \frac{1}{5040}\alpha^2(27n^2+42n+11)x^7 + \frac{1}{40320}\alpha(167n^3+297n^2+161n+15)x^8 \\ &\quad + \frac{1}{22680}n(13n^3+38n^2+23n+6)x^9, \\ &\quad \vdots \end{aligned} \quad (4.4)$$

where $f''(0) = \alpha < 0$ and p^i s are He's polynomials. The series solution is given as

$$\begin{aligned} f(x) &= x + \frac{\alpha x^2}{2} + \frac{nx^3}{3} + \left(\frac{1}{8}n\alpha + \frac{1}{24}\alpha \right) x^4 + \left(\frac{1}{30}n^2 + \frac{1}{40}n\alpha^2 + \frac{1}{120}\alpha^2 + \frac{1}{30}n \right) x^5 \\ &\quad + \left(\frac{19}{720}n^2\alpha + \frac{1}{240}\alpha + \frac{1}{40}n\alpha \right) x^6 \\ &\quad + \left(\frac{1}{120}n\alpha^2 + \frac{1}{315}n + \frac{2}{315}n^3 + \frac{11}{5040}\alpha^2 + \frac{3}{560}n^2\alpha^2 + \frac{2}{315}n^2 \right) x^7 \\ &\quad + \left(\frac{11}{40320}\alpha^3 + \frac{33}{4480}n^2\alpha + \frac{3}{4480}\alpha^3n^2 + \frac{23}{5760}n\alpha + \frac{1}{2688}\alpha + \frac{167}{40320}n^3\alpha + \frac{1}{960}\alpha^3n \right) x^8 \\ &\quad + \left(\frac{1}{3780}n + \frac{527}{362880}n^3\alpha^2 + \frac{19}{11340}n^3 + \frac{709}{362880}n\alpha^2 + \frac{23}{8064}n^2\alpha^2 + \frac{23}{22680}n^2 \right. \\ &\quad \left. + \frac{13}{22680}n^4 + \frac{43}{120960}\alpha^2 \right) x^9 + \dots \end{aligned} \quad (4.5)$$

Table 2: Numerical values for $\alpha = f''(0)$ for $n < 1$ by using diagonal Padé approximants [51, 59].

n	α
4	-2.483954032
10	-4.026385103
100	-12.84334315
1000	-40.65538218
5000	-104.8420672

Example 4.2 (see [51, 57]). Consider the following nonlinear differential equation which governs the unsteady flow of gas through a porous medium

$$y''(x) + \frac{2x}{\sqrt{1-\alpha y}} y'(x) = 0, \quad 0 < \alpha < 1 \quad (4.6)$$

with the following boundary conditions:

$$y(0) = 1, \quad \lim_{x \rightarrow \infty} y(x) = 0. \quad (4.7)$$

By applying the convex homotopy method we have

$$y_0 + p y_1 + \dots = y_0(x) - p \int \int_0^x \left(2x(1-\alpha)(y_0 + p y_1 + p^2 y_2 + \dots)^{-1/2} \right) dx dx. \quad (4.8)$$

By comparing the coefficient of like powers of p , the following approximants are obtained:

$$\begin{aligned} p^{(0)} : y_0(x) &= 1, \\ p^{(1)} : y_1(x) &= Ax, \\ p^{(2)} : y_2(x) &= \frac{A}{3\sqrt{1-\alpha}} x^3, \\ p^{(3)} : y_3(x) &= -\frac{\alpha A^2}{12(1-\alpha)^{3/2}} x^4 + \frac{A}{10(1-\alpha)} x^5, \\ p^{(4)} : y_4(x) &= -\frac{3\alpha^2 A^3}{80(1-\alpha)^{5/2}} x^5 + \frac{\alpha A^2}{15(1-\alpha)^2} x^6 + \dots, \\ &\vdots \end{aligned} \quad (4.9)$$

Table 3: [51, 57].

α	$B_{[2/2]} = y'(0)$	$B_{[3/3]} = y'(0)$
0.1	-3.556558821	-1.957208953
0.2	-2.441894334	-1.786475516
0.3	-1.928338405	-1.478270843
0.4	-1.606856838	-1.231801809
0.5	-1.373178096	-1.025529704
0.6	-1.185519607	-0.8400346085
0.7	-1.021411309	-0.6612047893
0.8	-0.8633400217	-0.4776697286
0.9	-0.6844600642	-0.2772628386

where $A = y'(0)$ and p^i s are He's polynomials. The series solution is given as

$$\begin{aligned}
 y(x) = 1 + Ax - \frac{A}{3\sqrt{1-\alpha}}x^3 - \frac{\alpha A^2}{12(1-\alpha)^{3/2}}x^4 + \left(\frac{A}{10(1-\alpha)} - \frac{3\alpha^2 A^3}{80(1-\alpha)^{5/2}} \right)x^5 \\
 + \left(\frac{\alpha A^2}{15(1-\alpha)^2} - \frac{\alpha^3 A^4}{48(1-\alpha)^{7/2}} \right)x^6 + \dots
 \end{aligned} \tag{4.10}$$

The diagonal Padé approximants [51, 57] can be applied to analyze the physical behavior. Based on this, the [2/2] Padé approximants produced the slope A to be

$$A = -\frac{2(1-\alpha)^{1/4}}{\sqrt{3\alpha}}, \tag{4.11}$$

and by using [3/3] Padé approximants we find

$$A = -\frac{\sqrt{(-4674\alpha + 8664)\sqrt{1-\alpha} - 144\gamma}}{57\alpha}, \tag{4.12}$$

where

$$\gamma = \sqrt{5(1-\alpha)(1309\alpha^2 - 2280\alpha + 1216)}. \tag{4.13}$$

Using (4.11)–(4.13) gives the values of the initial slope $A = y'(0)$ listed in Table 3. The formulas (4.11) and (4.12) suggest that the initial slope $A = y'(0)$ depends mainly on the parameter α , where $0 < \alpha < 1$. Table 3 exhibits the initial slopes $A = y'(0)$ for various values of α . Table 4 exhibits the values of $y(x)$ for $\alpha = 0.5$ for $x = 0.1$ to 1.0.

Table 4: [51, 57].

x	y^{kidder}	$y_{[2/2]}$	$y_{[3/3]}$
0.1	0.8816588283	0.8633060641	0.8979167028
0.2	0.7663076781	0.7301262261	0.7985228199
0.3	0.6565379995	0.6033054140	0.7041129703
0.4	0.5544024032	0.4848898717	0.6165037901
0.5	0.4613650295	0.3761603869	0.5370533796
0.6	0.3783109315	0.2777311628	0.4665625669
0.7	0.3055976546	0.1896843371	0.4062426033
0.8	0.2431325473	0.1117105165	0.3560801699
0.9	0.1904623681	0.04323673236	0.3179966614
1.0	0.1587689826	0.01646750847	0.2900255005

Example 4.3 (see [56]). Consider the following Thomas-Fermi (T-F) equation [6–13, 17, 31, 33, 34, 54] which arises in the mathematical modeling of various models in physics, astrophysics, solid state physics, nuclear charge in heavy atoms, and applied sciences:

$$y''(x) = \frac{y^{3/2}}{x^{1/2}}, \quad (4.14)$$

with boundary conditions

$$y(0) = 1, \quad \lim_{x \rightarrow \infty} y(x) = 0. \quad (4.15)$$

By applying the convex homotopy,

$$y_0 + py_1 + p^2y_2 + \dots = y_0(x) + p \iint_0^x \left(x^{-1/2} (y_0 + py_1 + p^2y_2 + \dots)^{3/2} \right) dx dx. \quad (4.16)$$

Now, we apply a slight modification in the conventional initial value and take $y_0(x) = 1$, instead of $y_0(x) = 1 + Bx$, where $B = y'(0)$. By comparing the coefficient of like powers of p , the following approximants are obtained

$$\begin{aligned}
p^{(0)} : y_0(x) &= 1, \\
p^{(1)} : y_1(x) &= Bx + \frac{4}{3}x^{3/2}, \\
p^{(2)} : y_2(x) &= \frac{2}{5}Bx^{5/2} + \frac{1}{3}x^3, \\
p^{(3)} : y_3(x) &= \frac{3}{70}B^2x^{7/2} + \frac{2}{15}Bx^4 + \frac{2}{27}x^{9/2} + \frac{1}{3}x^3, \\
p^{(4)} : y_4(x) &= \frac{3}{70}B^2x^{7/2} + \frac{2}{15}Bx^4 + \frac{2}{27}x^{9/2} - \frac{1}{252}B^3x^{9/2} + \frac{1}{175}B^2x^5 \\
&\quad + \frac{2}{27}x^{9/2} + \frac{3}{70}B^2x^{7/2}.
\end{aligned} \quad (4.17)$$

Table 5: Padé approximants and initial slopes $y'(0)$ [56]

Padé approximants	Initial slope $y'(0)$	Error (%)
[2/2]	-1.211413729	23.71
[4/4]	-1.550525919	2.36
[7/7]	-1.586021037	12.9×10^{-2}
[8/8]	-1.588076820	3.66×10^{-4}
[10/10]	-1.588076779	3.64×10^{-4}

The series solution is given as

$$\begin{aligned}
y(x) = & 1 + Bx + Bx + \frac{4}{3}x^{3/2} + \frac{2}{5}Bx^{5/2} + \frac{1}{3}x^3 + \frac{3}{70}B^2x^{7/2} + \frac{2}{15}Bx^4 + \frac{2}{27}x^{9/2} + \frac{3}{70}B^2x^{7/2} \\
& - \frac{1}{252}B^3x^{9/2} + \frac{1}{175}B^2x^5 + \frac{2}{27}x^{9/2} + \frac{3}{70}B^2x^{7/2} + \frac{1}{1056}B^4x^{11/2} + \frac{4}{1575}B^3x^6 \\
& + \frac{557}{100100}B^2x^{13/2} + \frac{4}{693}Bx^7 + \frac{101}{52650}x^{15/2} - \frac{3}{9152}B^5x^{13/2} - \frac{29}{24255}B^4x^7 \\
& - \frac{512}{351000}B^3x^{15/2} - \frac{46}{45045}B^2x^8 - \frac{113}{1178100}Bx^{17/2} + \frac{23}{473850}x^9 \dots, \\
& \vdots
\end{aligned} \tag{4.18}$$

Setting $x^{1/2} = t$, the series solution is obtained as

$$\begin{aligned}
y(t) = & 1 + Bt^2 + \frac{4}{3}t^3 + \frac{2}{5}Bt^5 + \frac{1}{3}t^6 + \frac{3}{70}B^2t^7 + \frac{2}{15}Bt^8 + \left(-\frac{1}{252}B^3 + \frac{2}{27}\right)t^9 + \frac{1}{175}B^2t^{10} \\
& + \left(\frac{1}{1056}B^4 + \frac{31}{1485}B\right)t^{11} + \left(\frac{4}{1575}B^3 + \frac{4}{405}\right)t^{12} + \left(-\frac{3}{9152}B^5 + \frac{557}{100100}B^2\right)t^{13} \\
& + \left(-\frac{29}{24255}B^4 + \frac{4}{693}B\right)t^{14} + \left(\frac{7}{499}B^6 - \frac{623}{351000}B^3 + \frac{101}{52650}\right)t^{15} \\
& + \left(\frac{68}{105105}B^4 - \frac{46}{45045}B^2\right)t^{16} + \left(-\frac{3}{43520}B^7 + \frac{153173}{116424000}B^4 - \frac{113}{1178100}B\right)t^{17} + \dots.
\end{aligned} \tag{4.19}$$

The diagonal Padé approximants can be applied [56] in order to study the mathematical behavior of the potential $y(x)$ and to determine the initial slope of the potential $y'(0)$.

Example 4.4 (see [42]). Consider the generalized variant of the Flierl-Petviashvili equation [37]

$$y'' + \frac{1}{x}y' - y^n - y^{n+1} = 0, \tag{4.20}$$

with boundary conditions

$$y(0) = \alpha, \quad y'(0) = 0, \quad y(\infty) = 0. \quad (4.21)$$

Using the transformation $u(x) = xy'(x)$, the generalized FP equation can be converted to the following first-order initial value problem:

$$u'(x) = x \left(\int_0^x \left(\frac{u(x)}{x} \right)^n + \left(\frac{u(x)}{x} \right)^{n+1} dx \right), \quad (4.22)$$

with initial conditions

$$u(0) = 0, \quad u(0) = 0. \quad (4.23)$$

By applying the convex homotopy, we have

$$\begin{aligned} & u_0 + pu_1 + p^2u_2 + \dots \\ & = p \int_0^s \left(x \left(\int_0^s \left(\frac{1}{x} (u_0 + pu_1 + p^2u_2 + \dots) \right)^n + \left(\frac{1}{x} (u_0 + pu_1 + p^2u_2 + \dots) \right)^{n+1} dx \right) \right) dx. \end{aligned} \quad (4.24)$$

The series solution after four iterations is given by

$$\begin{aligned} u(x) &= \frac{(\alpha^n + \alpha^{n+1})}{2} x^2 + \frac{(\alpha^n + \alpha^{n+1})(n\alpha^n + (n+1)\alpha^{n+1})}{16\alpha} x^4 \\ &+ \frac{(\alpha^n + \alpha^{n+1})(2n(3n-1)\alpha^{2n} + 2n(3n+1)\alpha^{2n+1} + (3n+1)(n+1)\alpha^{2n+2})}{384\alpha^2} x^6 \\ &+ \frac{(\alpha^n + \alpha^{n+1})(n(18n^2 - 29n + 12)\alpha^{3n} + n(54n^2 - 33n + 7)\alpha^{3n+1} + 2\mathfrak{A})}{18432\alpha^3} x^8 \\ &+ \dots, \end{aligned} \quad (4.25)$$

where \mathfrak{A} denote $(18n^2 + 7n + 1)(3n\alpha^{3n+2} + (n+1)\alpha^{3n+3})$, and the inverse transformation will yield

$$\begin{aligned} y(x) &= \alpha + \frac{(\alpha^n + \alpha^{n+1})}{4} x^2 + \frac{(\alpha^n + \alpha^{n+1})(n\alpha^n + (n+1)\alpha^{n+1})}{64\alpha} x^4 \\ &+ \frac{(\alpha^n + \alpha^{n+1})(2n(3n-1)\alpha^{2n} + 2n(3n+1)\alpha^{2n+1} + (3n+1)(n+1)\alpha^{2n+2})}{2304\alpha^2} x^6 \\ &+ \frac{(\alpha^n + \alpha^{n+1})(n(18n^2 - 29n + 12)\alpha^{3n} + n(54n^2 - 33n + 7)\alpha^{3n+1} + 2\mathfrak{A})}{147456\alpha^3} x^8 \\ &+ \dots, \end{aligned} \quad (4.26)$$

Table 6: Roots of the Padé approximants monopole [42] $\alpha, n = 1$.

Degree	Roots
[2/2]	-1.5
[4/4]	-2.50746
[6/6]	-2.390278
[8/8]	-2.392214

Table 7: Roots of the Padé approximants monopole [42] $\alpha, n = 3$.

Degree	Roots
[2/2]	-2.0
[4/4]	-2.0
[6/6]	-2.0
[8/8]	-2.0

Table 8: Roots of the Padé approximants monopole [42] α .

Degree	Roots
[2/2]	0.0
[4/4]	-.2197575908
[6/6]	-1.1918424398
[8/8]	-1.848997181

Table 9: Roots [42] of the Padé approximants [8/8] monopole for several values of n .

n	[8/8] roots	n	[8/8] roots
1	-2.392213866	7	-1.000708285
2	-2.0	8	-1.000601615
3	-1.848997181	9	-1.000523005
4	-1.286025892	10	-1.000462636
5	-1.001101141	11	-1.000262137
6	-1.000861533	$n \rightarrow \infty$	-1.0

where \mathfrak{A} denote $(18n^2 + 7n + 1)(3n\alpha^{3n+2} + (n+1)\alpha^{3n+3})$. Diagonal Padé approximants can be applied [42] to find the roots of the FP monopole α for $n \geq 1$.

Table 9 shows that the roots of the monopole α converge to -1 as n increases.

Example 4.5 (see [58, 59]). Consider the two-dimensional nonlinear inhomogeneous initial boundary value problem for the integro-differential equation related to the Blasius problem

$$y''(x) = \alpha - \frac{1}{2} \int_0^x y(t)y''(t)dt, \quad -\infty < x < 0, \quad (4.27)$$

with boundary conditions

$$\begin{aligned} y(0) &= 0, & y'(0) &= 1, \\ \lim_{x \rightarrow \infty} y'(x) &= 0, \end{aligned} \quad (4.28)$$

Table 10: Padé approximants and numerical value of α [53].

Padé approximant	α
[2/2]	0.5778502691
[3/3]	0.5163977793
[4/4]	0.5227030798

where the constant α is positive and defined by

$$y''(0) = \alpha, \quad \alpha > 0. \quad (4.29)$$

By applying the convex homotopy, we have

$$y_0 + py_1 + \dots = y_0(x) - p \iint_0^x \left(\int_0^x (y_0 + py_1 + \dots) \left(\frac{d^2 y_0}{dx^2} + p \frac{d^2 y_1}{dx^2} + \dots \right) dx dx \right). \quad (4.30)$$

Proceeding as before, the series solution is given as

$$\begin{aligned} y(x) = & x + \frac{1}{2}\alpha x^2 - \frac{1}{48}\alpha x^4 - \frac{1}{240}\alpha^2 x^5 + \frac{1}{960}\alpha x^6 + \frac{11}{20160}\alpha^2 x^7 + \left(\frac{11}{161280}\alpha^3 + \frac{1}{960}\alpha \right) x^8 \\ & - \frac{43}{967680}\alpha^2 x^9 + \left(\frac{1}{52960}\alpha - \frac{5}{387072}\alpha^3 \right) x^{10} + \left(\frac{587}{212889600}\alpha^2 - \frac{5}{4257792}\alpha^4 \right) x^{11} \\ & + \left(-\frac{1}{16220160}\alpha + \frac{1}{7257792}\alpha^3 \right) x^{12} + \dots, \end{aligned} \quad (4.31)$$

and consequently

$$\begin{aligned} y'(x) = & 1 + \alpha x - \frac{1}{12}\alpha x^3 - \frac{1}{48}\alpha^2 x^4 + \frac{1}{160}\alpha x^5 + \frac{11}{2880}\alpha^2 x^6 \left(\frac{11}{20160}\alpha^3 - \frac{1}{2688}\alpha \right) x^7 \\ & - \frac{43}{107520}\alpha^2 x^8 + 10 \left(\frac{1}{552960}\alpha - \frac{5}{387072}\alpha^3 \right) x^9 + 11 \left(\frac{587}{212889600}\alpha^2 - \frac{5}{4257792}\alpha^4 \right) x^{10} \\ & + 12 \left(-\frac{1}{16220160}\alpha + \frac{1}{725760}\alpha^3 \right) x^{11} + \dots. \end{aligned} \quad (4.32)$$

Now, we apply the diagonal Padé approximants to determine a numerical value for the constant α by using the given condition. Padé approximant of $y'(x)$ usually converges on the entire real axis [58, 59]. Moreover, $y'(x)$ is free of singularities on the real axis. Substituting the boundary conditions $y'(-\infty) = 0$ in each Padé approximant which vanishes if the coefficient of x with the highest power in the numerator vanishes. By solving the resulting polynomials of these coefficients, we obtain the values of α listed in Table 10 [58, 59].

5. Conclusion

In this paper, we applied a reliable combination of He's polynomials and the diagonal Padé approximants for obtaining approximate solutions of various singular and nonsingular boundary value problems of diversified physical nature. The proposed algorithm is employed without using linearization, discretization, transformation, or restrictive assumptions. The fact that the suggested technique solves nonlinear problems without using Adomian's polynomials is a clear advantage of this technique over the decomposition method.

Acknowledgment

The author is highly grateful to Brig (R) Qamar Zaman, the Vice Chancellor of HITEC University, Taxila Cantt, Pakistan for providing excellent research environment and facilities.

References

- [1] S. Abbasbandy, "A new application of He's variational iteration method for quadratic Riccati differential equation by using Adomian's polynomials," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 59–63, 2007.
- [2] S. Abbasbandy, "Numerical solution of non-linear Klein-Gordon equations by variational iteration method," *International Journal for Numerical Methods in Engineering*, vol. 70, no. 7, pp. 876–881, 2007.
- [3] M. A. Abdou and A. A. Soliman, "New applications of variational iteration method," *Physica D*, vol. 211, no. 1-2, pp. 1–8, 2005.
- [4] M. A. Abdou and A. A. Soliman, "Variational iteration method for solving Burger's and coupled Burger's equations," *Journal of Computational and Applied Mathematics*, vol. 181, no. 2, pp. 245–251, 2005.
- [5] T. A. Abassy, M. A. El-Tawil, and H. El Zoheiry, "Solving nonlinear partial differential equations using the modified variational iteration Padé technique," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 73–91, 2007.
- [6] G. Adomian, "Solution of the Thomas-Fermi equation," *Applied Mathematics Letters*, vol. 11, no. 3, pp. 131–133, 1998.
- [7] N. Anderson, A. M. Arthurs, and P. D. Robinson, "Complementary variational principles for a generalized diffusion equation," *Proceedings of the Royal Society A*, vol. 303, pp. 497–502, 1968.
- [8] B. Batiha, M. S. M. Noorani, and I. Hashim, "Variational iteration method for solving multispecies Lotka-Volterra equations," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 903–909, 2007.
- [9] J. Biazar and H. Ghazvini, "He's variational iteration method for fourth-order parabolic equations," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 1047–1054, 2007.
- [10] V. Bush and S. H. Caldwell, "Thomas-Fermi equation solution by the differential analyzer," *Physical Review*, vol. 38, no. 10, pp. 1898–1902, 1931.
- [11] B. L. Burrows and P. W. Core, "A variational-iterative approximate solution of the Thomas-Fermi equation," *Quarterly of Applied Mathematics*, vol. 42, no. 1, pp. 73–76, 1984.
- [12] C. Y. Chan and Y. C. Hon, "A constructive solution for a generalized Thomas-Fermi theory of ionized atoms," *Quarterly of Applied Mathematics*, vol. 45, no. 3, pp. 591–599, 1987.
- [13] A. Cedillo, "A perturbative approach to the Thomas-Fermi equation in terms of the density," *Journal of Mathematical Physics*, vol. 34, no. 7, pp. 2713–2717, 1993.
- [14] A. Golbabai and M. Javidi, "A variational iteration method for solving parabolic partial differential equations," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 987–992, 2007.
- [15] A. Ghorbani and J. Saberi-Nadjafi, "He's homotopy perturbation method for calculating Adomian polynomials," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 8, no. 2, pp. 229–232, 2007.
- [16] A. Ghorbani, "Beyond Adomian polynomials: He polynomials," *Chaos, Solitons and Fractals*, vol. 39, no. 3, pp. 1486–1492, 2009.

- [17] J.-H. He, "Variational approach to the Thomas-Fermi equation," *Applied Mathematics and Computation*, vol. 143, no. 2-3, pp. 533–535, 2003.
- [18] J.-H. He, "An elementary introduction to recently developed asymptotic methods and nanomechanics in textile engineering," *International Journal of Modern Physics B*, vol. 22, no. 21, pp. 3487–3578, 2008.
- [19] J.-H. He, "Some asymptotic methods for strongly nonlinear equations," *International Journal of Modern Physics B*, vol. 20, no. 10, pp. 1141–1199, 2006.
- [20] J.-H. He, "Homotopy perturbation method for solving boundary value problems," *Physics Letters A*, vol. 350, no. 1-2, pp. 87–88, 2006.
- [21] J.-H. He, "Comparison of homotopy perturbation method and homotopy analysis method," *Applied Mathematics and Computation*, vol. 156, no. 2, pp. 527–539, 2004.
- [22] J.-H. He, "Homotopy perturbation method for bifurcation of nonlinear problems," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 6, no. 2, pp. 207–208, 2005.
- [23] J.-H. He, "The homotopy perturbation method nonlinear oscillators with discontinuities," *Applied Mathematics and Computation*, vol. 151, no. 1, pp. 287–292, 2004.
- [24] J.-H. He, "A coupling method of a homotopy technique and a perturbation technique for non-linear problems," *International Journal of Non-Linear Mechanics*, vol. 35, no. 1, pp. 37–43, 2000.
- [25] J.-H. He, "Variational iteration method—a kind of non-linear analytical technique: some examples," *International Journal of Non-Linear Mechanics*, vol. 34, no. 4, pp. 699–708, 1999.
- [26] J.-H. He, "Variational iteration method for autonomous ordinary differential systems," *Applied Mathematics and Computation*, vol. 114, no. 2-3, pp. 115–123, 2000.
- [27] J.-H. He and X.-H. Wu, "Construction of solitary solution and compacton-like solution by variational iteration method," *Chaos, Solitons and Fractals*, vol. 29, no. 1, pp. 108–113, 2006.
- [28] J.-H. He, "Variational iteration method—some recent results and new interpretations," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 3–17, 2007.
- [29] J.-H. He and X.-H. Wu, "Variational iteration method: new development and applications," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 881–894, 2007.
- [30] J.-H. He, "The variational iteration method for eighth-order initial-boundary value problems," *Physica Scripta*, vol. 76, no. 6, pp. 680–682, 2007.
- [31] Y. C. Hon, "A decomposition method for the Thomas-Fermi equation," *Southeast Asian Bulletin of Mathematics*, vol. 20, no. 3, pp. 55–58, 1996.
- [32] M. Inokuti, H. Sekine, and T. Mura, "General use of the Lagrange multiplier in nonlinear mathematical physics," in *Variational Method in the Mechanics of Solids*, S. Nemat-Naseer, Ed., pp. 156–162, Pergamon Press, New York, NY, USA, 1978.
- [33] S. Kobayashi, T. Matsukuma, S. Nagai, and K. Umeda, "Some coefficients of the TFD function," *Journal of the Physical Society of Japan*, vol. 10, pp. 759–765, 1955.
- [34] B. J. Laurenzi, "An analytic solution to the Thomas-Fermi equation," *Journal of Mathematical Physics*, vol. 31, no. 10, pp. 2535–2537, 1990.
- [35] J. Lu, "Variational iteration method for solving two-point boundary value problems," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 92–95, 2007.
- [36] W. X. Ma and D. T. Zhou, "Explicit exact solution of a generalized KdV equation," *Acta Mathematica Scientia*, vol. 17, pp. 168–174, 1997.
- [37] W.-X. Ma and Y. You, "Solving the Korteweg-de Vries equation by its bilinear form: Wronskian solutions," *Transactions of the American Mathematical Society*, vol. 357, no. 5, pp. 1753–1778, 2005.
- [38] W. X. Ma and Y. You, "Rational solutions of the Toda lattice equation in Casoratian form," *Chaos, Solitons and Fractals*, vol. 22, no. 2, pp. 395–406, 2004.
- [39] W.-X. Ma, H. Wu, and J. He, "Partial differential equations possessing Frobenius integrable decompositions," *Physics Letters A*, vol. 364, no. 1, pp. 29–32, 2007.
- [40] S. Momani and S. Abuasad, "Application of He's variational iteration method to Helmholtz equation," *Chaos, Solitons and Fractals*, vol. 27, no. 5, pp. 1119–1123, 2006.
- [41] S. T. Mohyud-Din, M. A. Noor, and K. I. Noor, "Travelling wave solutions of seventh-order generalized KdV equations using He's polynomials," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 10, no. 2, pp. 227–233, 2009.
- [42] S. T. Mohyud-Din and M. A. Noor, "Homotopy perturbation method and Padé approximants for solving Flierl-Petviashvili equation," *Applications and Applied Mathematics*, vol. 3, no. 2, pp. 224–234, 2008.
- [43] M. A. Noor and S. T. Mohyud-Din, "Homotopy perturbation method for solving nonlinear higher-order boundary value problems," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 9, no. 4, pp. 395–408, 2008.

- [44] M. A. Noor and S. T. Mohyud-Din, "Variational iteration method for solving higher-order nonlinear boundary value problems using He's polynomials," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 9, no. 2, pp. 141–157, 2008.
- [45] M. A. Noor and S. T. Mohyud-Din, "Variational iteration technique for solving higher order boundary value problems," *Applied Mathematics and Computation*, vol. 189, no. 2, pp. 1929–1942, 2007.
- [46] M. A. Noor and S. T. Mohyud-Din, "An efficient method for fourth-order boundary value problems," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 1101–1111, 2007.
- [47] M. A. Noor and S. T. Mohyud-Din, "Modified variational iteration method for heat and wave-like equations," *Acta Applicandae Mathematicae*, vol. 104, no. 3, pp. 257–269, 2008.
- [48] M. A. Noor and S. T. Mohyud-Din, "Variational homotopy perturbation method for solving higher dimensional initial boundary value problems," *Mathematical Problems in Engineering*, vol. 2008, Article ID 696734, 11 pages, 2008.
- [49] M. A. Noor, K. I. Noor, and S. T. Mohyud-Din, "Modified variational iteration technique for solving singular fourth-order parabolic partial differential equations," *Nonlinear Analysis Series A: Theory, Methods & Applications*, vol. 71, pp. e630–e640, 2009.
- [50] M. A. Noor and S. T. Mohyud-Din, "Variational iteration method for unsteady flow of gas through a porous medium using He's polynomials and Pade approximants," *Computers and Mathematics with Applications*, vol. 58, pp. 2182–2189, 2009.
- [51] M. A. Noor and S. T. Mohyud-Din, "Solution of singular and nonsingular initial and boundary value problems by modified variational iteration method," *Mathematical Problems in Engineering*, vol. 2008, Article ID 917407, 23 pages, 2008.
- [52] N. H. Sweilam, "Harmonic wave generation in non linear thermoelasticity by variational iteration method and Adomian's method," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 64–72, 2007.
- [53] N. H. Sweilam, "Fourth order integro-differential equations using variational iteration method," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 1086–1091, 2007.
- [54] R. K. Sabirov, "Solution of the Thomas-Fermi-Dirac of the statistical model of an atom at small distances from the nucleus," *Optical Spectra*, vol. 75, no. 1, pp. 1–2, 1993.
- [55] M. Tatari and M. Dehghan, "On the convergence of He's variational iteration method," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 121–128, 2007.
- [56] A.-M. Wazwaz, "The modified decomposition method and Padé approximants for solving the Thomas-Fermi equation," *Applied Mathematics and Computation*, vol. 105, no. 1, pp. 11–19, 1999.
- [57] A.-M. Wazwaz, "The modified decomposition method applied to unsteady flow of gas through a porous medium," *Applied Mathematics and Computation*, vol. 118, no. 2-3, pp. 123–132, 2001.
- [58] A.-M. Wazwaz, "A reliable algorithm for solving boundary value problems for higher-order integro-differentiable equations," *Applied Mathematics and Computation*, vol. 118, no. 2-3, pp. 327–342, 2001.
- [59] A.-M. Wazwaz, "A study on a boundary-layer equation arising in an incompressible fluid," *Applied Mathematics and Computation*, vol. 87, no. 2-3, pp. 199–204, 1997.
- [60] L. Xu, "He's homotopy perturbation method for a boundary layer equation in unbounded domain," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 1067–1070, 2007.
- [61] L. Xu, "Variational iteration method for solving integral equations," *Computers & Mathematics with Applications*, vol. 54, no. 7-8, pp. 1071–1078, 2007.
- [62] X.-W. Zhou, Y.-X. Wen, and J.-H. He, "Exp-function method to solve the nonlinear dispersive $K(m, n)$ equations," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 9, no. 3, pp. 301–306, 2008.
- [63] S. T. Mohyud-Din and M. A. Noor, "Homotopy perturbation method for solving fourth-order boundary value problems," *Mathematical Problems in Engineering*, vol. 2007, Article ID 98602, 15 pages, 2007.
- [64] S. T. Mohyud-Din and M. A. Noor, "Homotopy perturbation method for solving partial differential equations," *Zeitschrift für Naturforschung A*, vol. 64, pp. 157–170, 2009.
- [65] S. T. Mohyud-Din, M. A. Noor, and K. I. Noor, "Parameter-expansion techniques for strongly nonlinear oscillators," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 10, no. 5, pp. 581–583, 2009.
- [66] S. T. Mohyud-Din, M. A. Noor, and K. I. Noor, "Traveling wave solutions of seventh-order generalized kdv equations by variational iteration method using adomian's polynomials," *International Journal of Modern Physics B*, vol. 23, no. 15, pp. 3265–3277, 2009.

- [67] S. T. Mohyud-Din, M. A. Noor, and K. I. Noor, "Some relatively new techniques for nonlinear problems," *Mathematical Problems in Engineering*, vol. 2009, Article ID 234849, 25 pages, 2009.
- [68] S. T. Mohyud-Din, M. A. Noor, and K. I. Noor, "Variational iteration method for Flierl-Petviashvili equations using He's polynomials and Padé approximants," *Journal of Applied Sciences*, vol. 6, no. 8, pp. 1139–1146, 2009.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

