

Research Article

Oregano Essential Oil Induces SOD1 and GSH Expression through Nrf2 Activation and Alleviates Hydrogen Peroxide-Induced Oxidative Damage in IPEC-J2 Cells

Yi Zou,¹ Jun Wang,¹ Jian Peng,^{1,2} and Hongkui Wei¹

¹Department of Animal Nutrition and Feed Science, College of Animal Science and Technology, Huazhong Agricultural University, Wuhan 430070, China

²The Cooperative Innovation Center for Sustainable Pig Production, Wuhan 430070, China

Correspondence should be addressed to Hongkui Wei; weihongkui@mail.hzau.edu.cn

Received 20 July 2016; Revised 29 September 2016; Accepted 14 November 2016

Academic Editor: Oluseyi O. Jimoh

Copyright © 2016 Yi Zou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oregano essential oil (OEO) has long been used to improve the health of animals, particularly their intestinal health. The health benefits of OEO are generally attributed to antioxidative actions, but the mechanisms remain unclear. Here, we investigate the antioxidative effects of OEO and their underlying molecular mechanisms in porcine small intestinal epithelial (IPEC-J2) cells. We found that OEO treatment prior to hydrogen peroxide (H₂O₂) exposure increased cell viability and prevented lactate dehydrogenase (LDH) release into the medium. H₂O₂-induced reactive oxygen species (ROS) and malondialdehyde (MDA) were remarkably suppressed by OEO. OEO dose-dependently increased mRNA and protein levels of the nuclear factor-erythroid 2-related factor-2 (Nrf2) target genes Cu/Zn-superoxide dismutase (SOD1) and g-glutamylcysteine ligase (GCLC, GLCM), as well as intracellular concentrations of SOD1 and glutathione. OEO also increased intranuclear expression of Nrf2 and the activity of an antioxidant response element reporter plasmid in IPEC-J2 cells. The OEO-induced expression of Nrf2-regulated genes and increased SOD1 and glutathione concentrations in IPEC-J2 cells were reduced by Nrf2 small interfering (si) RNAs, counteracting the protective effects of OEO against oxidative stress in IPEC-J2 cells. Our results suggest that OEO protects against H₂O₂-induced IPEC-J2 cell damage by inducing Nrf2 and related antioxidant enzymes.

1. Introduction

Oxidative stress is widely recognized as a state of imbalance between prooxidants and antioxidants. Excessive production of reactive oxygen species (ROS) and/or a deficiency of antioxidants result in endogenous oxidative stress [1]. The pathogenesis of various gastrointestinal diseases, such as peptic ulcers, gastrointestinal cancers, and inflammatory bowel disease, involves oxidative stress [2]. Oregano (*Origanum vulgare* L.) is an aromatic plant widely distributed throughout the Mediterranean area and Asia [3]. Oregano essential oil (OEO), a volatile oil, is a concentrate of natural plant products that contain the volatile aromatic compounds. This mixture of volatile aromatic compounds has several biological actions,

including antimicrobial, anti-inflammatory, and antioxidative activities [4]. In recent studies, OEO has been demonstrated to possess antioxidant activity and prevent oxidative induced intestinal damage in animals [5–7]. These studies also suggest that OEO suppresses oxidative stress by inducing antioxidant enzymes, in addition to scavenging radicals. These studies also suggested that OEO suppress oxidative stress by inducing antioxidant enzymes, in addition to having radical-scavenging effects. However, the ability of OEO to induce antioxidant enzymes and the potential mechanisms in intestinal have not been reported.

Nuclear factor-erythroid 2-related factor-2 (Nrf2) is a transcription factor that plays a central role in cellular defenses against oxidative and electrophilic insults through

the timely induction of antioxidant enzymes and related proteins [8]. Under normal conditions, Nrf2 is bound to Kelch-like ECH-associated protein 1 (Keap1) and is ubiquitinated and targeted for degradation [9]. Shear stress, dietary antioxidants, and other physiological stimuli that disrupt Keap1-Nrf2 interactions allow nuclear accumulation of Nrf2, resulting in the transcription of antioxidant enzymes such as γ -glutamyl cysteine ligase (GCL) and Cu/Zn-superoxide dismutase (SOD1) [10–12]. GCL is a heterodimer consisting of catalytic (GCLC) and modifier (GCLM) subunits, both of which are products of Nrf2 target genes. GCL has been extensively investigated for its ability to regulate the synthesis of glutathione (GSH), the most abundant natural cellular antioxidant, which plays an essential role in maintaining the cellular redox state [13]. SOD plays an important role in the defense mechanisms of biological cells exposed to oxygen [14], catalyzing the dismutation of superoxide anion radical (O_2^-) into an oxygen molecule and a hydrogen peroxide (H_2O_2) [15]. This reaction is recognized as an antioxidant system that protects cells from superoxide toxicity. Cu/Zn-SOD (SOD1) exists in the cytoplasm, lysosomes, and nuclear compartments of mammalian cells [16].

OEO has been shown to have antioxidant properties in *in vivo* studies, but its antioxidant role is not well understood. The objectives of this study were to investigate the oxidative damaging effects of H_2O_2 on growth of porcine small intestinal epithelial (IPEC-J2) cells *in vitro*, determine whether OEO has an antioxidative effect (restoring the cell redox state during oxidative stress) on IPEC-J2 cells, and, if OEO does have an antioxidative effect, to elucidate the underlying molecular mechanisms.

2. Materials and Methods

2.1. Chemicals and Regents. Dulbecco's Modified Eagle Medium (DMEM) and fetal bovine serum (FBS) were purchased from GIBCO BRL (Grand Island, NY, USA). OEO was obtained from Meritech Bioengineering Co. Ltd. (Guangzhou, China), and its composition is shown in Table 1. Typical chromatogram of oregano essential oil is shown in Supplementary Material available online at <http://dx.doi.org/10.1155/2016/5987183>. H_2O_2 solution, 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT), 2-(4-amidinophenyl)-6-indolecarbamidine dihydrochloride (DAPI), and 2',7'-dichlorofluorescein-diacetate (DCFH₂-DA) were obtained from Sigma-Aldrich (St. Louis, MO, USA). Antibodies against Nrf2, Keap1, and β -actin were obtained from Cell Signaling Technology (Beverly, MA, USA). Antibodies against GCLM, γ -GCLC, and SOD1 were obtained from Affinity Biosciences (Columbus, OH, USA). The antibody against proliferating cell nuclear antigen (PCNA) was obtained from BD Transduction Laboratories (San Diego, CA, USA). The HRP-conjugated secondary antibodies, anti-rabbit IgG and anti-mouse IgG, were purchased from Jackson ImmunoResearch (West Grove, PA, USA). IPEC-J2 cells were obtained from the Cell Bank of Type Culture Collection of the Chinese Academy of Sciences Institution (Chinese Academy of Sciences, China) and cultured at 37°C in a 5% CO₂ atmosphere

TABLE 1: Chemical composition of oregano essential oil.

Components ^a	Composition%
α -Thujene/ α -pinene	0.56
Camphene	0.08
β -Pinene	0.09
Sabinene	0.03
Myrcene	0.91
α -Phellandrene	0.09
α -Terpinene	0.50
Limonene	0.15
1,8-Cineole+ β -phellandrene	0.07
β -Ocimene	0.07
γ -Terpinene	4.54
3-Octanone	0.07
ρ -Cymene	3.11
Terpinolene	0.05
3-Octanol	0.11
1-Octen-3-ol	0.22
Dimethyl styrene	0.10
Trans-Sabinene hydrate	0.14
Linalool	0.32
Cis-Sabinene hydrate	0.03
1-Terpilool	0.05
Terpinen-4-ol	0.22
Carvacrol methyl ether	0.33
β -Caryophyllene	1.43
Dihydrocarvone	0.09
α -Humulene	0.08
α -Terpineol	0.21
Bomeol	0.33
β -Bisabolene	0.71
Caryophyllene oxide	0.16
Thymol	1.90
Carvacrol	79.92
Total	99.99

^aThe dates were provided by Meritech Bioengineering Co. Ltd.

in DMEM containing 10% FBS. OEO was dissolved in the serum-containing medium to achieve the final desired concentration.

2.2. MTT Assay. Cell viability was monitored using an MTT colorimetric assay. IPEC-J2 cells (2×10^4 per well in a 96-well plate) were incubated with or without OEO and H_2O_2 . Cells were washed with phosphate-buffered saline (PBS) and then incubated with 10 μ L of 5 mg/mL MTT in PBS for 4 h. The supernatant was removed and the culture resuspended with 150 μ L of isopropanol to dissolve MTT formazan. The absorbance was measured at 490 nm using a spectrophotometer (Biomate 5, Thermo Electron Corporation, Rochester, NY, USA). The effect of OEO and H_2O_2 on cell viability was assessed as the percentage of viable cells compared with vehicle-treated control cells, which were arbitrarily assigned a viability of 100%.

2.3. Lactate Dehydrogenase (LDH) Release Assay. The activity of LDH released into the culture medium through damaged membranes was measured spectrophotometrically using an LDH cytotoxicity assay kit (Nanjing Jiancheng Institute of Bioengineering, Nanjing, China) according to the manufacturer's protocol. The method is based on the LDH-catalyzed reduction of pyruvate to lactate by an equimolar amount of NADH. NADH formation is detected by the release of a chromogen that emits at 490 nm. IPEC-J2 cells (2×10^4 per well in a 96-well plate) were pretreated with different concentrations of OEO and then exposed to H_2O_2 . The absorbance was measured with an ELISA microplate reader, and any decrease in absorbance was monitored.

2.4. Measurement of ROS Generation. The intracellular accumulation of ROS was detected by fluorescence microscopy using DCFH₂-DA. IPEC-J2 cells (1×10^5 cells/well in a 6-well plate) were cultured in DMEM supplemented with 10% FBS; culture medium was renewed when cells reached 80% confluence. After OEO pretreatment (90% confluence) and H_2O_2 incubation, cell culture media were removed, and cells were washed with PBS and incubated with $10 \mu M$ DCFH₂-DA in fresh culture medium at $37^\circ C$ for 30 min. The acetate groups on DCFH₂-DA were removed by an intracellular esterase, trapping the probe inside IPEC-J2 cells. Intracellular ROS, as indicated by DCF fluorescence, was measured with a fluorescence microscope (ECLIOSE Ti, Nikon, Japan). The fluorescence intensity under each condition was quantified on a FACSCalibur flow cytometer (BD Biosciences, San Diego, CA, USA) using excitation and emission filters of 488 and 530 nm, respectively. Levels of ROS were measured in the supernatant of IPEC-J2 cells by chemiluminescence assay using luminol (5-amino-2,3-dihydro-1,4-phthalazinedione, Sigma, St. Louis, MO, USA) as described in our previous study [17]. The results were expressed as relative light units (RLU).

2.5. Assessment of Lipid Peroxidation. IPEC-J2 cells (1×10^5 per well in a 6-well plate) were incubated with or without OEO and H_2O_2 . The content of malondialdehyde (MDA) in IPEC-J2 cells was assayed using colorimetric methods with a spectrophotometer (Biomate 5). The assays were conducted with kits purchased from Nanjing Jiancheng Institute of Bioengineering according to the manufacturer's instructions. IPEC-J2 cells were washed twice with cold PBS and harvested in 5% metaphosphoric acid using a cell scraper. Harvested cells were lysed by sonication on ice, and the supernatant of the centrifugate ($1300 \times g$ at $4^\circ C$ for 15 min) was collected. Samples were vortexed with 8.1% sodium dodecyl sulfate, 20% acetic acid, and 0.8% 2-thiobarbituric acid and incubated for 1 h at $95^\circ C$ before butanol-pyridine 15:1 (v/v) was added. The mixture was shaken for 10 min and then centrifuged ($1300 \times g$ at $4^\circ C$). The butanol-pyridine layer was measured fluorometrically at 552 nm.

2.6. Quantitative PCR. IPEC-J2 cells (5×10^4 per well in a 24-well plate) were incubated with or without OEO. Total RNA was extracted from samples of IPEC-J2 cells using Trizol reagent (Invitrogen, Carlsbad, CA, USA) according to

the manufacturer's instructions. Total RNA was determined using a NanoDrop® ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). For cDNA synthesis, $2.5 \mu g$ of RNA was reverse-transcribed using the Prime Script RT reagent kit (Takara, Otsu, Japan) according to the manufacturer's protocol. The primers used in this study, synthesized according to our previous protocols or designed with Primer 5.0 according to pig gene sequences, were as follows: Nrf2 5'-GAAAGCCCAGTCTTCATTGC-3' (sense), 5'-TTGGAACCGTGCTAGTCTCA-3' (antisense); SOD1 5'-ACCTGG-GCAATGTG-ACTG-3' (sense), 5'-TCCAGC-ATTTCCCGTCT-3' (antisense); CAT 5'-AACTGTCCC-TTCCGTGCTA-3' (sense), 5'-CCTGGGTGACATTATC-TTCG-3' (antisense); GCLC 5'-CATTGCGACACTG-GAGAC-3' (sense), 5'-CAAACCA-TCCTACCCTTTGG-3' (antisense); GCLM 5'-ATTGTGCAGAGAGCCTGGTT-3' (sense), and 5'-ACAATACAACGGTTCAGGTGAGT-3' (antisense). Real-time PCR was performed as described in our previous study [6]. The relative expression of genes in the treatment group was normalized to the values of the control group.

2.7. Western Blotting Analysis. IPEC-2 cells (1×10^6 cells per well in a 6-well plate) were collected and whole-cell lysates were prepared in RIPA buffer containing protease inhibitors or phosphatase inhibitors (Thermo Scientific, Rockford, IL, USA). Nuclear lysates were prepared using a nuclear/cytosol fractionation kit (BestBio, Shanghai, China), according to the manufacturer's protocol. They were then centrifuged at $12,000 \times g$ at $4^\circ C$ for 10 min, and the supernatants were collected for assay. Protein concentrations were determined using a standard BCA protein assay, and $30 \mu g$ protein was loaded per sample/lane and separated on SDS-PAGE. Protein samples were then electrophoretically transferred to nitrocellulose membranes, which were blocked in TBST (5% nonfat milk, 10 mM Tris, 150 mM NaCl, and 0.05% Tween-20) for 2 h. The blots were then incubated with one of the following primary antibodies at $4^\circ C$ overnight: anti-Nrf2 (1:1000), anti-Keap1 (1:1000), anti-GCLM (1:1000), anti-GCLC (1:1000), anti-SOD1 (1:1000), anti-PCNA (1:5000), and anti-actin (1:1000). After three washes with Tris-buffered saline containing 0.1% Tween 20, blots were incubated with an HRP-conjugated secondary antibody, anti-rabbit IgG (1:15000), or anti-mouse IgG (1:15000), for 2 h, and then were washed again. Chemiluminescence detection was performed using ECL reagent (Thermo Scientific) according to the manufacturer's instructions. Specific bands were detected, analyzed, and quantified by Image J Software (NIH, Bethesda, MD, USA).

2.8. Intracellular GSH and SOD1 Measurement. For these assays, IPEC-J2 cell number was optimized at 1×10^6 per well in 6-well plate. Cells were washed twice with cold PBS and harvested in 5% metaphosphoric acid using a cell scraper. Harvested cells were lysed by sonication on ice, and the supernatant of the centrifugate ($1300 \times g$ at $4^\circ C$ for 15 min) was collected. GSH levels were measured colorimetrically using a GSH assay kit (Nanjing Jiancheng Institute of Bioengineering) according to the manufacturer's protocol

and normalized to the protein concentration in the lysate. This assay is based on the spectrophotometric measurement of 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), the product of a reaction with GSH. The yellow derivative 5'-thio-2-nitrobenzoic acid (TNB) was measured by detecting absorbance at 412 nm using a microplate reader. SOD1 activity was determined by measuring inhibition of the reduction of the water-soluble tetrazolium salt, WST-8 (2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulphophenyl)-2H-tetrazolium, monosodium salt), which produces a water-soluble formazan dye upon reduction with $O_2^{\cdot -}$. Determinations were performed with a Superoxide Dismutase Assay Kit (Beyotime Biotechnology, Shanghai, China) specific to SOD1.

2.9. Nrf2 Immunofluorescence. IPEC-J2 cells (1×10^5 per well) were seeded on diameter glass coverslips coated with poly-L-lysine in a 6-well plate. Cells were incubated with or without 10 $\mu\text{g}/\text{mL}$ of OEO for 24 h. After being washed with cold PBS for three times, cells were fixed with 4% paraformaldehyde for 15 min, permeabilized with Triton X-100 (0.1%) for 15 min, and blocked with bovine serum albumin (1%) for 30 min. After being washed with PBS, cells were incubated overnight with Nrf2 antibody diluted 1:100 at 4°C, followed by incubation with Alexa Fluor 568-conjugated secondary antibody for 1 h. DAPI staining was performed to define nuclear regions. The fluorescence images were captured by laser scanning confocal microscopy (Eclipse TI, Nikon, Tokyo, Japan).

2.10. Luciferase Reporter Assay. The antioxidant response element (ARE) luciferase reporter plasmid (pGL6 (pARE-luc)) was obtained from Beyotime Biotechnology. IPEC-J2 cells were plated into 6-well plates (1×10^5 cells/well) and cultured for 24 h. Cells were cotransfected with 0.5 μg of luciferase expression plasmid and 1 μg of PRL-TK plasmid (Promega, Madison, USA) as a normalization reference for transfection efficiency using SuperFect transfection reagent (Qiagen, Valencia, CA, USA) for 24 h and then stimulated with OEO 12 h. Cells were harvested, and firefly and Renilla luciferase activities were determined using a Dual-Luciferase Reporter assay system (Progema). The ratio (reporter/control luciferase activity) obtained from control cell lysate was set at one.

2.11. SiRNA Transfection in IPEC-J2 Cells. The double-stranded pig Nrf2 siRNA 5'-GCCCAUUGAUCUCUCUGA-UTT-3' (sense) and 5'-AUCAGAGAGAAUGGGCTT-3' (antisense) were synthesized by GenePharma Co. Ltd. (Shanghai, China). IPEC-J2 cells were plated into 6-well plates (1×10^5 cells/well) and cultured for 24 h. Cells were transfected with 100 nM siRNA against Nrf2 or scrambled duplex using LipofectAMINE 2000 (Invitrogen). After 24-hour incubation, fresh medium was added and the cells were cultured for another 24 h before treatment with OEO.

2.12. Statistical Analyses. Statistical analysis was performed using Prism software (Prism 5.0; GraphPad Software, La Jolla, CA, USA). Data were analyzed by Student's *t*-test, Student-Newman-Keuls multiple-comparison test, or two-way ANOVA using SAS v 8.2 (SAS Inst. Inc., Cary, NC, USA).

Student's *t*-test was used for comparisons of two groups, Student-Newman-Keuls multiple-comparison test was used for comparisons among groups, and OEO concentration and Nrf2 siRNA were used as factors in the two-way ANOVA. Values are presented as means \pm standard error of the mean (SEM), and those at $P < 0.05$ were considered significant.

3. Results

3.1. Preventive Effect of OEO on Oxidative Stress-Induced Toxicity in IPEC-J2 Cells. The cytotoxic effects of OEO on IPEC-J2 cells were evaluated by MTT assay. The percentage of viable cells was compared with that of the control cells. Cells were treated with concentrations of OEO ranging from 1.25 to 80 $\mu\text{g}/\text{mL}$ for 3, 6, 12, or 24 h, to clarify the role of the time of exposure. Low doses of OEO (1.25–10 $\mu\text{g}/\text{mL}$) did not cause cytotoxic effects at any duration of exposure. OEO doses of 20–80 $\mu\text{g}/\text{mL}$ for 24 h significantly reduced cell viability compared with untreated control cells (Figure 1(a)). Based on these cell viability data, subsequent experiments were performed using concentrations of OEO ≤ 10 $\mu\text{g}/\text{mL}$. Cells were treated with concentrations of H_2O_2 ranging from 0.2 to 1.2 mM for 3, 6, 12, or 24 h to clarify the role of the time of exposure. The results of cell exposure to H_2O_2 are shown in Figure 1(b). The viability of cells treated with 0.8 mM H_2O_2 for 24 h was approximately 70% of that of the control cells. Therefore, treatment with 0.8 mM H_2O_2 24 h was selected for subsequent experiments.

Pretreatment of IPEC-J2 cells with OEO significantly improved cell viability in the presence of H_2O_2 in a dose-dependent manner (Figure 1(c)). To further explore the protective effect of OEO, the release of LDH, an indicator of cell injury, was also examined in H_2O_2 -exposed IPEC-J2 cells. The results showed that pretreatment with OEO attenuated the H_2O_2 -induced increase in LDH, affirming its protective role (Figure 1(d)).

3.2. OEO Inhibits Oxidative Stress-Induced ROS and MDA Production in IPEC-J2 Cells. H_2O_2 -induced intracellular ROS production in IPEC-J2 cells was monitored by fluorescence microscopic analysis using DCFH₂-DA as a fluorescent probe. IPEC-J2 cells exposed to H_2O_2 (0.8 mM) for 24 h showed a significant increase in fluorescence, which is proportionate to the amount of ROS generated (Figure 2(a)). This ROS induction was substantially inhibited by OEO pretreatment (2.5–10 $\mu\text{g}/\text{mL}$) (Figure 2(b)). Levels of ROS were also measured in the supernatant of IPEC-J2 cells by chemiluminescence assay using luminol as a probe. Pretreatment with OEO inhibited ROS levels in IPEC-J2 cell supernatants in the presence of H_2O_2 in a dose-dependent manner (Figure 2(c)). Thus, the protective effects of OEO in IPEC-J2 cells are further evidenced by its inhibition of excessive ROS production.

MDA is the end product of lipoperoxidation and is considered to be an excellent indicator of oxidative stress [18]. To investigate the effect of OEO on cell lipid peroxidation in cultured IPEC-J2 cells, levels of MDA were measured in IPEC-J2 cells using the TBARS assay. As shown in Figure 2(d), MDA levels in IPEC-J2 cells were markedly increased in




FIGURE 1: Oregano essential oil (OEO) inhibits hydrogen peroxide- (H_2O_2 -) induced cytotoxicity in IPEC-J2 cells. (a) IPEC-J2 cells were incubated with OEO (1.25–80 $\mu\text{g}/\text{mL}$) for 3–24 h. Cell viability was determined by MTT assay. (b) IPEC-J2 cells were incubated with H_2O_2 (0.2–1.2 mmol) for 3–24 h. Cell viability was determined by MTT assay. (c) IPEC-J2 cells were incubated with OEO (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h and then incubated with 0.8 mM H_2O_2 for 24 h. The protective effects of OEO were evaluated by MTT assay. (d) IPEC-J2 cells were incubated with OEO (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h and then incubated with 0.8 mM H_2O_2 for 24 h. The protective effects of OEO were evaluated by lactate dehydrogenase (LDH) release assay. Values represent means \pm SEM, $n = 3$. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$, compared with untreated control cells; means without a common letter differ, $P < 0.05$.

response to H_2O_2 compared with the control group, whereas pretreatment with OEO significantly inhibited lipid peroxidation in a dose-dependent manner.

3.3. OEO Increases Nrf2-Regulated Antioxidant Enzyme Expression. We predicted that the protective effects of OEO against oxidative stress occurred as a result of the induction of antioxidant genes. The expression of antioxidant genes

SOD1, catalase (CAT), GCLM, and GCLC under different conditions was analyzed. IPEC-J2 cells were treated with 10 $\mu\text{g}/\text{mL}$ of OEO for 3, 6, 12, or 24 h. OEO caused a maximal increase in SOD1, CAT, GCLM, and GCLC mRNA expression levels at 12 h (Figures 3(a), 3(b), 3(c), and 3(d)). Treatment with OEO at 2.5–10 $\mu\text{g}/\text{mL}$ for 12 h dose-dependently increased GCLM, GCLC, and SOD1 mRNA expression levels (Figures 3(e), 3(f), 3(g), and 3(h)), and pretreatment with




FIGURE 2: Oregano essential oil (OEO) inhibits hydrogen peroxide- (H_2O_2 -) induced reactive oxygen species (ROS) and malondialdehyde (MDA) production in IPEC-J2 cells. (a) Cells were pretreated with OEO (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h and then incubated with 0.8 mM H_2O_2 for 24 h. Intracellular ROS levels were indicated by DCF fluorescence and detected by fluorescence microscopy (200x magnification). (b) Cells were pretreated with OEO (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h and then incubated with 0.8 mM H_2O_2 for 24 h. The fluorescence intensity of DCF-stained cells was quantified by flow cytometer. (c) Cells were pretreated with OEO (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h and then incubated with 0.8 mM H_2O_2 for 24 h. The levels of ROS in culture supernatant were measured by chemiluminescence. Chemiluminometric counts were obtained at 0.05 sec intervals for 5 sec, and the results were expressed as areas under the curve of relative light units (RLU) for 5 sec. (d) Cells were pretreated with OEO (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h and then incubated with 0.8 mM H_2O_2 for 24 h. The intracellular MDA levels were measured. Values represent means \pm SEM, $n = 3$. Means without a common letter differ, $P < 0.05$.

10 $\mu\text{g}/\text{mL}$ of OEO upregulated CAT gene expression by 1.24-fold, compared with the nontreatment group. OEO also dose-dependently increased intracellular protein concentrations of GCLM, GCLC, and SOD1 (Figure 3(i)) and increased intracellular GSH (Figure 3(j)) and SOD1 (Figure 3(k)) protein concentrations in a dose-dependent manner.

3.4. Activation of Nrf2 by OEO. Nrf2 is a major transcription factor involved in cellular protection against oxidative stress through ARE-mediated induction of antioxidant enzymes [19]. As shown in Figure 4(a), treatment with OEO at 2.5–10 $\mu\text{g}/\text{mL}$ for 12 h dose-dependently increased Nrf2 mRNA expression levels. Western blots demonstrated that OEO incubation (2.5–10 $\mu\text{g}/\text{mL}$) for 24 h increased the ratio of Nrf2 to Keap1 in a semi-dose-dependent manner (Figure 4(b)). To investigate OEO-mediated nuclear translocation of Nrf2, the amount of cytosolic and nuclear Nrf2 protein was determined in the respective fractions of IPEC-J2 cells. The results showed increased nuclear Nrf2 and decreased cytosolic Nrf2 with 10 $\mu\text{g}/\text{mL}$ of OEO alone (Figure 4(c)). For further confirmation, nuclear import of Nrf2 in control and treated cells was monitored by immunofluorescence. As illustrated in Figure 4(d), Nrf2 was located in the cytoplasm of control cells, but the amount of nuclear Nrf2 increased after OEO treatment (10 $\mu\text{g}/\text{mL}$). To evaluate the involvement of Nrf2 in OEO-induced ARE activity, IPEC-J2 cells were transfected

with Nrf2 expression plasmids and OEO enhanced the activation of ARE in a dose-dependent manner (Figure 4(e)).

3.5. Antioxidant Gene Induction by OEO is Dependent on Nrf2. Nrf2 was downregulated by treatment with Nrf2 siRNA, as shown in Figures 5(a) and 5(b). Treatment with Nrf2 siRNA approximately reduced Nrf2 mRNA expression and protein expression by 53% and 47%, respectively. The increasing expressions of Nrf2, SOD1, GCLM, and GCLC mRNA caused by the OEO (2.5–10 $\mu\text{g}/\text{mL}$) were suppressed by Nrf2 siRNA (Figures 5(c), 5(d), 5(e), and 5(f)). As shown in Figure 5(g), Nrf2 siRNA reduced baseline Nrf2 protein as well as OEO-induced Nrf2 protein. This finding coincided with the greater reduction of the OEO-induced (10 $\mu\text{g}/\text{mL}$) Nrf2, SOD1, GCLM, and GCLC protein expression levels (Figure 5(h)). Similarly, Nrf2 siRNA also reduced the intracellular GSH and SOD1 induced by OEO (Figures 5(i) and 5(j)). It is worth noting, however, that treatment with OEO still increased GCLM, GCLC, and Nrf2 mRNA expression levels and intracellular GSH under conditions of Nrf2 suppression (Figures 5(c), 5(d), 5(e), and 5(i)).

3.6. Cytoprotection Against Oxidative Stress by OEO is Dependent on Nrf2. The role of Nrf2 in the antioxidative effect of OEO against H_2O_2 -induced cell toxicity was investigated in Nrf2 knockdown IPEC-J2 cells. OEO (10 $\mu\text{g}/\text{mL}$) suppressed




FIGURE 3: Effects of oregano essential oil (OEO) on g-glutamylcysteine ligase (GCLM, GCLC), and Cu/Zn-superoxide dismutase (SOD1) expression in IPEC-J2 cells. (a–d) IPEC-J2 cells were incubated with OEO 10 µg/mL for 3–24 h. (e–h) IPEC-J2 cells were incubated with OEO (2.5–10 µg/mL) for 12 h. Relative mRNA expression levels were measured quantitatively using real-time RT-PCR. The results were normalized to β-actin and expressed as fold increase over control. (i) IPEC-J2 cells were incubated with OEO (2.5–10 µg/mL) for 24 h. GCLC, GCLM, and SOD1 were estimated by western blot. (j) IPEC-J2 cells were incubated with OEO (2.5–10 µg/mL) for 24 h, and intracellular glutathione (GSH) content was determined. (k) IPEC-J2 cells were incubated with OEO (2.5–10 µg/mL) for 24 h, and intracellular SOD1 content was determined. Values represent means ± SEM, n = 3. *P < 0.05, **P < 0.01, and ***P < 0.001, compared with untreated control cells. Means without a common letter differ, P < 0.05.

H₂O₂-induced cell toxicity in cells treated with control siRNA, but the protective effects were reduced in Nrf2 knockdown cells (Figure 6(a)). Furthermore, the reduction in intracellular ROS induced by OEO was inhibited by knockdown of Nrf2 (Figure 6(b)), indicating that this antioxidative effect of OEO was mediated by Nrf2.

4. Discussion

The gastrointestinal tract is prone to ROS attack as it is accessed by entities from the outside environment, with resident immune cells and intestinal flora as well as dietary factors as potential sources of ROS [2]. Oxidative stress has been widely implicated in intestinal injury under both in vivo [20, 21] and in vitro conditions [22, 23]. Recent studies have shown

that OEO reduces oxidative stress and increases antioxidant enzymes in the intestines of animals [5, 7]. Furthermore, in a previous study in our laboratory, we observed that OEO increases antioxidant enzyme expression and reduces ROS levels in the intestine of rats in a model of oxidative stress [6]. However, the detailed molecular mechanisms by which OEO improves antioxidant status are unknown. To elucidate the mechanisms by which these antioxidant effects are exerted, here we assessed the effects of OEO on antioxidant enzymes and molecules in porcine IPEC-J2 intestinal epithelial cells. The OEO used in the present study contains volatile, natural, complex compounds characterized by a strong odor, formed by *Origanum vulgare* L as secondary metabolites. The main components of the OEO were thymol and carvacrol, while their biogenetic precursors, p-Cymene and γ-Terpinene, were




FIGURE 4: Effects of oregano essential oil (OEO) on nuclear factor-erythroid 2-related factor-2 (Nrf2) activation in IPEC-J2 cells. (a) IPEC-J2 cells were incubated with OEO (2.5–10 $\mu\text{g/mL}$) for 12 h. Relative Nrf2 mRNA expression levels were measured quantitatively using real-time RT-PCR. (b) IPEC-J2 cells were incubated with OEO 10 $\mu\text{g/mL}$ for 24 h and then Nrf2 and Kelch-like ECH-associated protein 1 (Keap1) were estimated in IPEC-J2 cells. (c) IPEC-J2 cells were incubated with OEO 10 $\mu\text{g/mL}$ for 24 h; then Nrf2 was estimated in the cytosolic and nuclear fractions. Western blot results show the effects of OEO on protein levels of Nrf2 in the cytosolic and nuclear fractions. (d) Localization of endogenous Nrf2. IPEC-J2 cells were incubated with OEO 10 $\mu\text{g/mL}$ for 24 h and then subjected to immunohistochemical staining with antibodies specific for Nrf2 followed by incubation with Alexa Fluor 568-conjugated secondary antibodies. For each condition, an image of the cell nucleus stained with DAPI specific for nuclear proteins is also presented. The immunohistochemical staining was observed using fluorescence microscopy (200x magnification). (e) IPEC-J2 cells were incubated with OEO 10 $\mu\text{g/mL}$ for 24 h; then levels of Nrf2 and Keap1 were estimated in IPEC-J2 cells. (e) IPEC-J2 cells were cotransfected with a reporter plasmid (pGL6) and a control plasmid (pRL-TK). After transfection, IPEC-J2 cells were incubated with OEO (2.5–10 $\mu\text{g/mL}$) for 12 h; then luciferase activity was measured. Values represent means \pm SEM, $n = 3$. * $P < 0.05$ and *** $P < 0.001$, compared with untreated control cells. Means without a common letter differ, $P < 0.05$.

the most abundant monoterpenes (Table 1). Importantly, the OEO concentrations (2.5–10 $\mu\text{g/mL}$) used in the present study were noncytotoxic and very low. Favorable pharmacological properties of OEO have been demonstrated in several studies at similarly low concentrations. For instance, treatment of oxidized low density lipoprotein-activated THP-1 macrophages with OEO (10–30 $\mu\text{g/mL}$) results in an overall reduction in proinflammatory cytokine release [24]. Likewise, antioxidative effects of OEO components thymol and carvacrol (8 to 30 $\mu\text{g/mL}$) have been demonstrated in Caco-2 cells [25]. To our knowledge, this study is the first to demonstrate that OEO increases antioxidant enzyme expression levels and prevents oxidative stress-induced cell death through activation of Nrf2.

H_2O_2 is often used in studies of redox-regulated processes because it is relatively stable compared with other ROS [26]. It has been well documented that exposure to H_2O_2

triggers the rapid generation of ROS in various cell types [27–29], including IPEC-J2 cells [23]. Accumulation of ROS and impairment of the antioxidant defense system by H_2O_2 cause oxidative damage in cells [30]. In the present study, IPEC-J2 cells treated with H_2O_2 experienced a dose- and time-dependent loss of cell viability. However, pretreatment of cells with OEO at an appropriate concentration (2.5, 5, or 10 $\mu\text{g/mL}$) significantly reduced this loss of cell viability, affirmed by attenuation in LDH release. Moreover, pretreatment of cells with OEO suppressed H_2O_2 -induced ROS and MDA production, in accordance with an inhibition of oxidative stress. This inhibition of H_2O_2 -induced ROS generation and cell lipid peroxidation by pretreatment with OEO may occur via a direct antioxidant mechanism through free radical-scavenging activity. Polyphenolic compounds like OEO exert a wide range of antioxidant effects, acting as ROS scavengers and free radical reaction terminators [31,




FIGURE 5: Continued.




FIGURE 5: Effects of oregano essential oil (OEO) on g-glutamylcysteine ligase (GCLM, GCLC) and Cu/Zn-superoxide dismutase (SOD1) expression in nuclear factor-erythroid 2-related factor-2 (Nrf2) knockdown IPEC-J2 cells. (a, b) IPEC-J2 cells were treated with Nrf2 siRNA or control siRNA and incubated for 24 h. (a) Nrf2 mRNA expression was quantitated and normalized as for Figure 3(a). (b) Whole-cell lysates were subjected to western blotting. (c–f) IPEC-J2 cells were transfected with siRNA against Nrf2 or control siRNA. After 24 h, the cells were incubated with OEO (2.5–10 µg/mL) for an additional 12 h. Nrf2, GCLC, GCLM, and SOD1 mRNA expression levels were measured quantitatively using real-time RT-PCR. (g–h) IPEC-J2 cells were transfected with siRNA against Nrf2 or control siRNA. After 24 h, the cells were incubated with OEO 10 µg/mL for an additional 24 h. Nrf2, Kelch-like ECH-associated protein 1 (Keap1), GCLC, GCLM, and SOD1 were estimated by western blot. (i) Effect of OEO on intracellular glutathione (GSH) content in Nrf2 knockdown IPEC-J2 cells. OEO (2.5–10 µg/mL) was applied as described for Figure 3(j), and intracellular GSH content was measured. (j) Effect of OEO on intracellular SOD1 content in Nrf2 knockdown IPEC-J2 cells. OEO (2.5–10 µg/mL) was applied as described for Figure 3(k), and intracellular SOD1 content was measured. Values represent means \pm SEM, $n = 3$. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$, compared with control cells treated with control siRNA; ### $P < 0.001$, compared with corresponding cells treated with control siRNA. Means without a common letter differ, $P < 0.05$.




FIGURE 6: Nuclear factor-erythroid 2-related factor-2 (Nrf2) siRNA diminishes the protective effects of IPEC-J2 cells. IPEC-J2 cells were treated with Nrf2 siRNA or control siRNA and incubated for 24 h. Cells were incubated with oregano essential oil (OEO, 10 µg/mL for 24 h) and then incubated with H₂O₂ (0.8 mM) for 24 h. (a) Cell viability was determined by MTT assay. (b) Intracellular reactive oxygen species (ROS) levels were measured by DCF fluorescence microscopy. Values represent means \pm SEM, $n = 3$, significant at * $P < 0.05$, ** $P < 0.01$.

32]. Pretreatment of cultured IPEC-J2 cells with OEO also significantly increased and restored intracellular GSH and SOD1 levels, possibly as a result of the antioxidative action of OEO against H₂O₂-induced oxidative damage.

To protect against the harmful effects of ROS, antioxidant enzymes such as CAT and SOD detoxify ROS to safe

molecules [33]. SOD averts oxidative stress by catalyzing superoxide anions to H₂O₂ and CAT further reduces redox damage by catalyzing the reduction of H₂O₂ [34]. Thus, increased expression of SOD and CAT may protect against oxidative stress. The tripeptide GSH (γ -L-Glu-L-Cys-Gly) is the most abundant intracellular nonprotein thiol compound

in mammalian cells [35]. Importantly, GSH functions as an antioxidant with a crucial role in scavenging toxic free radicals [36]. Thus, endogenous GSH is consumed during oxidative stress, resulting in a depletion of cellular GSH [37]. GSH contains an unusual peptide linkage between the γ -carboxyl of glutamate and the α -amino group of cysteine [38]. The formation of γ -glutamylcysteine is the first step in GSH synthesis. It is catalyzed by GCL, a heterodimeric protein composed of catalytic (GCLC) and modifier (GCLM) subunits, which are encoded by separate genes in mammals [39]. GCLC exhibits all of the catalytic activity, but its association with GCLM alters its kinetic properties to enhance GCL activity [40]. Glycine is subsequently added to γ -glutamylcysteine in a second reaction that is catalyzed by glutathione synthase to form GSH [38].

In recent years, several studies have reported that plant-derived compounds activate Nrf2-dependent ARE activity and elevate the levels of cellular CAT, SOD1, and GSH, which exhibit cytoprotective effects against oxidative stress in various cells [30, 41]. The action of Nrf2 plays a key role in the adaptive response to oxidative stress and regulates ARE-driven antioxidant gene expression (including those encoding SOD1, CAT, GCLC, and GCLM) [42]. Under conditions of homeostasis, Nrf2 is retained in the cytosol, bound to a cluster of proteins that includes its cytosolic inhibitor, Keap1 [43, 44]. When stimulated, it dissociates from Keap1 and moves to the nucleus to bind with ARE to regulate the transcription of antioxidant genes [45]. Here, when we pretreated IPEC-J2 cells with OEO, Nrf2 dissociated from the inhibitory protein Keap1 and translocated to the nucleus, as indicated by a dramatic increase in nuclear Nrf2. Accumulated nuclear Nrf2 has been demonstrated to bind to the ARE and lead to transcriptional activation of antioxidant genes [46]. Our findings further confirm these molecular cascades by reporting increased ARE activity in IPEC-J2 cells treated with OEO. Activation of Nrf2 has been shown to regulate important cellular antioxidant genes, including SOD1, CAT, GCLC, and GCLM, and protect the cells from oxidative damage. These data are consistent with previous observations that plant extracts (including those from Willow bark, Ginkgo biloba, Nelumbo nucifera leaves, Piperaceae leaves, and Grape pomace) increase cellular antioxidant gene expression by increasing the levels of nuclear Nrf2 [30, 42, 47–49]. Knockdown of Nrf2 with siRNA inhibited not only OEO-induced expression of these antioxidant enzymes, but also its protective effect against oxidative stress-induced cellular damage in IPEC-J2 cells. In this study, while the increase in GCLM, GCLC, and Nrf2 mRNA expression levels and intracellular GSH caused by OEO was significantly suppressed by Nrf2 siRNA, treatment with OEO still increased GCLM, GCLC, and Nrf2 mRNA expression levels and intracellular GSH under conditions of Nrf2 suppression. However, the efficiency with which siRNAs inhibited Nrf2 was relatively low, with only 53% downregulation of Nrf2 gene expression and 47% downregulation of protein abundance. Alternatively, it has been shown that the forkhead box O (FOXO) family of transcription factors, tumor suppressor p53, and activator protein 1 (AP-1) also have major roles in preventing oxidative stress by inducing antioxidant gene expression [50–52]. Thus,

Nrf2 may not be the only factor modulating oxidative stress, and OEO may modulate oxidative stress by one or more different pathways.

5. Conclusions

In conclusion, this study provides evidence for the involvement of OEO in protecting against oxidative stress in IPEC-J2 cells and elucidates several underlying mechanisms. Treatment of IPEC-J2 cells with OEO enhanced SOD1 and GSH expression through activation of the Nrf2/ARE pathway. This mechanism may be pivotal to its antioxidative action against H_2O_2 -induced cell damage. ROS are essential factors in the pathogenesis of many gastrointestinal diseases. Our data support the need for further research into OEO activation of Nrf2 and the endogenous antioxidant response as a potential therapeutic approach for these diseases. OEO may indeed have broad applicability to therapies for oxidative stress-related intestinal diseases.

Competing Interests

The authors declare that they have no competing interests.

Acknowledgments

This study was supported financially by the International S&T Cooperation Projects of China (Grant no. 2013DFG32510) and the Key Technology Research and Development Program of Hubei Province (nos. 2014ABB014 and 2014ABC012). The authors are grateful to the colleagues for their helpful discussions.

References

- [1] C. Nathan and A. Cunningham-Bussel, “Beyond oxidative stress: an immunologist’s guide to reactive oxygen species,” *Nature Reviews Immunology*, vol. 13, no. 5, pp. 349–361, 2013.
- [2] A. Bhattacharyya, R. Chattopadhyay, S. Mitra, and S. E. Crowe, “Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases,” *Physiological Reviews*, vol. 94, no. 2, pp. 329–354, 2014.
- [3] M. Milos, J. Mastelic, and I. Jerkovic, “Chemical composition and antioxidant effect of glycosidically bound volatile compounds from oregano (*Origanum vulgare* L. ssp. *hirtum*),” *Food Chemistry*, vol. 71, no. 1, pp. 79–83, 2000.
- [4] F. Bakkali, S. Averbeck, D. Averbeck, and M. Idaomar, “Biological effects of essential oils—a review,” *Food and Chemical Toxicology*, vol. 46, no. 2, pp. 446–475, 2008.
- [5] T. Srihari, M. Sengottuvelan, and N. Nalini, “Dose-dependent effect of oregano (*Origanum vulgare* L.) on lipid peroxidation and antioxidant status in 1,2-dimethylhydrazine-induced rat colon carcinogenesis,” *Journal of Pharmacy and Pharmacology*, vol. 60, no. 6, pp. 787–794, 2008.
- [6] H.-K. Wei, G. Chen, R.-J. Wang, and J. Peng, “Oregano essential oil decreased susceptibility to oxidative stress-induced dysfunction of intestinal epithelial barrier in rats,” *Journal of Functional Foods*, vol. 18, pp. 1191–1199, 2015.
- [7] K. Mueller, N. M. Blum, H. Kluge, and A. S. Mueller, “Influence of broccoli extract and various essential oils on performance and expression of xenobiotic- and antioxidant enzymes in

- broiler chickens," *British Journal of Nutrition*, vol. 108, no. 4, pp. 588–602, 2012.
- [8] J. Kim, Y.-N. Cha, and Y.-J. Surh, "A protective role of nuclear factor-erythroid 2-related factor-2 (Nrf2) in inflammatory disorders," *Mutation Research—Fundamental & Molecular Mechanisms of Mutagenesis*, vol. 690, no. 1-2, pp. 12–23, 2010.
- [9] M. McMahon, N. Thomas, K. Itoh, M. Yamamoto, and J. D. Hayes, "Dimerization of substrate adaptors can facilitate Cullin-mediated ubiquitylation of proteins by a 'tethering' mechanism: a two-site interaction model for the Nrf2-Keap1 complex," *Journal of Biological Chemistry*, vol. 281, no. 34, pp. 24756–24768, 2006.
- [10] R. Masella, R. Di Benedetto, R. Vari, C. Filesi, and C. Giovannini, "Novel mechanisms of natural antioxidant compounds in biological systems: involvement of glutathione and glutathione-related enzymes," *Journal of Nutritional Biochemistry*, vol. 16, no. 10, pp. 577–586, 2005.
- [11] M. McMahon, K. Itoh, M. Yamamoto, and J. D. Hayes, "Keap1-dependent proteasomal degradation of transcription factor Nrf2 contributes to the negative regulation of antioxidant response element-driven gene expression," *The Journal of Biological Chemistry*, vol. 278, no. 24, pp. 21592–21600, 2003.
- [12] H. Dreger, K. Westphal, A. Weller et al., "Nrf2-dependent upregulation of antioxidative enzymes: a novel pathway for proteasome inhibitor-mediated cardioprotection," *Cardiovascular Research*, vol. 83, no. 2, pp. 354–361, 2009.
- [13] M. Valko, D. Leibfritz, J. Moncol, M. T. D. Cronin, M. Mazur, and J. Telser, "Free radicals and antioxidants in normal physiological functions and human disease," *International Journal of Biochemistry & Cell Biology*, vol. 39, no. 1, pp. 44–84, 2007.
- [14] J. M. McCord and I. Fridovich, "Superoxide dismutase. An enzymic function for erythrocyte protein (hemocuprein)," *The Journal of Biological Chemistry*, vol. 244, no. 22, pp. 6049–6055, 1969.
- [15] I. Fridovich, "Superoxide radical and superoxide dismutases," *Annual Review of Biochemistry*, vol. 64, pp. 97–112, 1995.
- [16] J. D. Crapo, T. Oury, C. Rabouille, J. W. Slot, and L.-Y. Chang, "Copper,zinc superoxide dismutase is primarily a cytosolic protein in human cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 89, no. 21, pp. 10405–10409, 1992.
- [17] Y. Zou, H. K. Wei, Q. Xiang, J. Wang, Y. Zhou, and J. Peng, "Protective effect of quercetin on pig intestinal integrity after transport stress is associated with regulation oxidative status and inflammation," *Journal of Veterinary Medical Science*, vol. 78, no. 9, pp. 1487–1494, 2016.
- [18] E. Ho, K. Karimi Galougahi, C.-C. Liu, R. Bhindi, and G. A. Figtree, "Biological markers of oxidative stress: applications to cardiovascular research and practice," *Redox Biology*, vol. 1, no. 1, pp. 483–491, 2013.
- [19] H.-Y. Cho, S. P. Reddy, A. DeBiase, M. Yamamoto, and S. R. Kleiberger, "Gene expression profiling of NRF2-mediated protection against oxidative injury," *Free Radical Biology & Medicine*, vol. 38, no. 3, pp. 325–343, 2005.
- [20] E. Aghdassi, J. Carrier, J. Cullen, M. Tischler, and J. P. Allard, "Effect of iron supplementation on oxidative stress and intestinal inflammation in rats with acute colitis," *Digestive Diseases & Sciences*, vol. 46, no. 5, pp. 1088–1094, 2001.
- [21] J. Piao, Y. Nakatsu, M. Ohno, K.-I. Taguchi, and T. Tszuzuki, "Mismatch repair deficient mice show susceptibility to oxidative stress-induced intestinal carcinogenesis," *International Journal of Biological Sciences*, vol. 10, no. 1, pp. 73–79, 2013.
- [22] M. K. Amani, A. T. Sané, G. Gouspillou et al., "Sa1432 CFTR knockout induces oxidative stress, inflammation and mitochondrial dysfunctions in Caco-2/15 intestinal epithelial cells," *Gastroenterology*, vol. 150, no. 4, pp. S313–S314, 2016.
- [23] X. Cai, X. Chen, X. Wang et al., "Pre-protective effect of lipoic acid on injury induced by H₂O₂ in IPEC-J2 cells," *Molecular & Cellular Biochemistry*, vol. 378, no. 1-2, pp. 73–81, 2013.
- [24] A. Ocaña-Fuentes, E. Arranz-Gutiérrez, F. J. Señorans, and G. Reglero, "Supercritical fluid extraction of oregano (*Origanum vulgare*) essential oils: anti-inflammatory properties based on cytokine response on THP-1 macrophages," *Food and Chemical Toxicology*, vol. 48, no. 6, pp. 1568–1575, 2010.
- [25] M. Llana-Ruiz-Cabello, D. Gutiérrez-Praena, M. Puerto, S. Pichardo, Á. Jos, and A. M. Cameán, "In vitro pro-oxidant/antioxidant role of carvacrol, thymol and their mixture in the intestinal Caco-2 cell line," *Toxicology In Vitro*, vol. 29, no. 4, pp. 647–656, 2015.
- [26] E. Paszti-Gere, E. Csibrik-Nemeth, K. Szeker, R. Csizinszky, C. Jakab, and P. Galfi, "Acute oxidative stress affects IL-8 and TNF- α expression in IPEC-J2 porcine epithelial cells," *Inflammation*, vol. 35, no. 3, pp. 994–1004, 2012.
- [27] C. Osera, M. Amadio, S. Falone et al., "Pre-exposure of neuroblastoma cell line to pulsed electromagnetic field prevents H₂O₂-induced ROS production by increasing MnSOD activity," *Bioelectromagnetics*, vol. 36, no. 3, pp. 219–232, 2015.
- [28] J.-L. Wu, H.-Y. Wang, Y.-L. Cheng, C. Du, and H. Qian, "Neuroprotective effects of torularhodin against H₂O₂-induced oxidative injury and apoptosis in PC12 cells," *Pharmazie*, vol. 70, no. 1, pp. 17–23, 2015.
- [29] Y. Hu, N. Zhang, Q. Fan et al., "Protective efficacy of carnolic acid against hydrogen peroxide induced oxidative injury in HepG2 cells through the SIRT1 pathway," *Canadian Journal of Physiology and Pharmacology*, vol. 93, no. 8, pp. 625–631, 2015.
- [30] J.-Y. Je and D.-B. Lee, "Nelumbo nucifera leaves protect hydrogen peroxide-induced hepatic damage via antioxidant enzymes and HO-1/Nrf2 activation," *Food & Function*, vol. 6, no. 6, pp. 1911–1918, 2015.
- [31] R. Amorati, M. C. Foti, and L. Valgimigli, "Antioxidant activity of essential oils," *Journal of Agricultural and Food Chemistry*, vol. 61, no. 46, pp. 10835–10847, 2013.
- [32] T. Kulisic, A. Radonic, V. Katalinic, and M. Milos, "Use of different methods for testing antioxidative activity of oregano essential oil," *Food Chemistry*, vol. 85, no. 4, pp. 633–640, 2004.
- [33] J. Nordberg and E. S. J. Arnér, "Reactive oxygen species, antioxidants, and the mammalian thioredoxin system," *Free Radical Biology and Medicine*, vol. 31, no. 11, pp. 1287–1312, 2001.
- [34] A. Kunwar and K. Priyadarsini, "Free radicals, oxidative stress and importance of antioxidants in human health," *Journal of Medical & Allied Sciences*, vol. 1, no. 2, pp. 53–60, 2011.
- [35] H. Sies, "Glutathione and its role in cellular functions," *Free Radical Biology and Medicine*, vol. 27, no. 9-10, pp. 916–921, 1999.
- [36] S. Bharath, M. Hsu, D. Kaur, S. Rajagopalan, and J. K. Andersen, "Glutathione, iron and Parkinson's disease," *Biochemical Pharmacology*, vol. 64, no. 5-6, pp. 1037–1048, 2002.
- [37] S. M. Kumar, K. Swaminathan, D. L. Clemens, and A. Dey, "GSH protects against oxidative stress and toxicity in VL-17A cells exposed to high glucose," *European Journal of Nutrition*, vol. 54, no. 2, pp. 223–234, 2014.
- [38] A. K. Sikalidis, K. M. Mazor, J.-I. Lee, H. B. Roman, L. L. Hirschberger, and M. H. Stipanuk, "Upregulation of capacity for glutathione synthesis in response to amino acid deprivation:

- regulation of glutamate-cysteine ligase subunits,” *Amino Acids*, vol. 46, no. 5, pp. 1285–1296, 2014.
- [39] S. C. Lu, “Glutathione synthesis,” *Biochimica et Biophysica Acta (BBA)—General Subjects*, vol. 1830, no. 5, pp. 3143–3153, 2013.
- [40] M. E. Anderson, “Glutathione: an overview of biosynthesis and modulation,” *Chemico-Biological Interactions*, vol. 111-112, pp. 1–14, 1998.
- [41] X. Jin, K. Wang, H. Liu, F. Hu, F. Zhao, and J. Liu, “Protection of bovine mammary epithelial cells from hydrogen peroxide-induced oxidative cell damage by resveratrol,” *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 2572175, 15 pages, 2016.
- [42] W. N. Wan Hasan, M.-K. Kwak, S. Makpol, W. Z. W. Ngah, and Y. A. M. Yusof, “Piper betle induces phase I & II genes through Nrf2/ARE signaling pathway in mouse embryonic fibroblasts derived from wild type and Nrf2 knockout cells,” *BMC Complementary and Alternative Medicine*, vol. 14, article no. 72, 2014.
- [43] Y.-C. Hseu, H.-W. Lo, M. Korivi, Y.-C. Tsai, M.-J. Tang, and H.-L. Yang, “Dermato-protective properties of ergothioneine through induction of Nrf2/ARE-mediated antioxidant genes in UVA-irradiated Human keratinocytes,” *Free Radical Biology and Medicine*, vol. 86, pp. 102–117, 2015.
- [44] S. Tanigawa, M. Fujii, and D.-X. Hou, “Action of Nrf2 and Keap1 in ARE-mediated NQO1 expression by quercetin,” *Free Radical Biology and Medicine*, vol. 42, no. 11, pp. 1690–1703, 2007.
- [45] Q. Zhong, M. Mishra, and R. A. Kowluru, “Transcription factor Nrf2-mediated antioxidant defense system in the development of diabetic retinopathy,” *Investigative Ophthalmology & Visual Science*, vol. 54, no. 6, pp. 3941–3948, 2013.
- [46] S. K. Niture, R. Khatri, and A. K. Jaiswal, “Regulation of Nrf2—an update,” *Free Radical Biology and Medicine*, vol. 66, pp. 36–44, 2014.
- [47] A. Ishikado, Y. Sono, M. Matsumoto et al., “Willow bark extract increases antioxidant enzymes and reduces oxidative stress through activation of Nrf2 in vascular endothelial cells and *Caenorhabditis elegans*,” *Free Radical Biology and Medicine*, vol. 65, pp. 1506–1515, 2013.
- [48] N. Goutzourelas, D. Stagos, A. Housmekeridou et al., “Grape pomace extract exerts antioxidant effects through an increase in GCS levels and GST activity in muscle and endothelial cells,” *International Journal of Molecular Medicine*, vol. 36, no. 2, pp. 433–441, 2015.
- [49] X.-P. Liu, C. E. P. Goldring, I. M. Copple et al., “Extract of Ginkgo biloba induces phase 2 genes through Keap1-Nrf2-ARE signaling pathway,” *Life Sciences*, vol. 80, no. 17, pp. 1586–1591, 2007.
- [50] H. Zhang and H. J. Forman, “Submicromolar concentrations of 4-hydroxynonenal induce glutamate cysteine ligase expression in HBE1 cells,” *Redox Report*, vol. 12, no. 1, pp. 101–106, 2013.
- [51] K. E. Iles and R.-M. Liu, “Mechanisms of glutamate cysteine ligase (GCL) induction by 4-hydroxynonenal,” *Free Radical Biology and Medicine*, vol. 38, no. 5, pp. 547–556, 2005.
- [52] C. Gorrini, I. S. Harris, and T. W. Mak, “Modulation of oxidative stress as an anticancer strategy,” *Nature Reviews Drug Discovery*, vol. 12, no. 12, pp. 931–947, 2013.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

