

Supplementary Table S1

Table S1. References related to the wound healing studies of diabetic foot

ulcers.[1-8]

Reference	Species	STZ dose (mg/kg)	Period (d)	Blood glucose level (mg/dL)
Theranostics. 2019, 9, 477-490	Male C57BL/6 mice (18-22 g)	60	5	250
J Diabetes Res. 2019, 2019, Article ID 1897174	Male Sprague-Dawley rats	65	14	~540
Wound Repair Regen. 2019, 27, 69-79	Male Wistar rats (4 months old)	50	7	300
Int Wound J. 2019, 16, 144-152	CD hairless rats (9-to10-week-old)	65	-	350
Sci Rep. 2017, 7, 11631	Female albino Wistar rats (5-day-old)	70	63	300
J Diabetes Res. 2016, 2016, Article ID 5782904	Male Sprague-Dawley rats (300–325 g)	70	28	~570
Drug Des Dev Ther. 2016, 2016, 1715-1730	Male Wistar rats (150-250 g)	65	3	~180
Comp Med. 2012, 62, 37-48	Male Wistar rats (250-350 g)	65	8	250

References

[1] R.-F. Chen, C.-H. Chang, C.-T. Wang et al., "Modulation of vascular endothelial growth factor and mitogen-activated protein kinase-related pathway involved in extracorporeal shockwave therapy accelerate diabetic wound healing" Wound Repair and Regeneration, vol. 27, pp. 69-79, 2019.

- [2] J.J. Mendes, C.I. Leandro, D.P. Bonaparte and A.L. Pinto, "A rat model of diabetic wound infection for the evaluation of topical antimicrobial therapies" *Comparative Medicine*, vol. 62, pp. 37-48, 2012.
- [3] A.A. Muhammad, P. Arulselvan, P.S. Cheah, F. Abas and S. Fakurazi, "Evaluation of wound healing properties of bioactive aqueous fraction from *Moringa oleifera* Lam on experimentally induced diabetic animal model" *Drug Design, Development and Therapy*, vol. 10, pp. 1715-1730, 2016.
- [4] J. Ning, H. Zhao, B. Chen et al., "Argon Mitigates Impaired Wound Healing Process and Enhances Wound Healing In Vitro and In Vivo" *Theranostics*, vol. 9, pp. 477-490, 2019.
- [5] S. Qiu, Y. Jia, Y. Sun et al., "Von Hippel-Lindau (VHL) Protein Antagonist VH298 Improves Wound Healing in Streptozotocin-Induced Hyperglycaemic Rats by Activating Hypoxia-Inducible Factor- (HIF-) 1 Signalling" *Journal of diabetes research*, vol. 2019, pp. Article ID 1897174, 2019.
- [6] M. Saidian, J.R.T. Lakey, A. Ponticorvo et al., "Characterisation of impaired wound healing in a preclinical model of induced diabetes using wide-field imaging and conventional immunohistochemistry assays" *International wound journal*, vol. 16, pp. 144-152, 2019.
- [7] C.O.-L. Yu, K.-S. Leung, J.L. Jiang et al., "Low-Magnitude High-Frequency Vibration Accelerated the Foot Wound Healing of n5-streptozotocin-induced Diabetic Rats by Enhancing Glucose Transporter 4 and Blood Microcirculation" *Scientific Reports*, vol. 7, pp. 11631-11631, 2017.
- [8] Y. Zhang, S.A. McClain, H.-M. Lee et al., "A Novel Chemically Modified Curcumin "Normalizes" Wound-Healing in Rats with Experimentally Induced Type I Diabetes: Initial Studies" *Journal of Diabetes Research*, vol. 2016, pp. Article ID 5782904, 2016.