

Myocardial adipose triglyceride lipase overexpression protects against burn-induced cardiac lipid accumulation and injury

Lingfei Li^{1,2}, Xingyue Zhang^{1,2}, Qiong Zhang^{1,2}, Jiezhai Jia^{1,2}, Junhui Zhang^{1,2}, Dongxia Zhang^{1,2}, Huapei Song^{1,2}, Bing Chen³, Jiongyu Hu^{2,3*}, Yuesheng Huang^{1,2*}

¹Institute of Burn Research, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing, China.

²State Key Laboratory of Trauma, Burns and Combined Injury, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing, China.


³Endocrinology Department, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing, China.

*Corresponding authors:

Yuesheng Huang, Institute of Burn Research, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing, 400038, China. Phone and fax: +86 023 68766023. E-mail: yshuang1958@163.com;

Jiongyu Hu, Endocrinology Department, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing, 400038, China. Phone and fax: +86 023 68773162. E-mail: jiongyuhu@163.com.

Supplementary Information



Supplementary Figure S1 Detection of fasting blood glucose. Fasting blood glucose (fasting 8 hours before measurement) was determined between WT, ATGL-Het and MHC-ATGL mice with or without burn injury. $n = 8$. Data are shown as the mean \pm SEM. $*P < 0.05$. P values were derived from one-way ANOVA with the appropriate posttest.