Review Article

The Role of Antioxidants in Ameliorating Cyclophosphamide-Induced Cardiotoxicity

**Muluken Altaye Ayza , Kaleab Alemayehu Zewdie , Bekalu Amare Tesfaye ,
Dawit Zewdu Wondafrash , and Abera Hadgu Berhe **

Department of Pharmacology and Toxicology, School of Pharmacy, Mekelle University, Mekelle, Ethiopia

Correspondence should be addressed to Muluken Altaye Ayza; mulealto1@gmail.com

Received 10 January 2020; Revised 9 April 2020; Accepted 21 April 2020; Published 11 May 2020

Academic Editor: Shane Thomas

Copyright © 2020 Muluken Altaye Ayza et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The chemotherapeutic and immunosuppressive agent cyclophosphamide has previously been shown to induce complications within the setting of bone marrow transplantation. More recently, cardiotoxicity has been shown to be a dose-limiting factor during cyclophosphamide therapy, and cardiooncology is getting wider attention. Though mechanism of cyclophosphamide-induced cardiotoxicity is not completely understood, it is thought to encompass oxidative and nitrative stress. As such, this review focuses on antioxidants and their role in preventing or ameliorating cyclophosphamide-induced cardiotoxicity. It will give special emphasis to the cardioprotective effects of natural, plant-derived antioxidants that have garnered significant interest in recent times.

1. Introduction

1.1. Drug-Induced Cardiotoxicity. Drug-induced cardiotoxicity poses a serious risk to human health, and cardiooncology is currently becoming an important concern [1]. Antineoplastic treatments led to increased overall and progression-free survival in the management of an increasing number of malignancies [2]. However, as cancer survival has improved with advancing therapies, late cardiovascular adverse effects have become an important management issue, mainly in childhood cancers, leukaemia, lymphoma, and breast cancer. In patients diagnosed with early stage breast cancer, cardiovascular disease is the major cause of mortality [3]. Even though anticancer drugs are targeted against malignant cells, they are also toxic to normal cells [4].

Patients who survived cancer, when compared to their healthy counterparts, are at an increased risk of cardiovascular-related mortality, which might be due to myocardial infarction with coronary artery disease, cardiomyopathy with congestive heart failure, and cerebrovascular events [5, 6]. Patients on cancer chemotherapy can be considered as a stage A heart failure group, patients with

increased risk of heart failure and do not have structural heart disease [7, 8]. Total dose of the anticancer agent patient received, rate of drug administration, extent of radiation of the mediastinum, age, being female, previous history of heart disease, and increased blood pressure are risk factors to develop cardiotoxicity [9, 10].

Antineoplastic agents are well known to cause a wide array of toxicities including cardiac dysfunction leading to heart failure, arrhythmias, myocardial ischemia, hypertension, thromboembolism, myocarditis, and pericarditis [11]. Anthracyclines are the best known of the chemotherapeutic agents that cause cardiotoxicity. In addition, alkylating drugs, including cisplatin, cyclophosphamide, ifosfamide, carmustine, chlormethine, busulfan, and mitomycin, are also linked with cardiac toxicity [9].

1.2. Cyclophosphamide. Cyclophosphamide is an alkylating, anticancer agent which was first characterized in experiments on rat tumors. It is an oxazaphosphorine-substituted nitrogen mustard, with strong cytotoxic and immunosuppressive activity [12]. It is the mainstay of most preparative regimens for organ transplant and a

FIGURE 1: Major metabolic pathway of cyclophosphamide.

broadly active anticancer, immunosuppressive agent used in combination chemotherapy for Hodgkin's disease, non-Hodgkin's lymphoma, leukaemia, rheumatoid arthritis, Burkitt's lymphoma, lupus erythematosus, multiple sclerosis, neuroblastoma, multiple myeloma, endometrial cancer, breast cancer, and lung cancer. At high dosages, cyclophosphamide can be used alone or in combination with bone marrow transplant in the management of solid tumors and lymphomas [9, 13].

The electrophilic nature of the alkyl group enables the drug to react with nucleophilic moieties of DNA or proteins, and this leads to the covalent transfer of an alkyl group. Cyclophosphamide is a prodrug that requires an activation step by cytochromes (P450) in the liver [14]. As shown in Figure 1, the introduction of the hydroxyl group to the oxazaphosphorine ring generates 4-hydroxycyclophosphamide, which cooccurs in equilibrium with its isomer, aldophosphamide. Then, aldophosphamide is converted into two compounds, phosphoramidate mustard and acrolein (Figure 1) [15].

Phosphoramidate mustard forms a highly reactive cyclic aziridinium cation, which can react with the N(7) of the guanine and with cytidine from the DNA. Due to the two reactive moieties in the molecule, intrastrand and interstrand cross-links can be formed [16]. This leads to inhibition of DNA replication and apoptosis, with the active metabolites also having cell-cycle-independent activity. The specific mechanism of action of the compound used in managing autoimmune diseases has been postulated to include apoptosis, B-cell suppression, which will lead to decreased immunoglobulin G production and decreased production of adhesion molecules and cytokines [12].

Acrolein is the cause of hemorrhagic cystitis, one of the major toxicities of cyclophosphamide therapy. Other toxicities include bone marrow suppression, cardiotoxicity, gonadal toxicity, and carcinogenesis, with cumulative doses being the principal risk factor [15]. Additionally, administration of a single, large dose of cyclophosphamide is capable of causing hemorrhagic cell death, leading to heart failure or even death [17].

2. Pathophysiology of Cyclophosphamide-Induced Cardiotoxicity

Cyclophosphamide-induced cardiac damage is dose dependent, and the total dose of an individual course is the best indicator of toxicity, with patients who receive greater than 150 mg/kg or 1.55 g/m²/day, which are at a high risk for cardiotoxicity [18]. The dose-limiting factor during cyclophosphamide therapy is cardiotoxicity [19], which is irreversible [20]. Fatal cardiomyopathy has been reported among 2–17% of patients taking cyclophosphamide. It is dependent on the regimen and the particular patient population characteristics [21]. Overall, cyclophosphamide-induced cardiotoxicity affects between 7 and 28% of patients taking the drug [13].

The pathophysiology of cyclophosphamide-induced cardiac damage is poorly understood [10], although its metabolites are thought to induce oxidative stress and direct endothelial capillary damage with resultant extravasation of proteins, erythrocytes, and toxic metabolites. In the presence of toxic metabolites, breakdown of endothelial cells contributes to direct damage to the myocardium and capillary blood vessels resulting in edema, interstitial hemorrhage, and formation of microthrombosis [22, 23].

Endothelial cells are more susceptible to cyclophosphamide-induced damage than other cells ([24]); this might be associated with their high proliferation rate [25]; cyclophosphamide-induced reactive oxygen species generation can also lead to a reduction in nitric oxide bioavailability, thus leading to compromised endothelial function [26].

Molecular mechanisms of cyclophosphamide-mediated cardiac damage are currently being postulated, potentially leading to better preventative strategies to treat cardiotoxicity. It has been shown that treatments with cyclophosphamide inhibited heart-type fatty acid-binding proteins and carnitine palmitoyltransferase-I gene expression in cardiac tissues [27]. Inhibition of these pathways leads to decreased production of adenosine triphosphate and accumulation of toxic metabolites from fatty acid oxidation, consequently leading to cardiomyopathy [28]. Heart-type fatty acid-

FIGURE 2: Molecular mechanisms involved in cyclophosphamide-induced cardiotoxicity.

binding protein can be used as an early diagnostic marker of chemotherapy-induced cardiotoxicity [29]. In addition, carnitine deficiency can aggravate cardiotoxicity and it is important to monitor serum and urinary carnitine levels [30]. Carnitine supplementation showed beneficial effects in various cyclophosphamide-induced toxicities [31–34].

Cyclophosphamide administration affects the ability of the heart mitochondria to retain accumulated calcium [35]. Calcium leak from sarcoplasmic reticulum can lead to mitochondrial calcium overload, leading to reduced production of adenosine triphosphate and increased release of ROS [36]. It is reported that improving mitochondrial function through supplementation of lupeol and its ester can protect heart from cyclophosphamide-induced toxicity [37].

Cyclophosphamide is found to promote proinflammatory cytokines [38, 39]. It enhanced nuclear factor-kappa B (NF- κ B) phosphorylation, both expression and serum levels of cyclooxygenase-2 (COX-2), tumor necrosis factor-alpha (TNF- α), and interleukin-1 beta (IL-1 β) [40, 41]. Nuclear erythroid 2-related factor 2 (Nrf2) and NF- κ B are considered as an important molecular target for the anti-inflammatory and antioxidant chemicals for cytoprotection during cyclophosphamide therapy [42, 43]. It has been reported that inhibition of the NF- κ B/TNF- α pathway prevented cyclophosphamide-induced multiple organ toxicity including the heart, kidney, and liver [44, 45].

It has been reported that p53 expression plays an important role in apoptosis [46, 47]. Reduction in apoptosis, infarct size, and hemodynamic parameter improvement can be achieved by inhibiting p53 [48]. Cyclophosphamide-induced activation of p53 protein is considered as one of

the possible mechanisms for cardiomyopathy, and it is reported that probucol supplementation restored cyclophosphamide-induced upregulation of p53 and reversed apoptosis in cardiomyocytes [49].

Cyclophosphamide activates the p38 mitogen-activated protein kinase (p38-MAPK) pathway, and it can induce an oxidative injury [50]. Cyclophosphamide-enhanced proinflammatory/proapoptotic activities are reported to result in cardiomyopathy, myocardial infarction, and heart failure [13]. Rutin attenuated cyclophosphamide-induced oxidative stress and inflammation through downregulating TNF- α , IL-6, and expressions of p38-MAPK, NF- κ B, and COX-2 (Figure 2) [45].

Cyclophosphamide induces the calcineurin-mediated dephosphorylation of nuclear factor of activated T-cell (NFAT); it belongs to the family of calcium-regulated transcription factors. Unphosphorylated/active GSK-3 β phosphorylates eIF2, NFAT, and c-jun and thus contributes significantly to cardiac hypertrophy inhibition/protection [13]. Cyclosporine A prevented NFAT nuclear translocation and reversed cyclophosphamide-induced cardiac damage [35, 51].

In general, mechanisms of cyclophosphamide-induced cardiotoxicity encompass oxidative and nitrative stress, protein adduct formation which leads to cardiomyocyte inflammation, altered calcium homeostasis, programmed cell death, swelling of the cardiomyocytes, nuclear splitting, vacuolization, and alteration in signaling pathways. These events result in diseases of the heart muscle including heart failure, if left undiagnosed or untreated, and may result in death [13]. Further supporting a role of cyclophosphamide-induced

oxidative stress in the evident cardiotoxicity of the compound, exposure of rats to cyclophosphamide resulted in reduced pulmonary glutathione (GSH) content, GSH reductase (GRx), glucose-6-phosphate dehydrogenase, GSH peroxidase (GPx), and superoxide dismutase (SOD) activities [24].

3. Current Management for Cyclophosphamide-Induced Cardiotoxicity

Clinical management of cardiovascular diseases (CVDs) involves multiple drugs (angiotensin-converting enzyme inhibitors, blockers of angiotensin-II receptor, calcium channel blockers, β -blockers, aldosterone inhibitors, aspirin, statins, and warfarin), and others include diuretics, digoxin, and nitrates [52–54]. It is a common practice to use those medications in combination for the management of CVDs, and these lead to increased side effect and drug interactions [55].

These same preventive strategies can be considered for ischemia, heart failure, arrhythmia, hypertension, and arterial thromboembolism associated with cyclophosphamide-induced cardiotoxicity. Primary prevention may include widespread treatment of all patients who are potentially on cardiotoxic cancer treatments or early diagnosis of subclinical cardiac injury with targeted treatment.

According to the Canadian Cardiovascular Society recommendation, even though the recommendation is weak, patients believed to be at a high risk for cancer treatment-related left ventricular dysfunction, an angiotensin-converting enzyme inhibitor, angiotensin receptor blocker, and/or β -blocker, and/or statin can be considered to decrease the risk of cardiac damage [56]. Valsartan, an angiotensin receptor blocker, showed a strong effect in preventing acute cyclophosphamide-, doxorubicin-, vincristine-, and prednisolone-induced cardiotoxicity [57]. A nonselective β -blocker, carvedilol, with antioxidant activity and nebivolol, a selective β -blocker with a nitric oxide donor capacity, were reported to have an advantageous effect on antineoplastic-associated cardiac damage [58].

Mild to moderate heart failure and small pericardial effusions generally resolve within a few days to weeks after stoppage of cyclophosphamide. In the presence of suspected hemorrhagic myocarditis, cardiac tamponade, and cardiogenic shock, timely recognition and involvement of the intensive care unit or coronary care unit are vital. These patients need aggressive monitoring and circulatory support [22].

4. Natural Antioxidants for the Management of Cyclophosphamide-Induced Cardiotoxicity

The use of plants and plant-based products in the treatment of ailments has been known to mankind from ancient times [59]. Various natural antioxidants have originated from medicinal plants, which are used for the treatment of different ailments throughout the world, and there has been a significant interest in finding natural antioxidants from plant sources [60]. Diseases and drug-induced toxicities with the

underlying cause of oxidative stress can be effectively managed with plants having antioxidant activity. Apart from being rich sources of antioxidants, phytochemicals are also known to impede the progression of cardiac tissue damage [59]. These compounds could serve as one of the valuable sources in industrial pharmaceutical research and can be treated as a complementary and alternative medicine.

Various medicinal plants showed cardioprotective activity against cyclophosphamide-induced cardiotoxicity in different preclinical studies (Table 1). In addition, xanthine-oxidase inhibitors (allopurinol and febuxostat) and nicorandil (vasodilatory drug used to treat angina) were also found to reverse cardiac damages induced by cyclophosphamide in male Wistar rats (Table 2).

5. Future Hopes and Hurdles Associated with Cardioprotective Antioxidants

Antioxidants such as flavonoids, flavones, isoflavones, anthocyanin, catechins, and isocatechins are the responsible ones for the antioxidant activity of spices and herb [90]. These led supplementation of antioxidants to be a popular practice to maintain optimal body function [91]. Polyphenols may reduce cholesterol absorption and upregulate hepatic mRNA abundance for the LDL receptor, reductions in plasma TG, yielding a reduced amount of LDL in circulation, and polyphenols were found to exert anti-inflammatory effects, thereby reducing the formation of cytokines involved in cellular adhesion [92]. The production of vasodilating factors like nitric oxide, endothelium-derived hyperpolarizing factor, and prostacyclin was enhanced by plant polyphenols. These plant phenols were also found to inhibit the production of vasoconstrictor endothelin-1 in endothelial cells and inhibit the expression of two main proangiogenic factors, matrix metalloproteinase-2, and vascular endothelial growth factor in smooth muscle cells [93]. Flavonoids can also improve endothelial function, and the primary mechanism for this is that the effect is nitric oxide production [94].

Even though the results were not posted, currently, there are different agents under clinical trial, including enalapril for prevention of chemotherapy-induced cardiotoxicity in high-risk patients (NCT00292526), nutritional supplement sulforaphane on doxorubicin-associated cardiac dysfunction (NCT03934905), estimation of the effects of ACE inhibitors and β blockers in the management of cardiotoxicity in oncologic patients (NCT02818517), cardiotoxicity prevention in breast cancer patients treated with anthracyclines and/or trastuzumab using bisoprolol and ramipril (NCT02236806), carvedilol effect in preventing chemotherapy-induced cardiotoxicity (NCT01724450), prevention of chemotherapy-induced cardiotoxicity in children with bone tumors and acute myeloid leukaemia using capoten (captopril) (NCT03389724), and statins to prevent the cardiotoxicity from anthracyclines (NCT02943590), and others are under investigation. These agents might be the future hopes for the management of chemotherapy-induced cardiotoxicity.

Even though antioxidants like flavonoids have a great hope in the future clinical scenario of cardioprotection [95], the importance of antioxidants is currently in question due

TABLE 1: Effect of medicinal plants and isolates against cyclophosphamide-induced cardiotoxicity.

References	Animals used	Method and intervention	Major findings
Asiri [49]	Male Wistar albino rats	Rats were administered with the same doses of corn oil (control) and probucol (61 mg/kg/day, i.p.), respectively, for one week before and one week after a single dose of CP (200 mg/kg, i.p.).	Probulcol prevented the development of CP-induced cardiotoxicity by a mechanism related, at least in part, to its ability to increase mRNA expression of antioxidant genes and to decrease apoptosis in cardiac tissues with the consequent improvement in mitochondrial oxidative phosphorylation and energy production. Concurrent administration of SLY and CUR with CP resulted significantly lower biochemical parameters and histopathological and immunohistochemical results than in the CP-only group. It can be concluded that the natural antioxidant SLY and CUR might have protective effects against CP-induced cardiotoxicity and oxidative stress in rats.
Avci et al. [61]	Female Wistar albino rats	Animals were treated with 100 mg/kg/day silymarin (SLY) by oral gavage for 14 days and 30 mg/kg/day CP intraperitoneally starting from the seventh day and 100 mg/kg/day curcumin (CUR) by gavage for 14 days plus 30 mg/kg/day CP intraperitoneally starting from the seventh day.	<i>Croton macrostachyus</i> reversed CP-induced elevations of cardiac troponin, ALT, AST, ALP, TC, and TG. These findings were further supported by histopathological findings. Treatment with the extract reduced the serum biomarkers (CK-MB, ALT, AST, ALP, TC, and TG) and increased the tissue antioxidant level. Histopathology of heart tissue was also improved. Mangiferin treatment resulted in decrement of the serum cardiac biomarkers (AST, ALT, ALP, CK-MB, CK-NAC, and LDH). Mangiferin increased tissue antioxidant levels (SOD, CAT, and GSH), and animals showed improvement in lipid profile, ECG parameters, histological score, and mortality. Vitamin E in a single oral dose failed to inhibit acute cardiotoxic activity of doxorubicin but suspended further progression of the heart muscle damage over the time. On the contrary, vitamin E did not attain cardioprotection against doxorubicin and CP in combination.
Ayza et al. [62]	Either sex of Sprague Dawley rats	Animals were treated with a single dose of CP (200 mg/kg, i.p.) on the first day followed by hydromethanolic crude extract and solvent fractions of <i>Croton macrostachyus</i> for 10 days.	
Baniya et al. [63]	Male Wistar rats	Animals received a single dose of CP (200 mg/kg) on the first day followed by ethanolic extract of <i>Citrus grandis</i> (L.) Osbeck at 250 and 500 mg/kg, p.o. for 10 days.	
Bhatt et al. [64]	Either sex of Wistar rats	Cyclophosphamide (200 mg/kg, i.p.) toxicity was induced on day 1. Then, rats were treated with 100 mg/kg of mangiferin for 10 days.	
Byelogrlic et al. [65]	Female BalbC/NIH mice	Animals were treated with vitamin E (100 IU/kg, orally) 24 hr before single bolus doses of doxorubicin (10 mg/kg, intravenously), or doxorubicin and CP (150 mg/kg, i.p.).	

TABLE 1: Continued.

References	Animals used	Method and intervention	Major findings
Cetik et al. [66]	Sprague-Dawley rats	Carvacrol administration was started three days before the CP application and continued till the end of experiment (six days).	Carvacrol at both the doses increased the GSH levels close to the control group GSH levels. Carvacrol at 5.0 and 10 mg/kg doses lowered the levels of serum ALT, AST, LDH, and CK-MB. Reduced inflammation and lipid peroxidation in the heart tissue and increase of serum GSH and total antioxidant capacity (TAS) levels were found when carvacrol was applied.
Chakraborty et al. [67]	Either sex Wistar albino rats	Rats were subjected to CP toxicity with the dose of (200 mg/kg i.p.) on day first. Then, treated with green tea extract (GTE) along with hydrochlorothiazide.	GTE dose dependently reduced CP-induced myocardial toxicity. Green tea when combined with hydrochlorothiazide reduced the associated side effects and exhibited myocardial protection.
Chakraborty et al. [68]	Male Wistar albino rats	Rats were treated with combination of curcumin (100, 50, 25 mg/kg, p.o.) and piperine (20 mg/kg, p.o.) for 10 days. All treated groups were subjected to CP (200 mg/kg, i.p.) toxicity on day 1.	Piperine incorporation with the doses of 50 and 25 mg/kg with curcumin exhibited a significant beneficial effect compared to the curcumin alone-treated group. Treatment with curcumin and piperine significantly modified the markers.
Conklin et al. [69]	Glutathione S-transferase (GSTP) wild-type (WT) and GSTP-P1/P2 null mice	To examine CP cardiotoxicity, WT and GSTP-null mice were treated with saline (control) or 100, 200, and 300 mg/kg CP.	According to the findings, myocardial GSTP levels are likely to be key determinants of CP cardiotoxicity. GSTP is a highly regulated enzyme that is readily induced by different environmental factors, diet constituents such as garlic organosulfur compounds, coffee, and chemopreventive agents such as selenocysteine conjugates. Therefore, GSTP induction by such agents could attenuate CP toxicity, and conversely, disturbed metabolic states such as obesity, which are associated with downregulation of GSTP, and could enhance the cardiotoxicity of CP treatment.
El-Agamy et al. [70]	Male Wistar rats	Cardiotoxicity was induced by single injection of CP (200 mg/kg, i.p.). Methyl palmitate (MP) was administered at two different dose levels (300 and 400 mg/kg) for 10 days before and 7 days after CP injection.	Animals treated with MP showed significant attenuation of ECG changes. MP supplementation significantly lowered the elevated cardiac markers and improved cardiac lesions, which was more prominent at the higher dose. MP treatment significantly decreased MDA content and enhanced the antioxidant parameters (SOD and GSH), and it significantly decreased the expression of TLR4 and NF- κ B p65. MP supplementation suppressed inflammatory

TABLE 1: Continued.

References	Animals used	Method and intervention	Major findings
Gado et al. [71]	Male Swiss albino rats	Curcumin (200 mg/kg, i.p.) was administered for 8 consecutive days followed by a single dose of CP (150 mg/kg, i.p.).	cytokines (TNF- α and nitrite/nitrate) and reduced apoptosis. Serum LDH and CPK were decreased significantly with the curcumin administration. Curcumin treatment significantly decreased MDA, NO(x), and restore GSH level in the cardiac tissue. Histological alterations were also found to be improved.
Gunes et al. [72]	Male Sprague Dawley rats	Animals received respective selenium (Se) doses (0.5 or 1 mg/kg) for 6 days and then a single dose of CP administered on the sixth day. On day 7, the animals were sacrificed.	Based on microscopic evaluation, tissue damage was noticeably lower in CP plus Se groups. Additionally, 1 mg/kg Se was more protective than 0.5 mg/kg Se. It can be concluded that Se can be a potential candidate to ameliorate CP-induced cardiotoxicity which may be related to its antioxidant activity.
Iqbal et al. [73]	Male Swiss albino mice	Animals were treated with nerolidol (NER) (200 and 400 mg/kg p.o.) and fenofibrate (FF) 80 mg/kg, p.o. for 14 days along with a single dose of CP 200 mg/kg i.p. on the 7th day.	NER 400 significantly reversed cardiotoxic effects of CP and showed cardioprotective activity which was comparable with FF 80. However, NER 200 did not show significant cardioprotective activity.
Mansour and Hasan [74]	Male Wistar albino rats	Rats were pretreated with N-acetylcysteine (200 mg/kg) for 5 days; 1 hour after the last dose, rats were injected with CP (200 mg/kg).	Treatment with N-acetylcysteine significantly decreased serum levels of ALT, AST, CK, and LDH. Decrease in the NOx, MDA levels and TNF- α , SOD, catalase, GSHPx, and GST levels were increased.
Mythili et al. [75]	Male Wistar albino rats	Rats received single injection of CP (200 mg/kg, i.p.) to induce cardiotoxicity, then followed by dl- α -lipoic acid treatment (25 mg/kg for 10 days).	Normalized lipid peroxidation and antioxidant defenses were observed in the dl- α -lipoic acid-treated rats.
Mythili et al. [76]	Male Wistar albino rats	Rats were injected with a single dose of CP (200 mg/kg, i.p.) to induce cardiotoxicity, and then rats were treated with lipoic acid (25 mg/kg, orally for 10 days).	Treatment with lipoic acid reversed the abnormalities in the lipid levels and activities of lipid-metabolizing enzymes to near normalcy.
Mythili et al. [77]	Male Wistar albino rats	Rats received CP (200 mg/kg i.p.), which is immediately followed by lipoic acid (25 mg/kg orally for 10 days).	Lipoic acid effectively reversed abnormal biochemical changes to near normalcy. Based on the results, lipoic acid showed a protective role of lipoic acid in CP-induced cardiotoxicity.

TABLE 1: Continued.

References	Animals used	Method and intervention	Major findings
Nagi et al. [78]	Male Wistar albino rats	Rats received thymoquinone (50 mg/l in drinking water) for 5 days before a single dose of CP (200 mg/kg, i.p.) and continued thereafter until day 12. On day 13, animals were sacrificed.	Thymoquinone reversed CP-induced increase in serum CK-MB and LDH. Complete reversal of the CP-induced increase in serum cholesterol, triglycerides, urea, and creatinine to the control values. CP-induced increase in TBARS and NO(x) and a decrease in GSH, GPx, and CAT were reversed by thymoquinone supplementation. Thymoquinone supplementation to CP-treated rats completely reversed the increase in TNF- α induced by CP.
Omole et al. [79]	Male Wistar rats	Rats were pretreated with 200 and 400 mg/kg/d Kolavorin, orally for 14 days followed by CP (50 mg/kg/d, i.p.) for 3 days.	Kolavorin pretreatment increased food consumption, body weight, and attenuated the biochemical and histological changes. It was reported that kolavorin inhibited oxidative stress and preserved the activity of antioxidant enzymes.
Sekeroğlu et al. [80]	Male Swiss albino mice	After treatment with <i>Viscum album</i> and quercetin for 7 days, rats were administered CP (40 mg/kg, i.p) on days 8 and 9 of the experiment. Total treatment period was 10 days.	Treatments decreased the levels of antioxidant enzymes, glutathione-S-transferases; reduced glutathione and mitotic index were observed. Quercetin completely and <i>Viscum album</i> partly ameliorated almost all of the examined parameters when given together with CP.
Senthilkumar et al. [81]	Male albino Wistar rats	Animals were cotreated with CP intraperitoneally dissolved in saline, in a dose of 150 mg/kg b.w. and different doses of squalene for the first 2 days, and squalene treatment was followed continuously, daily for 10 days up to the end of the experimental period.	Squalene oral treatment exerted protection to the heart, kidney, and liver at a dose of 0.4 ml/day/rat. Histopathological examinations also confirmed the protective efficacy of squalene. It can be concluded that squalene may be efficacious as a cytoprotectant in CP-induced toxicities.
Shalaby et al. [82]	Male Sprague-Dawley albino rats	Rats received <i>Zingiber officinale</i> 200 mg/kg/day orally followed by a single dose of CP (150 mg/kg i.p.).	Results showed significant improvement in the <i>Zingiber officinale</i> -treated group. Based on their conclusion, the cardiotoxic effect of CP might be prevented by <i>Zingiber officinale</i> supplementation.
Shanmugarajan et al. [83]	Male Wistar rats	Rats were treated with the methanolic leaf extract of <i>Ficus hispida</i> Linn. for 10 consecutive days following CP-induced oxidative myocardial injury on the first day.	Treatment with <i>Ficus hispida</i> Linn. decreased serum cardiac biomarkers (CPK, LDH, AST, and ALT), and these were increased in the heart tissue. <i>Ficus hispida</i> Linn. increased the levels of enzymic antioxidants (SOD, CAT, GPx, GSH, and GRx).

TABLE 1: Continued.

References	Animals used	Method and intervention	Major findings
Song et al. [84]	Male ICR mice	Animals were injected with a single dose of CP (200 mg/kg i.p.) followed by the intragastric treatment with ferulic acid (FA) (50, 100 mg/kg) for 7 consecutive days.	FA significantly decreased the serum levels of cardiac biomarkers, IL-6, IL-1 β , and TNF- α in CP-injected mice. Additionally, FA effectively reduced the total numbers of WBCs, RBCs, platelets, and hemoglobin content. FA also attenuated the histological changes of the heart tissues caused by CP. Moreover, western blot demonstrated that FA inhibited the phosphorylations of the NF- κ B signaling pathway in CP-stimulated cardiac tissues.
Sudharsan et al. [85]	Male Wistar albino rats	Rats were injected with a single dose of CP (200 mg/kg, i.p.) and treated with lupeol and lupeol linoleate (50 mg/kg).	Lupeol and its ester reversed alterations of serum lipoproteins and lipid fractions in both serum and cardiac tissue. It was found that lupeol linoleate was more effective than lupeol.
Swamy et al. [86]	Male Wistar albino rats	Cardiotoxicity was induced by administering CP (200 mg/kg, i.p.) single injection. <i>Saraca indica</i> (200 and 400 mg/kg, p.o.) was daily for 10 days.	Treatment with <i>Saraca indica</i> reversed the status of cardiac biomarkers (CK, CK-MB, LDH, AST, ALT, and ALP) ECG, oxidative enzymes (GSH, SOD, and CAT), and lipid profile.
Todorova et al. [87]	Male Fischer 344 rats	After 2 d of prefeeding with glutamine (GLN) or glycine (GLY) by gavage, the rats were randomized into one of six groups receiving a lethal intraperitoneal dose of CP (450 mg/kg), a sublethal dose of CP (200 mg/kg), or saline (control).	The results showed that dietary GLN decreased cardiac necrosis and maintained normal cardiac GSH levels. GLN protected against the acute cardiotoxic effects of CP and significantly improved the short-term survival after lethal and sublethal doses of CP.

TABLE 2: The effect of currently available drugs against cyclophosphamide-induced cardiotoxicity.

References	Animals used	Methods and intervention	Main findings
El-Sheikh et al. [88]	Male Wistar rats	Rats were treated with allopurinol (ALL) 100 mg/kg/day or febuxostat (FEB) 10 mg/kg/day which were administered orally to rats in the presence and absence of CP (200 mg/kg i.p. single dose at the ninth day) treatment.	Based on the results, both xanthine oxidase (XO) inhibitors, ALL and FEB, ameliorated CP-induced cardiotoxicity. Though only FEB showed protective activity against CP-induced myelotoxicity, however ALL might aggravate myelotoxicity. Based on the findings, ROS and XO enzymatic pathways may largely participate in the mechanism of pathogenesis of cardiac and bone marrow toxicities related to CP exposure.
Refaie et al. [89]	Male Wistar albino rats	Rats were administered with nicorandil (NIC) (3 mg/kg/day) alone and coadministered with nitro- ω -l-arginine (L-NNA) (25 mg/kg/day) and glibenclamide (GP) (5 mg/kg/day) orally for 5 days and injected with CP (150 mg/kg/day) i.p. on 4 th and 5 th days.	NIC reversed CP-induced cardiotoxicity by its potassium channel opening effect, stimulating eNOS gene expression, anti-inflammatory, antiapoptotic, and antioxidant properties. Cotreatment with GP or L-NNA decreased the protective effect of NIC.

to their less effectiveness in an *in vivo* study. These failures of antioxidants in preventing/treating diseases have become the main obstacle in the clinical scenario [96].

As concluded by Guallar et al., known antioxidants like vitamin E, β -carotene, vitamin A and B supplements, and folic acid are ineffective for the prevention of mortality and morbidity due to chronic diseases [97].

This failure might be due to different reasons including antioxidant-related reasons including testing incorrect antioxidant or combination of antioxidants; there might be differences between synthetic and dietary source antioxidants, reductive stress (i.e., too much antioxidant capacity), and it may also be related to patient or clinical trials [98].

Unconjugated flavonoid plasma level rarely exceeds $1\ \mu\text{M}$, and metabolites of flavonoids have lower antioxidant activity. Since plasma total antioxidant capacities (TAC) are often in the range of 1 mM or more, it is difficult to picture how an additional $1\ \mu\text{M}$ polyphenol could exert an *in vivo* antioxidant effect. Antioxidants like flavonoids and other phenols are complex molecules and have multiple potential targets/actions in addition to antioxidant activity. These may include inhibition of different enzymes including cyclooxygenase, lipoxygenase, xanthine oxidase, matrix metalloproteinases, angiotensin-converting enzyme, proteasome, and cytochrome P450, affecting signal transduction pathways. Flavonoids may also interact with cellular drug transport systems [99]. These issues need to be addressed in the future.

6. Conclusion

Cyclophosphamide is a known anticancer and immunosuppressive agent that becomes effective after metabolic activation in the liver. Its wider clinical application is currently limited by its toxicity. Cardiotoxicity, which is associated with oxidative and nitrative stress, is one of the toxicities limiting the clinical use of cyclophosphamide. Different natural, plant-derived antioxidants (summarized in this review) showed significant cardioprotective effects in *in vivo* preclinical studies. However, further investigations aimed at improving their efficacy are required. Facilitating translational clinical research on those shown to be safe and effective in the preclinical studies should also be considered, lest the evidences from the preclinical studies would only be left to be discussed in scientific meetings and publications.

Abbreviations

CAT: Catalase
 CP: Cyclophosphamide
 CVDs: Cardiovascular diseases
 GPx: Glutathione peroxidase
 GRx: Glutathione reductase
 GSH: Glutathione.

Conflicts of Interest

The authors declare no conflict of interest.

References

- [1] C. Cadeddu Dessalvi, M. Deidda, D. Mele et al., "Chemotherapy-induced cardiotoxicity: new insights into mechanisms, monitoring, and prevention," *Journal of Cardiovascular Medicine*, vol. 19, no. 7, pp. 315–323, 2018.
- [2] G. Varricchi, P. Ameri, C. Cadeddu et al., "Antineoplastic drug-induced cardiotoxicity: a redox perspective," *Frontiers in Physiology*, vol. 9, no. 167, 2018.
- [3] R. Conyers, B. Costello, A. la Gerche et al., "Chemotherapy-related cardiotoxicity: are Australian practitioners missing the point?," *Internal Medicine Journal*, vol. 47, no. 10, pp. 1166–1172, 2017.
- [4] S. Kumar, R. K. Gupta, and N. Samal, "Cyclophosphamide cardiotoxicity," *The National Medical Journal of India*, vol. 5, no. 1, pp. 17–19, 1992.
- [5] M. Csapo and L. Lazar, "Chemotherapy-induced cardiotoxicity: pathophysiology and prevention," *Medicine and Pharmacy Reports*, vol. 87, no. 3, pp. 135–142, 2014.
- [6] R. W. Loar, C. V. Noel, H. Tunuguntla, J. L. Colquitt, and R. H. Pignatelli, "State of the art review: chemotherapy-induced cardiotoxicity in children," *Congenital Heart Disease*, vol. 13, no. 1, pp. 5–15, 2018.
- [7] D. Cardinale and C. M. Cipolla, "Assessment of cardiotoxicity with cardiac biomarkers in cancer patients," *Herz*, vol. 36, no. 4, pp. 325–332, 2011.
- [8] L. B. Daniels, P. Clopton, K. Jiang, B. Greenberg, and A. S. Maisel, "Prognosis of stage A or B heart failure patients with elevated B-type natriuretic peptide levels," *Journal of Cardiac Failure*, vol. 16, no. 2, pp. 93–98, 2010.
- [9] V. B. Pai and M. C. Nahata, "Cardiotoxicity of chemotherapeutic agents: incidence, treatment and prevention," *Drug Safety*, vol. 22, no. 4, pp. 263–302, 2000.
- [10] I. Taniguchi, "Clinical significance of cyclophosphamide-induced cardiotoxicity," *Internal Medicine*, vol. 44, no. 2, pp. 89–90, 2005.
- [11] A. Colombo, C. A. Meroni, C. M. Cipolla, and D. Cardinale, "Managing cardiotoxicity of chemotherapy," *Current Treatment Options in Cardiovascular Medicine*, vol. 15, no. 4, pp. 410–424, 2013.
- [12] J. Kim and J. J. Chan, "Cyclophosphamide in dermatology," *Australasian Journal of Dermatology*, vol. 58, no. 1, pp. 5–17, 2017.
- [13] A. Iqbal, M. K. Iqbal, S. Sharma et al., "Molecular mechanism involved in cyclophosphamide-induced cardiotoxicity: old drug with a new vision," *Life Sciences*, vol. 218, pp. 112–131, 2019.
- [14] S. Puyo, D. Montaudon, and P. Pourquier, "From old alkylating agents to new minor groove binders," *Critical Reviews in Oncology/Hematology*, vol. 89, no. 1, pp. 43–61, 2014.
- [15] A. Emadi, R. J. Jones, and R. A. Brodsky, "Cyclophosphamide and cancer: golden anniversary," *Nature Reviews Clinical Oncology*, vol. 6, no. 11, pp. 638–647, 2009.
- [16] M. Ahlmann and G. Hempel, "The effect of cyclophosphamide on the immune system: implications for clinical cancer therapy," *Cancer Chemotherapy and Pharmacology*, vol. 78, no. 4, pp. 661–671, 2016.
- [17] M. Molinaro, P. Ameri, G. Marone et al., "Recent advances on pathophysiology, diagnostic and therapeutic insights in cardiac dysfunction induced by antineoplastic drugs," *BioMed Research International*, vol. 2015, no. 10, pp. 1–14, 2015.

- [18] S. Kusumoto, H. Kawano, T. Hayashi et al., "Cyclophosphamide-induced cardiotoxicity with a prolonged clinical course diagnosed on an endomyocardial biopsy," *Internal Medicine*, vol. 52, no. 20, pp. 2311–2315, 2013.
- [19] Y. Kanda, T. Matsumura, K. Maki, T. Matsuyama, S. Chiba, and H. Hirai, "Fatal cardiac toxicity in two patients receiving same-day administration of cyclophosphamide and cytarabine as conditioning for hematopoietic stem cell transplantation," *Haematologica*, vol. 86, no. 9, pp. 1002–1003, 2001.
- [20] K. W. Michelle, R. F. Clark, and J. A. Rey, *Pharmacology*, Lippincott Williams & Wilkins, a Wolters Kluwer business, Republic of China, 5th edn edition, 2012.
- [21] S. Ishida, N. Doki, N. Shingai et al., "The clinical features of fatal cyclophosphamide-induced cardiotoxicity in a conditioning regimen for allogeneic hematopoietic stem cell transplantation (allo-HSCT)," *Annals of Hematology*, vol. 95, no. 7, pp. 1145–1150, 2016.
- [22] S. Dhesi, M. P. Chu, G. Blevins et al., "Cyclophosphamide-induced cardiomyopathy: a case report, review, and recommendations for management," *Journal of Investigative Medicine High Impact Case Reports*, vol. 1, no. 1, article 232470961348034, 2013.
- [23] P. Morandi, P. A. Ruffini, G. M. Benvenuto, R. Raimondi, and V. Fossier, "Cardiac toxicity of high-dose chemotherapy," *Bone Marrow Transplantation*, vol. 35, no. 4, pp. 323–334, 2005.
- [24] B. Ranchoux, S. Günther, R. Quarck et al., "Chemotherapy-induced pulmonary hypertension: role of alkylating agents," *The American Journal of Pathology*, vol. 185, no. 2, pp. 356–371, 2015.
- [25] I. Mikaelian, A. Buness, M. C. de Vera-Mudry et al., "Primary endothelial damage is the mechanism of cardiotoxicity of tubulin-binding drugs," *Toxicological Sciences*, vol. 117, no. 1, pp. 144–151, 2010.
- [26] A. Sandoo, G. D. Kitas, and A. R. Carmichael, "Endothelial dysfunction as a determinant of trastuzumab-mediated cardiotoxicity in patients with breast cancer," *Anticancer Research*, vol. 34, no. 3, pp. 1147–1151, 2014.
- [27] M. M. Sayed-Ahmed, M. L. Aldelemy, O. A. Al-Shabanah et al., "Inhibition of gene expression of carnitine palmitoyl-transferase I and heart fatty acid binding protein in cyclophosphamide and ifosfamide-induced acute cardiotoxic rat models," *Cardiovascular Toxicology*, vol. 14, no. 3, pp. 232–242, 2014.
- [28] M. Sayed-Ahmed, A. M. Rishk, S. Solmon, and S. Abdel-Aleem, "Protection by L-carnitine against the inhibition of gene expression of heart fatty acid binding protein by chronic administration of doxorubicin," *Journal of the Egyptian National Cancer Institute*, vol. 12, pp. 275–281, 2000.
- [29] A. H. ElGhandour, M. el Sorady, S. Azab, and M. ElRahman, "Human heart-type fatty acid-binding protein as an early diagnostic marker of doxorubicin cardiac toxicity," *Hematology Reviews*, vol. 1, no. 1, p. 6, 2009.
- [30] A. G. Fatani, A. Q. Darweesh, L. Rizwan, A. M. Aleisa, O. A. Al-Shabanah, and M. M. Sayed-Ahmed, "Carnitine deficiency aggravates cyclophosphamide-induced cardiotoxicity in rats," *Chemotherapy*, vol. 56, no. 1, pp. 71–81, 2010.
- [31] A. Çetinkaya, B. Kantarçeken, E. Bülbüloğlu, and E. Kurutaş, "The effects of L-carnitine on cyclophosphamide-induced oxidative liver and intestinal damage in rats," *Türkiye Klinikleri tıp Bilimleri Dergisi*, vol. 29, no. 5, pp. 1161–1167, 2009.
- [32] E. H. M. Hassanein, A. M. Abo-Youssef, B. A. S. Messiha, and R. A. M. Heme, "Protective effects of montelukast and L-carnitine on cyclophosphamide-induced lung injury," *UK Journal of Pharmaceutical Biosciences*, vol. 3, no. 1, pp. 30–36, 2016.
- [33] M. K. Mahabady, H. N. Varzi, and S. Z. Jahromi, "L-Carnitine protect against cyclophosphamide induced skeletal and neural tube malformations in rat fetuses," *Acta Medica Iranica*, vol. 53, no. 11, pp. 703–710, 2015.
- [34] B. Zhu, Y. F. Zheng, Y. Y. Zhang et al., "Protective effect of L-carnitine in cyclophosphamide-induced germ cell apoptosis," *Journal of Zhejiang University-SCIENCE B*, vol. 16, no. 9, pp. 780–787, 2015.
- [35] I. A. Al-Nasser, "In vivo prevention of cyclophosphamide-induced Ca²⁺ dependent damage of rat heart and liver mitochondria by cyclosporin A," *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, vol. 121, no. 3, pp. 209–214, 1998.
- [36] G. Santulli, W. Xie, S. R. Reiken, and A. R. Marks, "Mitochondrial calcium overload is a key determinant in heart failure," *Proceedings of the National Academy of Sciences*, vol. 112, no. 36, pp. 11389–11394, 2015.
- [37] P. T. Sudharsan, Y. Mythili, E. Selvakumar, and P. Varalakshmi, "Lupeol and its ester exhibit protective role against cyclophosphamide-induced cardiac mitochondrial toxicity," *Journal of Cardiovascular Pharmacology*, vol. 47, no. 2, pp. 205–210, 2006.
- [38] S. Jnaneshwari, M. Hemshekhar, R. M. Thushara et al., "Sesamol ameliorates cyclophosphamide-induced hepatotoxicity by modulating oxidative stress and inflammatory mediators," *Anti-Cancer Agents in Medicinal Chemistry*, vol. 14, no. 7, pp. 975–983, 2014.
- [39] D. F. Mansour, D. O. Saleh, and R. E. Mostafa, "Genistein ameliorates cyclophosphamide - induced hepatotoxicity by modulation of oxidative stress and inflammatory mediators," *Open Access Macedonian Journal of Medical Sciences*, vol. 5, no. 7, pp. 836–843, 2017.
- [40] S. H. Aladaileh, M. H. Abukhalil, S. A. M. Saghir et al., "Galanin activates Nrf2 signaling and attenuates oxidative damage, inflammation, and apoptosis in a rat model of cyclophosphamide-induced hepatotoxicity," *Biomolecules*, vol. 9, no. 8, p. 346, 2019.
- [41] A. M. Mahmoud, M. O. Germoush, M. F. Alotaibi, and O. E. Hussein, "Possible involvement of Nrf2 and PPAR γ up-regulation in the protective effect of umbelliferone against cyclophosphamide-induced hepatotoxicity," *Biomedicine & Pharmacotherapy*, vol. 86, pp. 297–306, 2017.
- [42] P. R. Gore, C. P. Prajapati, U. B. Mahajan et al., "Protective effect of thymoquinone against cyclophosphamide-induced hemorrhagic cystitis through inhibiting DNA damage and upregulation of Nrf2 expression," *International Journal of Biological Sciences*, vol. 12, no. 8, pp. 944–953, 2016.
- [43] D. N. Tripathi and G. B. Jena, "Effect of melatonin on the expression of Nrf2 and NF- κ B during cyclophosphamide-induced urinary bladder injury in rat," *Journal of Pineal Research*, vol. 48, no. 4, pp. 324–331, 2010.
- [44] A. A. El-Sheikh, M. A. Morsy, and A. M. Okasha, "Inhibition of NF- κ B/TNF- α pathway may be involved in the protective effect of resveratrol against cyclophosphamide-induced multi-organ toxicity," *Immunopharmacology and Immunotoxicology*, vol. 39, no. 4, pp. 180–187, 2017.

- [45] S. Nafees, S. Rashid, N. Ali, S. K. Hasan, and S. Sultana, "Rutin ameliorates cyclophosphamide induced oxidative stress and inflammation in Wistar rats: role of NF κ B/MAPK pathway," *Chemico-Biological Interactions*, vol. 231, pp. 98–107, 2015.
- [46] A. Morita, S. Yamamoto, B. Wang et al., "Sodium orthovanadate inhibits p53-mediated apoptosis," *Cancer Research*, vol. 70, no. 1, pp. 257–265, 2010.
- [47] B. Saleme, V. Gurtu, Y. Zhang et al., "Tissue-specific regulation of p53 by PKM2 is redox dependent and provides a therapeutic target for anthracycline-induced cardiotoxicity," *Science Translational Medicine*, vol. 11, no. 478, article eaau8866, 2019.
- [48] A. V. Gudkov and E. A. Komarova, "Prospective therapeutic applications of p53 inhibitors," *Biochemical and Biophysical Research Communications*, vol. 331, no. 3, pp. 726–736, 2005.
- [49] Y. A. Asiri, "Probulcol attenuates cyclophosphamide-induced oxidative apoptosis, p53 and Bax signal expression in rat cardiac tissues," *Oxidative Medicine and Cellular Longevity*, vol. 3, no. 5, pp. 308–316, 2010.
- [50] A. A. El-Kholy, M. A. Elkablawy, and D. S. El-Agamy, "Lutein mitigates cyclophosphamide induced lung and liver injury via NF- κ B/MAPK dependent mechanism," *Biomedicine & Pharmacotherapy*, vol. 92, pp. 519–527, 2017.
- [51] R. Gómez-Sintes and J. J. Lucas, "NFAT/Fas signaling mediates the neuronal apoptosis and motor side effects of GSK-3 inhibition in a mouse model of lithium therapy," *The Journal of Clinical Investigation*, vol. 120, no. 7, pp. 2432–2445, 2010.
- [52] N. M. Albert, "Improving medication adherence in chronic cardiovascular disease," *Critical Care Nurse*, vol. 28, no. 5, pp. 54–64, 2008.
- [53] J. Perk, Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR), G. De Backer et al., "European guidelines on cardiovascular disease prevention in clinical practice (version 2012)," *International Journal of Behavioral Medicine*, vol. 19, no. 4, pp. 403–488, 2012.
- [54] G. Van Camp, "Cardiovascular disease prevention," *Acta Clinica Belgica*, vol. 69, no. 6, pp. 407–411, 2014.
- [55] X. Rossello, S. J. Pocock, and D. G. Julian, "Long-term use of cardiovascular drugs: challenges for research and for patient care," *Journal of the American College of Cardiology*, vol. 66, no. 11, pp. 1273–1285, 2015.
- [56] S. A. Virani, S. Dent, C. Brezden-Masley et al., "Canadian Cardiovascular Society guidelines for evaluation and management of cardiovascular complications of cancer therapy," *Canadian Journal of Cardiology*, vol. 32, no. 7, pp. 831–841, 2016.
- [57] H. Nakamae, K. Tsumura, Y. Terada et al., "Notable effects of angiotensin II receptor blocker, valsartan, on acute cardiotoxic changes after standard chemotherapy with cyclophosphamide, doxorubicin, vincristine, and prednisolone," *Cancer*, vol. 104, no. 11, pp. 2492–2498, 2005.
- [58] G. Curigliano, D. Cardinale, S. Dent et al., "Cardiotoxicity of anticancer treatments: epidemiology, detection, and management," *CA: A Cancer Journal for Clinicians*, vol. 66, no. 4, pp. 309–325, 2016.
- [59] M. Koneru, N. Nalban, B. D. Sahu, and R. Sistla, "Natural products against drug-induced cardiotoxicity," *Cardioprotective Natural Products: Promises And Hopes*, G. Brahmachari, Ed., pp. 121–147, 2017.
- [60] M. Rafieian-Kopaei, "Medicinal plants and the human needs," *Journal of Herbmmed Pharmacology*, vol. 1, no. 1, pp. 1–2, 2012.
- [61] H. Avci, E. T. Epikmen, E. Ipek et al., "Protective effects of silymarin and curcumin on cyclophosphamide-induced cardiotoxicity," *Experimental and Toxicologic Pathology*, vol. 69, no. 5, pp. 317–327, 2017.
- [62] M. A. Ayza, R. Balasubramanian, and A. H. Berhe, "Cardioprotective effect of Croton macrostachyus stem bark extract and solvent fractions on cyclophosphamide-induced cardiotoxicity in rats," *Evidence-Based Complementary and Alternative Medicine*, vol. 2020, 13 pages, 2020.
- [63] S. Baniya, D. Dhananjaya, A. Acharya, B. Dangi, and A. Sapkota, "Cardioprotective activity of ethanolic extract of Citrus grandis (L.) Osbeck peel on doxorubicin and cyclophosphamide induced cardiotoxicity in albino rats," *International Journal of Pharmaceutical Sciences and Drug Research*, vol. 7, pp. 354–360, 2015.
- [64] L. Bhatt, B. Sebastian, and V. Joshi, "Mangiferin protects rat myocardial tissue against cyclophosphamide induced cardiotoxicity," *Journal of Ayurveda and Integrative Medicine*, vol. 8, no. 2, pp. 62–67, 2017.
- [65] S. K. Bjelogrić, J. Radic, V. Jovic, and S. Radulovic, "Activity of d,l-alpha-Tocopherol (vitamin e) against cardiotoxicity induced by doxorubicin and doxorubicin with cyclophosphamide in mice," *Basic Clinical Pharmacology Toxicology*, vol. 97, no. 5, pp. 311–319, 2005.
- [66] S. Cetik, A. Ayhanci, and V. Sahinturk, "Protective effect of carvacrol against oxidative stress and heart injury in cyclophosphamide-induced cardiotoxicity in rat," *Brazilian Archives of Biology and Technology*, vol. 58, no. 4, pp. 569–576, 2015.
- [67] M. Chakraborty, J. V. Kamath, and A. Bhattacharjee, "Pharmacodynamic interaction of green tea extract with hydrochlorothiazide against cyclophosphamide-induced myocardial damage," *Toxicology International*, vol. 21, no. 2, pp. 196–202, 2014.
- [68] M. Chakraborty, A. Bhattacharjee, and J. V. Kamath, "Cardioprotective effect of curcumin and piperine combination against cyclophosphamide-induced cardiotoxicity," *Indian Journal of Pharmacology*, vol. 49, no. 1, pp. 65–70, 2017.
- [69] D. J. Conklin, P. Haberzettl, G. Jagatheesan et al., "Glutathione S-transferase P protects against cyclophosphamide-induced cardiotoxicity in mice," *Toxicology and Applied Pharmacology*, vol. 285, no. 2, pp. 136–148, 2015.
- [70] D. S. El-Agamy, M. A. Elkablawy, and H. M. Abo-Haded, "Modulation of cyclophosphamide-induced cardiotoxicity by methyl palmitate," *Cancer Chemotherapy and Pharmacology*, vol. 79, no. 2, pp. 399–409, 2017.
- [71] A. Gado, A. Adam, and B. Aldahmash, "Cardiotoxicity induced by cyclophosphamide in rats: protective effect of curcumin," *Journal of Research in Environmental Science and Toxicology*, vol. 2, no. 4, pp. 87–95, 2013.
- [72] S. Gunes, V. Sahinturk, P. Karasati, I. K. Sahin, and A. Ayhanci, "Cardioprotective effect of selenium against cyclophosphamide-induced cardiotoxicity in rats," *Biological Trace Element Research*, vol. 177, no. 1, pp. 107–114, 2017.
- [73] A. Iqbal, S. Sharma, M. A. Ansari et al., "Nerolidol attenuates cyclophosphamide-induced cardiac inflammation, apoptosis and fibrosis in Swiss Albino mice," *European Journal of Pharmacology*, vol. 863, p. 172666, 2019.
- [74] H. H. Mansour, S. M. el kiki, and H. F. Hasan, "Protective effect of N-acetylcysteine on cyclophosphamide-induced cardiotoxicity in rats," *Environmental Toxicology and Pharmacology*, vol. 40, no. 2, pp. 417–422, 2015.

- [75] Y. Mythili, P. Sudharsan, E. Selvakumar, and P. Varalakshmi, "Protective effect of DL- α -lipoic acid on cyclophosphamide induced oxidative cardiac injury," *Chemico-Biological Interactions*, vol. 151, no. 1, pp. 13–19, 2004.
- [76] Y. Mythili, P. T. Sudharsan, V. Sudhahar, and P. Varalakshmi, "Protective effect of DL- α -lipoic acid on cyclophosphamide induced hyperlipidemic cardiomyopathy," *European Journal of Pharmacology*, vol. 543, no. 1-3, pp. 92–96, 2006.
- [77] Y. Mythili, P. T. Sudharsan, G. Amudha, and P. Varalakshmi, "Effect of dl- α -lipoic acid on cyclophosphamide induced lysosomal changes in oxidative cardiotoxicity," *Life Sciences*, vol. 80, no. 21, pp. 1993–1998, 2007.
- [78] M. N. Nagi, O. A. Al-Shabanah, M. M. Hafez, and M. M. Sayed-Ahmed, "Thymoquinone supplementation attenuates cyclophosphamide-induced cardiotoxicity in rats," *Journal of Biochemical and Molecular Toxicology*, vol. 25, no. 3, pp. 135–142, 2011.
- [79] J. G. Omole, O. A. Ayoka, Q. K. Alabi et al., "Protective effect of kolaviron on cyclophosphamide-induced cardiac toxicity in rats," *Journal of Evidence-Based Integrative Medicine*, vol. 23, p. 215658721875764, 2018.
- [80] V. Sekeroğlu, B. Aydin, and Z. Sekeroğlu, "Viscum album L. extract and quercetin reduce cyclophosphamide-induced cardiotoxicity, urotoxicity and genotoxicity in mice," *Asian Pacific Journal of Cancer Prevention*, vol. 12, no. 11, pp. 2925–2931, 2011.
- [81] S. Senthilkumar, T. Devaki, B. M. Manohar, and M. S. Babu, "Effect of squalene on cyclophosphamide-induced toxicity," *Clinica Chimica Acta*, vol. 364, no. 1-2, pp. 335–342, 2006.
- [82] M. M. S. Nashwa, R. H. M. Abeer, E. S. M. A. Nagah, E. I. Nahla, and H. A. H. M. Noura, "The possible protective effect of Zingiber officinale extract on cyclophosphamide-induced cardiotoxicity in adult male albino rats," *Journal of Toxicology and Environmental Health Sciences*, vol. 11, no. 4, pp. 38–49, 2019.
- [83] T. Shanmugarajan, M. Arunsundar, I. Somasundaram, E. Krishnakumar, D. Sivaraman, and V. Ravichandiran, "Cardioprotective effect of Ficus hispida Linn. on cyclophosphamide provoked oxidative myocardial injury in a rat model," *International Journal of Pharmacology*, vol. 4, no. 2, pp. 1–10, 2008.
- [84] Y. Song, C. Zhang, C. Wang et al., "Ferulic acid against cyclophosphamide-induced heart toxicity in mice by inhibiting NF- κ B pathway," *Evidence-Based Complementary and Alternative Medicine*, vol. 2016, 8 pages, 2016.
- [85] P. T. Sudharsan, Y. Mythili, V. Sudhahar, and P. Varalakshmi, "Role of lupeol and its ester on cyclophosphamide-induced hyperlipidaemic cardiomyopathy in rats," *Journal of Pharmacy and Pharmacology*, vol. 57, no. 11, pp. 1437–1444, 2005.
- [86] A. H. M. V. Swamy, U. M. Patel, B. C. Koti, P. C. Gadad, N. L. Patel, and A. H. M. Thippeswamy, "Cardioprotective effect of Saraca indica against cyclophosphamide induced cardiotoxicity in rats: a biochemical, electrocardiographic and histopathological study," *Indian Journal of Pharmacology*, vol. 45, no. 1, pp. 44–48, 2013.
- [87] V. Todorova, D. Vanderpool, S. Blossom et al., "Oral glutamine protects against cyclophosphamide-induced cardiotoxicity in experimental rats through increase of cardiac glutathione," *Nutrition*, vol. 25, no. 7-8, pp. 812–817, 2009.
- [88] A. A. El-Sheikh, W. Y. Abdelzاهر, A. A. Gad, and S. A. Abdel-Gaber, "Purine versus non-purine xanthine oxidase inhibitors against cyclophosphamide-induced cardiac and bone marrow toxicity in rats," *Human & Experimental Toxicology*, vol. 39, no. 3, pp. 249–261, 2020.
- [89] M. M. M. Refaie, S. Shehata, M. el-Hussieny, W. M. Abdelraheem, and A. M. A. Bayoumi, "Role of ATP-sensitive potassium channel (KATP) and eNOS in mediating the protective effect of nicorandil in cyclophosphamide-induced cardiotoxicity," *Cardiovascular Toxicology*, vol. 20, no. 1, pp. 71–81, 2020.
- [90] V. Sindhi, V. Gupta, K. Sharma, S. Bhatnagar, R. Kumari, and N. Dhaka, "Potential applications of antioxidants – A review," *Journal of Pharmacy Research*, vol. 7, no. 9, pp. 828–835, 2013.
- [91] E. B. Kurutas, "The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state," *Nutrition Journal*, vol. 15, no. 1, p. 71, 2016.
- [92] T. L. Zern and M. L. Fernandez, "Cardioprotective effects of dietary polyphenols," *The Journal of Nutrition*, vol. 135, no. 10, pp. 2291–2294, 2005.
- [93] J.-C. Stoclet, T. Chataigneau, M. Ndiaye et al., "Vascular protection by dietary polyphenols," *European Journal of Pharmacology*, vol. 500, no. 1-3, pp. 299–313, 2004.
- [94] K. D. Croft, "Dietary polyphenols: antioxidants or not?," *Archives of biochemistry and biophysics*, vol. 595, pp. 120–124, 2016.
- [95] M. Sandu, L. M. Birsă, and L. G. Bahrin, "Flavonoids—small molecules, high hopes," *Acta Chemica Iasi*, vol. 25, no. 1, pp. 6–23, 2017.
- [96] L. Shen and H.-F. Ji, "Is antioxidant supplement beneficial? New avenue to explore," *Trends in Food Science & Technology*, vol. 68, pp. 51–55, 2017.
- [97] E. Guallar, S. Stranges, C. Mulrow, L. J. Appel, and E. R. Miller, "Enough is enough: stop wasting money on vitamin and mineral supplements," *Annals of Internal Medicine*, vol. 159, no. 12, pp. 850–851, 2013.
- [98] J. A. Leopold, "Antioxidants and coronary artery disease: from pathophysiology to preventive therapy," *Coronary Artery Disease*, vol. 26, no. 2, pp. 176–183, 2015.
- [99] B. Halliwell, "Dietary polyphenols: good, bad, or indifferent for your health?," *Cardiovascular Research*, vol. 73, no. 2, pp. 341–347, 2007.