Research Article

The Effects of Hypoxia-Reoxygenation in Mouse Digital Flexor Tendon-Derived Cells

Chen Chen ¹, Wei Feng Mao ^{1,2} and Ya Fang Wu ¹

¹Department of Hand Surgery, Affiliated Hospital of Nantong University, Nantong, Jiangsu, China

²Department of Anatomy, Medical School, Nantong University, Nantong, Jiangsu, China

Correspondence should be addressed to Wei Feng Mao; maoweifeng@ntu.edu.cn and Ya Fang Wu; yafang.wu@ntu.edu.cn

Received 27 July 2020; Revised 12 November 2020; Accepted 25 November 2020; Published 15 December 2020

Academic Editor: Mariella B. Freitas

Copyright © 2020 Chen Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. Ischemia-reperfusion injury refers to the exacerbated and irreversible tissue damage caused by blood flow restoration after a period of ischemia. The hypoxia-reoxygenation (H/R) model in vitro is ideal for studying ischemia-reperfusion injury at the cellular level. We employed this model and investigated the effects of cobalt chloride- (CoCl₂-) induced H/R in cells derived from mouse digital flexor tendons. **Materials and Methods.** Various H/R conditions were simulated via treatment of tendon-derived cells with different concentrations of CoCl₂ for 24 h, followed by removal of CoCl₂ to restore a normal oxygen state for up to 96 h. Cell viability was measured using the Cell Counting Kit-8 (CCK-8) assay. Cell growth was determined via observation of cell morphology and proliferation. Oxidative stress markers and mitochondrial activity were detected. The expression levels of hypoxia-inducible factor- (HIF-) 1 α , vascular endothelial growth factor-A (VEGF-A), collagen I, and collagen III were determined using Western blot (WB), real-time PCR, and immunofluorescence staining. Cellular apoptosis was analyzed via flow cytometry, and the expression of apoptosis-related proteins Bax and bcl-2 was examined using WB. **Results.** The cells treated with low concentrations of CoCl₂ showed significantly increased cell viability after reoxygenation. The increase in cell viability was even more pronounced in cells that had been treated with high concentrations of CoCl₂. Under H/R conditions, cell morphology and growth were unchanged, while oxidative stress reaction was induced and mitochondrial activity was increased. H/R exerted opposite effects on the expression of HIF-1 α mRNA and protein. Meanwhile, the expression of VEGF-A was upregulated, whereas collagen type I and type III were significantly downregulated. The level of cellular apoptosis did not show significant changes during H/R, despite the significantly increased Bax protein and reduced bcl-2 protein levels that led to an increase in the Bax/bcl-2 ratio during reoxygenation. **Conclusions.** Tendon-derived cells were highly tolerant to the hypoxic environments induced by CoCl₂. Reoxygenation after hypoxia preconditioning promoted cell viability, especially in cells treated with high concentrations of CoCl₂. H/R conditions caused oxidative stress responses but did not affect cell growth. The H/R process had a notable impact on collagen production and expression of apoptosis-related proteins by tendon-derived cells, while the level of cellular apoptosis remained unchanged.

1. Introduction

Tendon injury is frequently accompanied by a lack of blood supply, exposing the tendon to a hypoxic condition. Oxygen tension, which is influenced by blood supply, is vital to cell survival and proliferation. In vitro studies have shown that low oxygen tension enhances the expansion capacity of newborn pig tenocytes without affecting the cellular phenotype and functions [1], and hypoxic conditions significantly improve the cell proliferation and self-renewal capacity of

human tendon stem cells [2, 3]. Other studies, however, have shown that hypoxia inhibits cell proliferation and alters the synthesis of matrix components in synovial tissue [4]. Cobalt chloride- (CoCl₂-) induced hypoxia has been shown to alter the expression of the bcl-2 family proteins and trigger caspase cascade apoptosis [5, 6]. Collectively, the conflicting results in these studies underscore the complicated effects of hypoxia on cell growth.

Most tissues subjected to oxygen deprivation after injury undergo reperfusion [7–10]. Reperfusion injury, which is

manifested by blood flow-deprived and oxygen-starved organs following blood flow restoration and tissue reoxygenation, is a tissue response to stimulate oxidative metabolism [11–15]. Tissues with reperfusion injury share some of the characteristic features of injury responses to oxygen deprivation and subsequent reoxygenation, such as necrosis, apoptosis, impaired microvascular function, and edema. It has been shown that considerable vascular remodeling takes place during tendon healing to enhance perfusion [16, 17]. Stange et al. reported a peak in changes of relative blood volume 7 days after *in situ* freezing-induced tendinopathy in a rat patellar tendon [18]. A significant uniform hypervascularization occurred during the healing process of chronic Achilles tendinopathy after operative tenolysis in rabbits [17]. The increased vascularization facilitates reoxygenation during tendon repair. Therefore, the process of ischemia/reperfusion after tendon injury, which causes hypoxia-reoxygenation (H/R) changes at the cellular level, presents a major impediment for tendon recovery [19]. However, the sensitivity of tendon-derived cells to hypoxia and the effect of H/R on cell growth and characteristics have been rarely addressed.

In this study, we aimed to establish an *in vitro* model of cobalt chloride- (CoCl_2 -) induced hypoxia-reoxygenation (H/R) in mouse digital flexor tendon-derived cells and investigate the effects of H/R on these cells [20]. Hypoxic conditions were simulated by treating cultured cells with CoCl_2 , whereas reoxygenation conditions were simulated by replacing the culture media following CoCl_2 treatment. The principle behind CoCl_2 -induced hypoxia is the replacement of Fe^{2+} by Co^{2+} in heme-based oxygen sensors, which prevents the oxygen sensor from combining with oxygen [21]. Compared with culturing cells in a low-oxygen incubator, the mimic of the hypoxic niche chemically exhibits several advantages, including simple preparation of CoCl_2 , flexible dosage adjustment, treatment suspension at any time, and no impact on oxygen levels for cells that require normal oxygen tension [22]. To have better control over the degree of hypoxia, we decided to employ different CoCl_2 concentrations. Following H/R, cell viability, cell growth, oxidative stress markers, mitochondrial activity, and expression of HIF-1 α , vascular endothelial growth factor-A (VEGF-A), collagen I, and collagen III were assessed. Cellular apoptosis and expression of apoptosis-related protein markers Bax and bcl-2 were analyzed.

2. Materials and Methods

2.1. Experiment Design. We first confirmed the success of the CoCl_2 -induced hypoxia model by examining the expression of HIF-1 α protein in tendon-derived cells treated with low-to-high concentrations of CoCl_2 (0.05, 0.1, 0.2, and 0.4 mM), according to previous studies [20, 23, 24]. Next, the cells were divided into the following groups: control (cells cultured in regular medium), H (cells treated with a concentration gradient of CoCl_2 for 24 h), and R (cells treated with CoCl_2 for 24 h, followed by culturing in regular culture medium for 24 h, 48 h, 72 h, and 96 h (R24h, R48h, R72h, and R96h)). Cell viability was compared across the control, H, and R groups, as well as among different concentrations

of CoCl_2 . Then, we determined an appropriate concentration of CoCl_2 for the hypoxia model to perform the following experiments: cell growth, oxidative stress markers, mitochondrial activity, mRNA and protein expression of HIF-1 α , VEGF-A, collagen I, and collagen III, cellular apoptosis, and expression of the apoptosis-related proteins Bax and bcl-2.

2.2. Isolation of Mouse Flexor Tendon Cells and H/R Model.

The animal procedures were approved by the Administration Committee of Experimental Animals of our university. All animal experiments were carried out in accordance with the Experimental Animal Management Ordinance (National Science and Technology Committee of China) and the Guide for the Care and Use of Laboratory Animals (National Institutes of Health (NIH), Bethesda, MD, USA). Tendon-derived cells were isolated from mouse flexor digitorum profundus (FDP) tendons (C57BL/6, female, 4 weeks old). The mice were sacrificed by cervical dislocation, and the FDP tendons of the index, middle, and ring fingers of the hind paws were harvested. The tendon samples were washed with sterile phosphate-buffered saline (PBS) and then cut into small pieces. The cut samples were digested in a mixture of 3 mg/ml type I collagenase (Gibco, Thermo Fisher Scientific, Grand Island, NY) and 4 mg/ml dispase (Gibco, Thermo Fisher Scientific) for 2 h at 37°C. Undigested tissues and debris were filtered through a 70 μm cell strainer (Merck Millipore, Cork, Ireland). The released cells in the filtrate were centrifuged at 600 g for 10 min and resuspended in low-glucose Dulbecco's modified Eagle's medium (LG-DMEM; Gibco, Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS), 100 U/ml penicillin, and 100 $\mu\text{g}/\text{ml}$ streptomycin. The isolated nucleated cells were plated as passage 0 (P0) at a seeding density of 500 cells/ cm^2 , as determined in a previous study [25]. The cells were cultured at 37°C with 5% CO_2 . After 2 days, the cells were washed twice with PBS to remove nonadherent cells. After 10 days, the cells at 95% confluence were collected via local trypsin digestion and split 1:3 for subsequent passages. The medium was changed twice a week. Cells at P3 were used for all the experiments.

2.3. Cell Viability. The cells were seeded in 96-well plates at a density of 5000 cells/well and cultured in DMEM. After 48 h, different concentrations of CoCl_2 (0.05, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, and 0.4 mM) were added to the cells for 24 h. Subsequently, the cell media were replaced with regular DMEM, and the cells were cultured for 24 h, 48 h, 72 h, and 96 h. Cell Counting Kit-8 (CCK-8; meilunbio, China) was used to evaluate the viability of cells subjected to H/R. Briefly, fresh medium containing CCK-8 solution was added to the 96-well plates containing cells cultured for different time periods. The cells were then incubated for 1.5 h at 37°C. The absorbance of the supernatants (100 μl) was measured at 450 nm using an automatic microplate reader (Bio-Rad).

2.4. Cell Growth. The cells were seeded in a 6-well plate at 5000 cells/ cm^2 and in a 96-well plate at 5000 cells/well. After 48 h, cells were treated with CoCl_2 (concentration determined by

above cell viability measurement) for 24 h, followed by reoxygenation for 24 h, 48 h, 72 h, and 96 h. The morphological changes of cells in the 6-well plate before and after H/R treatment were observed under a phase-contrast microscope (Olympus IX71). The cells in the 96-well plate were harvested using a 0.05% trypsin (Invitrogen) solution, and the cell numbers were counted for growth curve construction before and after H/R treatment.

2.5. Oxidative Stress Markers. The intracellular malondialdehyde (MDA) levels were measured using a Lipid Peroxidation MDA Assay Kit (Beyotime, Shanghai, China) following the manufacturer's instructions. First, cells were seeded in 6-well plates (5×10^4 cells/well) and continuously grew until the confluence reached 90%. After H/R treatment, cells were collected and lysed by cell lysis buffer (Beyotime, Shanghai, China) and centrifuged at 10,000 g for 15 min. The supernatants were reacted with the thiobarbituric acid (TBA), and the reaction products were measured spectrophotometrically at 532 nm.

A superoxide dismutase (SOD) assay was performed using the Total Superoxide Dismutase Assay Kit with WST-8 (Beyotime, Shanghai, China). Cells were seeded in 6-well plates (5×10^4 cells/well) and continuously grew until the confluence reached 90%. The cells were treated with H/R conditions₂. Cells were washed with cold PBS before the addition of SOD buffer solution. Protein concentration was measured using the Enhanced BCA Protein Assay Kit (Beyotime, Shanghai, China). The samples (20 μ l) were mixed with the WST-8/enzyme solution (160 μ l) and the reaction starting solution (20 μ l) and incubated at 37°C for 30 minutes. The absorbance at 450 nm was measured using a microplate reader (Bio-Rad).

2.6. MitoTracker Red Staining. Cells were stained with MitoSOX Red Mitochondrial Superoxide Indicator (Yeasen, Shanghai, China) and Hoechst 33342 (Cell Signaling Technologies, Danvers, MA). Cells at 80-90% confluence were treated with H/R conditions₂. Then, the cells were incubated with MitoSOX Red Mitochondrial Superoxide Indicator for 10 min at 37°C. Then, the cells were washed with PBS and stained with Hoechst 33342 labeling solution (1:10000) for 10 min at room temperature. The cells were observed under a fluorescence microscope (Olympus IX71).

2.7. Western Blot Analysis. The cells were rinsed twice with cold PBS and then lysed in RIPA buffer (Beyotime, Shanghai, China) containing 1% (v/v) phenylmethanesulfonyl fluoride (PMSF) (Beyotime, Shanghai, China) and 1% (v/v) protease and phosphatase inhibitor cocktails (Roche, Mannheim, Germany). The Enhanced BCA Protein Assay Kit (Beyotime, Shanghai, China) was used to measure protein concentrations. The samples were subjected to 10% SDS-polyacrylamide gel electrophoresis and subsequently transferred onto a polyvinylidene difluoride membrane (Millipore Corp., Billerica, Mass.) under 100 volts for 2 h. The membranes were incubated in Tris-buffered saline (TBS: 10 mM Tris-HCl, pH 7.4, 150 mM NaCl) and 0.1% (v/v) Tween 20 (TBS-Tween) containing 5% (w/v) dried milk

FIGURE 1: Protein expression of HIF-1 α after treatment with different concentrations of CoCl₂ for 24 h. * $P < 0.05$ vs. control group; ** $P < 0.005$ vs. control group.

for 1 h and then incubated with the respective primary antibody overnight at 4°C. The primary antibodies against HIF-1 α , VEGF-A, collagen I, collagen III, Bax, and bcl-2 are described in Table S1. Next, the filters were washed 3 times, 8 min each, with TBS-T and then incubated with a conjugated affinity-purified secondary antibody labeled with IRDye 800v for 1 h at room temperature. Afterwards, the membranes were washed again, and the protein bands were detected with an Odyssey imager (LI-COR, Inc., Lincoln, NE). The intensity of each band was quantified with ImageJ software (NIH, Bethesda, MD, USA). The expression levels of proteins were normalized to β -actin.

2.8. Real-Time PCR. Total RNA of cells was isolated using TRIzol® Reagent (Ambion, Life Technologies). First-strand cDNA were reverse-transcribed with HiScript II qRT Super-Mix plus gDNA wiper (R223-01, Vazyme, China). QRT-PCR was carried out using 1x AceQ qPCR SYBR Green Master Mix (Q111-02, Vazyme, China) according to the manufacturer's instructions. Specific primers were synthesized for the HIF-1 α , VEGF-A, collagen I, and collagen III mRNAs. GAPDH was used as an internal reference, and the geometric mean of its expression was used for normalization. The sequences of qPCR primers are listed in Table S2. Relative quantification of the target genes was performed in triplicate and analyzed using the $2^{-\Delta\Delta C_t}$ method.

2.9. Immunocytochemistry. The cells were plated at a density of 1×10^4 cells/well on coverglasses in 24-well plates for 2 days. The expression of 4 protein markers, including HIF-1 α , VEGF-A, collagen I, and collagen III, was measured using immunocytochemistry. Briefly, cells were fixed in 4% paraformaldehyde for 30 min at room temperature. Following washing, the cells were permeabilized with 0.25% Triton X-100 in PBS and blocked with 1% bovine serum albumin (BSA) for 1 h at room temperature. The cells were then incubated with antibodies against HIF-1 α , VEGF-A,

FIGURE 2: Changes in cell viability of tendon-derived cells under H/R conditions induced by a concentration gradient of CoCl₂. (a) Cell viability under CoCl₂-induced hypoxia. (b–e) Cell viability after reoxygenation for 24 h, 48 h, 72 h, and 96 h. * $P < 0.05$ vs. control group; ** $P < 0.005$ vs. control group.

FIGURE 3: Changes in cell viability of tendon-derived cells under H/R at each concentration of CoCl₂. (a–h) Cell viability increases along the CoCl₂ concentration gradient (0.05–0.4 mM). * $P < 0.05$ vs. control group; ** $P < 0.005$ vs. control group; # $P < 0.05$ vs. H group; ## $P < 0.005$ vs. H group.

collagen I, and collagen III overnight at 4°C. After 3 washes with PBS, the cells were incubated with Alexa Fluor488 Goat anti-rabbit IgG (H+L) for 2 h at room temperature. Finally, cells were counterstained with Hoechst. The stained cells were then examined under a fluorescence microscope (Olympus IX71).

2.10. Flow Cytometry Analysis. Annexin V labeling was used in conjunction with flow cytometry to detect phosphatidylserine on the outer membrane of apoptotic cells. First, the binding buffer was diluted to 1: 4 with deionized water (4 ml binding buffer+12 ml deionized water). Then, the cells were harvested using 0.05% trypsin with no EDTA solution

FIGURE 4: Cell morphology and growth curves of tendon-derived cells when treated with 0.3 mM CoCl_2 . (a) Cell morphology under H/R conditions. Scale bar: 200 μm . (b) Cell growth curves under H/R conditions.

(Invitrogen), washed twice with 4°C prechilled PBS, and resuspended with binding buffer. Next, the cell concentration was adjusted to 1×10^6 cells / ml. The cell suspension (100 μl) was placed in a 5 ml flow tube, and 5 μl annexin V/Alexa Fluor 488 and 10 μl propidium iodide solution were added. After mixing, the mixture was incubated away from light for 15 min at room temperature. After incubation, 400 μl of PBS was added to the reaction tube, and the cells were analyzed using flow cytometry (Attune NxT, Applied Biosystems, CA, USA). For each sample, approximately 10,000 events were counted and analyzed using the FlowJo software (FlowJo, Ashland, OR, USA).

2.11. Statistical Analysis. All experiments were repeated at least 3 times for replication. The data were expressed as mean $\pm$ SD. The results were analyzed by one-way ANOVA with post hoc Tukey testing or by the *t*-test for pairwise comparisons. All data were analyzed using Prism 5.0b (GraphPad

Software, La Jolla, CA, USA). $P < 0.05$ was considered statistically significant.

3. Results

3.1. CoCl_2 -Induced Hypoxia Elevates HIF-1 α Expression. The protein expression of HIF-1 α significantly increased in cells treated with CoCl_2 at concentrations of 0.05 and 0.1 mM ($P < 0.05$) and further increased with higher CoCl_2 concentrations of 0.2 and 0.4 mM ($P < 0.005$) (Figure 1), which confirmed the success of the CoCl_2 -induced hypoxia model in tendon-derived cells.

3.2. H/R Increases Cell Viability. Under hypoxic conditions induced by different concentrations of CoCl_2 , the viability of tendon-derived cells showed an initial increase followed by a gradual decrease with the increased CoCl_2 concentrations. A significant difference was observed only at 0.1 mM of CoCl_2 when compared with the control group ($P < 0.05$)

FIGURE 5: (a) Cellular malondialdehyde (MDA) levels. (b) Superoxide dismutase (SOD) activity. (c) The intensity of mitochondria in different groups. Scale bar: 100 μ m. * P < 0.05 vs. control group; ** P < 0.005 vs. control group; # P < 0.05 vs. H group; ## P < 0.005 vs. H group.

(Figure 2(a)). After reoxygenation for 24 h (R24h), cell viability significantly increased in cells treated with 0.15 mM and higher CoCl_2 concentrations (P < 0.05) (Figure 2(b)). At R48h, cell viability significantly increased in cells treated with 0.2–0.4 mM CoCl_2 (P < 0.05 or P < 0.005) (Figure 2(c)). At R72h, a significant increase in cell viability was only observed in cells treated with 0.25 mM of CoCl_2 (P < 0.05) (Figure 2(d)). The cell viability of all CoCl_2 -treated cells was similar to that in the control group at R96h (Figure 2(e)).

At each concentration, cell viability was compared across the control, hypoxia, and reoxygenation groups. Compared with the control group, the cells treated with low concentrations of CoCl_2 showed significant increases in cell viability under hypoxia, with a 14% increase at 0.05 mM and 15% at 0.1 mM (P < 0.05) (Figures 3(a) and 3(b)). The cells treated with higher concentrations of CoCl_2 at 0.15, 0.2, 0.25, and 0.3 mM exhibited significantly enhanced cell viability after reoxygenation for 24 h–72 h (P < 0.05 or P < 0.005) (Figures 3(c)–3(f)). At the 2 highest CoCl_2 concentrations, cell viability significantly increased at R48h compared with

those in the control and H groups (P < 0.005) (Figures 3(g) and 3(h)).

3.3. H/R Does Not Inhibit Cell Growth. During the experiment, we observed relatively healthy growth of cells treated with 0.3 mM CoCl_2 after reoxygenation. The cells were shrunk slightly when under hypoxic conditions, but the morphology returned to normal after reoxygenation (Figure 4(a)). Therefore, we selected 0.3 mM CoCl_2 to simulate hypoxia for subsequent experiments. Cell growth curves revealed a slightly increased cell proliferation under H/R compared with that of the control group (Figure 4(b)).

3.4. H/R Mediates Oxidative Stress and Mitochondrial Activity. To determine the impact of H/R on oxidative stress, the intracellular levels of MDA and SOD were measured. The MDA levels significantly increased during reoxygenation compared with those in the control and H groups (P < 0.005) (Figure 5(a)). The SOD levels significantly decreased under H/R compared with that in the control group

FIGURE 6: Protein expression of (a) HIF-1 α , (b) VEGF-A, (c) collagen I, and (d) collagen III when cells were under H/R conditions at 0.3 mM CoCl₂. * $P < 0.05$ vs. control group; ** $P < 0.005$ vs. control group; # $P < 0.05$ vs. H group; ## $P < 0.005$ vs. H group.

($P < 0.005$) (Figure 5(b)). The MitoTracker Red staining showed that the intensity of bioactive mitochondria was higher in the H and R groups than in the control group (Figure 5(c)).

3.5. H/R Enhances Protein Expression of HIF-1 α and VEGF-A and Decreases Expression of Collagens I and III. The protein expression of HIF-1 α significantly increased when the cells were under hypoxia and returned to normal levels after reoxygenation ($P < 0.05$) (Figure 6(a)). By comparison, VEGF-A levels showed a similar but not significant increase under hypoxia and at R24h and returned to normal from R48h and after (Figure 6(b)). Both collagen I and collagen III protein levels significantly decreased at R48h, R72h, and R96h compared with those of the control group and H group ($P < 0.05$ or $P < 0.005$) (Figures 6(c) and 6(d)).

3.6. H/R Regulates mRNA Expression of HIF-1 α , VEGF-A, Collagen I, and Collagen III. Under H/R conditions, the mRNA expression of HIF-1 α was significantly suppressed (Figure 7(a)). The VEGF-A mRNA level significantly

increased when the cells were under hypoxia and at R24h compared with that of the control group ($P < 0.05$) but significantly decreased when the cells were at R72h and R96h compared with that of the H group ($P < 0.05$) (Figure 7(b)). The mRNA expression of collagen I significantly increased under hypoxia, returned to normal levels at R24h, and significantly decreased from R48h to R96h compared with those of the control and H groups ($P < 0.05$ or $P < 0.005$) (Figure 7(c)). The collagen III mRNA levels also significantly increased under hypoxia compared with that of the control group but remained at normal levels with different periods of reoxygenation ($P < 0.05$) (Figure 7(d)).

3.7. Immunocytochemical Staining. Figure 8(a) shows the representative fluorescence micrographs of HIF-1 α , VEGF-A, collagen I, and collagen III when the cells were under H/R conditions. We calculated the percentage of positive cells via cell counting Figure 8(b). Compared with the control group, the percentage of HIF-1 α -positive cells significantly increased under hypoxia (Figure 8), whereas that of VEGF-A-positive cells was significantly increased under hypoxia,

FIGURE 7: mRNA expression of (a) HIF-1 α , (b) VEGF-A, (c) collagen I, and (d) collagen III when cells were under H/R conditions induced by 0.3 mM CoCl₂. * $P < 0.05$ vs. control group; ** $P < 0.005$ vs. control group; # $P < 0.05$ vs. H group; ## $P < 0.005$ vs. H group.

at R24h, and at R48h ($P < 0.05$) (Figure 8(CID="C002" value="B")). The percentages of collagen I- and collagen III-positive cells were significantly lower at R48h, R72h, and R96h compared with those in the control and H groups ($P < 0.05$ or $P < 0.005$) (Figures 8(CID="C004" value="C") and 8(CID="C006" value="D")).

3.8. H/R Does Not Affect Cellular Apoptosis but Alters Protein Expression of Bax and bcl-2. value="Flo" value="w" cytometry was conducted to investigate the apoptotic levels of cells under H/R conditions. The flow cytometric analysis showed that no significant differences in the number of apoptotic cells were found across the control, hypoxia, and reoxygenation groups (Figures 9(a) and 9(b)). Meanwhile, the ratio of Bax/bcl-2 protein expression was significantly upregulated after reoxygenation compared with those of the control and hypoxia groups ($P < 0.05$), with a peak at R48h (Figure 9(c)).

4. Discussion

The current study uncovered the cellular responses of tendon-derived cells to hypoxia and subsequent reoxygenation. Our results showed that different concentrations of CoCl₂ exerted distinct effects on cell viability. Hypoxia induced by low CoCl₂ concentrations (0.05 and 0.1 mM) enhanced cell viability, while further increases in CoCl₂ con-

centrations (0.15–0.4 mM) led to a gradual decrease in cell viability. With 24 h–72 h of reoxygenation, cell viability increased in cells treated with high concentrations of CoCl₂. Cell viability returned to the same level as the control group after 96 h of reoxygenation, irrespective of the initial CoCl₂ concentrations.

Previous studies have demonstrated that different cell lines show diverse tolerance profiles to CoCl₂-induced H/R. Zhang et al. demonstrated that CoCl₂ did not promote or attenuate the viability of A498 cells at low concentrations (0.05–0.2 mM), but when the concentration was increased to 0.25 mM, cell activity gradually declined [20]. The study by Shi et al. in a human ovarian carcinoma cell line showed that CoCl₂-induced (0.15 mM) hypoxia inhibited cell proliferation, which was subsequently recovered with reoxygenation [23]. Tong et al. showed that HepG2 cells treated with different concentrations of CoCl₂ (0.05, 0.1, 0.15, and 0.2 mM) exhibited significantly repressed cell viability in a concentration-dependent manner [24]. However, our data showed that cell viability was enhanced by CoCl₂-induced hypoxia, even when the concentration of CoCl₂ reached 0.3 mM. Furthermore, during the early period of reoxygenation, cell viability further increased, especially in cells treated with high concentrations of CoCl₂ (0.25–0.4 mM). In addition, H/R had no obvious negative effects on cell growth. These results suggested that tendon-derived cells

FIGURE 8: (a) Representative fluorescence micrographs of HIF-1α, VEGF-A, collagen I, and collagen III in cells under H/R conditions induced by 0.3 mM CoCl₂. (b) Percentages of positive cells stained with HIF-1α (A), VEGF-A (B), collagen I (C), and collagen III (D). Scale bar: 100 μm. **P* < 0.05 vs. control group; ***P* < 0.005 vs. control group; #*P* < 0.05 vs. H group; ##*P* < 0.005 vs. H group.

were highly tolerant to CoCl₂-induced hypoxia and the reoxygenation.

H/R conditions can cause oxidative stress, which is a major mechanism involved in the pathogenesis of H/R injury [26–29]. Therefore, we examined the levels of two oxidative stress markers, MDA and SOD, and the activity of bioactive mitochondria in cells under H/R. MDA, a byproduct of lipid oxidation, is a biomarker of oxidative stress of cells [30]. SOD is the main antioxidant enzyme in cells that plays important roles in scavenging oxygen free radicals and resisting the damage of oxygen free radicals

[31]. Mitochondria, accounting for the majority of oxygen consumption, can help to tune cellular and organismal hypoxia responses [32]. Our results showed that the MDA level increased, the SOD level decreased, and the activity of bioactive mitochondria was enhanced under H/R, indicating that H/R induced notable oxidative stress in the tendon-derived cells. However, the oxidative stress did not affect the cell growth in this study. We speculate that the increased mitochondrial activity may retain the metabolic demand of the cells and induce an adaptive response to oxidative stress [33–36].

FIGURE 9: Flow cytometric analysis of cellular apoptosis and ratios of Bax/bcl-2 proteins: (a) representative graphs of flow cytometric assays showing the apoptosis of different groups; (b) the histogram showing the ratio of apoptotic cells; (c) expression ratio of Bax/bcl-2 protein. * $P < 0.05$ vs. control group; ** $P < 0.005$ vs. control group. # $P < 0.05$ vs. H group; ## $P < 0.005$ vs. H group.

HIF-1 α is a key transcription factor in response to hypoxic stress and is widely expressed in mammals, including humans, under hypoxic conditions [37, 38]. In the present study, the HIF-1 α protein level increased notably in the cells under hypoxia and then returned to the same level in the control cells immediately after reoxygenation. We noted, however, that the changes in HIF-1 α mRNA and protein expression were inconsistent. Specifically, HIF-1 α mRNA expression decreased significantly during hypoxia and reoxygenation, which is inconsistent with previous studies [39, 40]. A possible explanation is that the expression of HIF-1 α mRNA changes prior to the expression of protein, and the peak of mRNA expression occurred in less than 24 hours. Then, the inhibited HIF-1 α mRNA expression caused the recovery of the HIF-1 α protein level in the cells under reoxygenation. These results indicate that tendon-derived cells can regulate the expression of HIF-1 α rapidly under H/R conditions. It may be beneficial to the adaptation of tendon-derived cells to oxygen deprivation, leading to the high tolerance of this type of cells to H/R.

VEGF-A is an essential growth factor for most tissues in their response to traumatic injuries involving disrupted blood supply. Previous studies demonstrated that the expres-

sion of the VEGF gene significantly increased during the early period of tendon healing and that intraoperative delivery of VEGF notably enhanced tendon healing in a chicken model, indicating the importance of VEGF in tendon healing [41, 42]. In our study, we found that VEGF-A was upregulated under hypoxia when HIF-1 α was activated, and its levels returned to normal during reoxygenation along with the passage of time. Our results were consistent with previous studies investigating the role of the HIF-1 α /VEGF pathway in hypoxia. Liang et al. showed that hypoxia markedly upregulated VEGF-A and that appropriate vascular response might be essential for normal repair and remodeling. VEGF-A elevation is a rapid and strong response to hypoxic insult typically seen in most tissues [43]. Therefore, hypoxia-promoted expression of VEGF may be a self-protective mechanism after tendon injury.

Collagen is the major substance secreted by tendon cells. It is the main component of tendon ECM, and thus, synthesis of collagen is essential for maintaining tendon structure [44, 45]. A normal tendon comprises 65%–85% of collagen I, which is the most abundant collagen, followed by collagen III [46, 47]. Collagen III is highly upregulated during tendon healing [48]. Collagen III primarily forms an abundant but

disorganized collagen matrix in the proliferative phase of tendon healing. Webster et al. showed that when tendon-derived cells were cultured at low oxygen tensions, cellular metabolism was depressed and both total protein and collagen production were reduced [49]. It was hypothesized that the hypoxic environment might not have satisfied the physiological oxygen requirements of the cells, and it may have deleterious effects on collagen production [50]. A study also reported that there was an abrupt reduction of collagens I and III in the early stages of Achilles tendinitis; the tendon subsequently developed substantial impairment and Achilles tendinitis eventually occurred [51]. Our results showed that the expression of collagens I and III was downregulated during reoxygenation. Therefore, the in vitro reoxygenation process may correspond to the in vivo repair process after ischemia. Interestingly, the decline in the expression of collagen appears to contradict the increased cell viability. We speculate that although cell growth is not affected, other cellular functions are inhibited under hypoxic stress.

Apoptosis plays a critical role in the homeostasis of normal tissue [52]. In human rotator cuff tendinopathy, HIF-1 α accumulation was associated with cellular apoptosis, which provided an early support for the role of hypoxia-induced damage to cell loss via apoptosis [53]. However, Sasabe et al. reported that forced expression of HIF-1 α suppressed hypoxia-induced apoptosis of human oral squamous cell carcinoma cell lines [54]. Based on flow cytometry analysis in the present study, CoCl₂-induced H/R had a minimal effect on the cellular apoptosis. However, the protein expression of Bax was upregulated while that of bcl-2 was downregulated, resulting in a significant increase in the Bax/bcl-2 ratio during reoxygenation. The Bax/bcl-2 ratio reached its peak at 48 h after reoxygenation and began to decrease thereafter. This suggested that although the apoptotic pathway was activated, the cells did not actually undergo apoptosis within the observed period. We speculate that tendon-derived cells are highly resistant to H/R-induced apoptosis, and their inherent cell characteristics may prevent or even reverse the occurrence of apoptosis. Another possibility is that the initiation of the apoptotic mechanism precedes the occurrence of apoptosis. To this regard, extending the observation time may allow us to obtain a more complete picture of apoptotic events after H/R.

5. Conclusions

Collectively, the present study successfully established a CoCl₂-induced H/R model in tendon-derived cells, which provided a framework for future studies to understand the tendon-specific features of this widely observed stress response mechanism. The data in this study showed that hypoxia followed by reoxygenation for a certain period of time promoted cell viability in a concentration-dependent manner. Tendon-derived cells exhibited considerable tolerance to hypoxia. H/R caused oxidative stress responses but did not affect cell growth. H/R altered the expression of HIF-1 α , VEGF-A, collagen I, and collagen III. Cellular apo-

ptosis was not affected by H/R, but the Bax/bcl-2 ratio increased during reoxygenation.

Data Availability

All data generated or analyzed during this study are included in this published article.

Conflicts of Interest

All authors have read the journal's policy on disclosure of potential conflicts of interest and declare no conflict of interest.

Acknowledgments

This work was supported by the Jiangsu Provincial Medical Youth Talent (grant number QNRC2016701), Natural Science Foundation of Jiangsu Province (grant number BK20181203), and Six Talent Peaks Project in Jiangsu Province (grant number WSW-048).

Supplementary Materials

Table S1: antibodies used in the Western blot assay and immunocytochemical staining. Table S2: sequences of qPCR primers. (*Supplementary Materials*)

References

- [1] Y. Zhang, B. Wang, W. J. Zhang, G. Zhou, Y. Cao, and W. Liu, "Enhanced proliferation capacity of porcine tenocytes in low O₂ tension culture," *Biotechnology Letters*, vol. 32, no. 2, pp. 181–187, 2010.
- [2] J. Zhang and J. H. Wang, "Human tendon stem cells better maintain their stemness in hypoxic culture conditions," *PLoS One*, vol. 8, no. 4, article e61424, 2013.
- [3] Y. Yu, L. Lin, Y. Zhou et al., "Effect of hypoxia on self-renewal capacity and differentiation in human tendon-derived stem cells," *Medical Science Monitor*, vol. 23, pp. 1334–1339, 2017.
- [4] D. Rempel and S. O. Abrahamsson, "The effects of reduced oxygen tension on cell proliferation and matrix synthesis in synovium and tendon explants from the rabbit carpal tunnel: an experimental study in vitro," *Journal of Orthopaedic Research*, vol. 19, no. 1, pp. 143–148, 2001.
- [5] F. K. Fung, B. Y. Law, and A. C. Lo, "Lutein attenuates both apoptosis and autophagy upon cobalt (II) chloride-induced hypoxia in rat Müller cells," *PLoS One*, vol. 11, no. 12, article e0167828, 2016.
- [6] H. Song, I. Y. Han, Y. Kim et al., "The NADPH oxidase inhibitor DPI can abolish hypoxia-induced apoptosis of human kidney proximal tubular epithelial cells through Bcl 2 up-regulation via ERK activation without ROS reduction," *Life Sciences*, vol. 126, pp. 69–75, 2015.
- [7] S. Guo, A. Tjärnlund-Wolf, W. Deng et al., "Comparative transcriptome of neurons after oxygen-glucose deprivation: potential differences in neuroprotection versus reperfusion," *Journal of Cerebral Blood Flow and Metabolism*, vol. 38, no. 12, pp. 2236–2250, 2018.
- [8] Y. Zhang, Q. Pan, and Y. Liu, "CXCL16 silencing alleviates hepatic ischemia reperfusion injury during liver transplantation

- by inhibiting p 38 phosphorylation," *Pathology, Research and Practice*, vol. 216, no. 5, p. 152913, 2020.
- [9] W. Hou, X. Zhu, J. Liu, and J. Map, "Inhibition of miR-153 ameliorates ischemia/reperfusion-induced cardiomyocytes apoptosis by regulating Nrf2/HO-1 signaling in rats," *Biomedical Engineering Online*, vol. 19, no. 1, p. 15, 2020.
 - [10] D. Fang, Z. Li, Q. Zhong-ming et al., "Expression of bystin in reactive astrocytes induced by ischemia/reperfusion and chemical hypoxia in vitro," *Biochimica et Biophysica Acta*, vol. 1782, no. 11, pp. 658–663, 2008.
 - [11] J. M. Downey, "Free radicals and their involvement during long-term myocardial ischemia and reperfusion," *Annual Review of Physiology*, vol. 52, no. 1, pp. 487–504, 1990.
 - [12] J. E. Jung, G. S. Kim, H. Chen et al., "Reperfusion and neurovascular dysfunction in stroke: from basic mechanisms to potential strategies for neuroprotection," *Molecular Neurobiology*, vol. 41, no. 2-3, pp. 172–179, 2010.
 - [13] Y. Zhai, H. Petrowsky, J. C. Hong, R. W. Busuttil, and J. W. Kupiec-Weglinski, "Ischaemia-reperfusion injury in liver transplantation—from bench to bedside," *Nature Reviews. Gastroenterology & Hepatology*, vol. 10, no. 2, pp. 79–89, 2013.
 - [14] W. Z. Wang, R. C. Baynosa, and W. A. Zamboni, "Update on ischemia-reperfusion injury for the plastic surgeon: 2011," *Plastic and Reconstructive Surgery*, vol. 128, no. 6, pp. 685e–692e, 2011.
 - [15] D. N. Granger and P. R. Kvietys, "Reperfusion injury and reactive oxygen species: the evolution of a concept," *Redox Biology*, vol. 6, pp. 524–551, 2015.
 - [16] P. Sharma and N. Maffulli, "Tendon injury and tendinopathy: healing and repair," *The Journal of Bone and Joint Surgery. American Volume*, vol. 87, no. 1, pp. 187–202, 2005.
 - [17] T. Friedrich, W. Schmidt, D. Jungmichel, L. C. Horn, and C. Josten, "Histopathology in rabbit Achilles tendon after operative tenolysis (longitudinal fiber incisions)," *Scandinavian Journal of Medicine & Science in Sports*, vol. 11, no. 1, pp. 4–8, 2001.
 - [18] R. Stange, H. Sahin, B. Wieskötter et al., "In vivo monitoring of angiogenesis during tendon repair: a novel MRI-based technique in a rat patellar tendon model," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 23, no. 8, pp. 2433–2439, 2015.
 - [19] M. Aslam, K. D. Schluter, S. Rohrbach et al., "Hypoxia-reoxygenation-induced endothelial barrier failure: role of RhoA, Rac1 and myosin light chain kinase," *The Journal of Physiology*, vol. 591, no. 2, pp. 461–473, 2013.
 - [20] N. Zhang, B. Hong, C. Zhou et al., "Cobalt chloride-induced hypoxia induces epithelial-mesenchymal transition in renal carcinoma cell lines," *Annals of Clinical and Laboratory Science*, vol. 47, no. 1, pp. 40–46, 2017.
 - [21] T. Kietzmann, H. Schmidt, I. Probst, and K. Jungermann, "Modulation of the glucagon-dependent activation of the phosphoenolpyruvate carboxykinase gene by oxygen in rat hepatocyte cultures. Evidence for a heme protein as oxygen sensor," *FEBS Letter*, vol. 311, no. 3, pp. 251–255, 1992.
 - [22] A. Menon, P. Creo, M. Piccoli et al., "Chemical activation of the hypoxia-inducible factor reversibly reduces tendon stem cell proliferation, inhibits their differentiation, and maintains cell undifferentiation," *Stem Cells International*, vol. 2018, Article ID 9468085, 13 pages, 2018.
 - [23] J. Shi, Y. Wan, and W. Di, "Effect of hypoxia and reoxygenation on cell invasion and adhesion in human ovarian carcinoma cells," *Oncology Reports*, vol. 20, no. 4, pp. 803–807, 2008.
 - [24] Y. Tong, K. Tong, Q. Zhu et al., "Cobalt chloride induced apoptosis by inhibiting GPC3 expression via the HIF-1 α /c-Myc axis in HepG2 cells," *Oncotargets and Therapy*, vol. Volume 12, pp. 10663–10670, 2019.
 - [25] Y. F. Wu, C. Chen, J. B. Tang, and W. F. Mao, "Growth and stem cell characteristics of tendon-derived cells with different initial seeding densities: an in vitro study in mouse flexor tendon cells," *Stem Cells and Development*, vol. 29, no. 15, pp. 1016–1025, 2020.
 - [26] N. Ming, H. S. T. Na, J. L. He, Q. T. Meng, and Z. Y. Xia, "Propofol alleviates oxidative stress via upregulating lncRNA-TUG1/Brg1 pathway in hypoxia/reoxygenation hepatic cells," *Journal of Biochemistry*, vol. 166, no. 5, pp. 415–421, 2019.
 - [27] Y. Hu, Z. Mao, L. Xu et al., "Protective effect of dioscin against intestinal ischemia/reperfusion injury via adjusting miR-351-5p-mediated oxidative stress," *Pharmacological Research*, vol. 137, pp. 56–63, 2018.
 - [28] N. E. Hoffman, B. A. Miller, J. F. Wang et al., "Ca²⁺ entry via Trpm 2 is essential for cardiac myocyte bioenergetics maintenance," *American Journal of Physiology. Heart and Circulatory Physiology*, vol. 308, no. 6, pp. H637–H650, 2015.
 - [29] M. J. Berridge, P. Lipp, and M. D. Bootman, "The versatility and universality of calcium signalling," *Nature Reviews. Molecular Cell Biology*, vol. 1, no. 1, pp. 11–21, 2000.
 - [30] P. Eschwege, V. Paradis, M. Conti et al., "In situ detection of lipid peroxidation by-products as markers of renal ischemia injuries in rat kidneys," *The Journal of Urology*, vol. 162, no. 2, pp. 553–557, 1999.
 - [31] H. Chen, P. H. Chow, S. K. Cheng, A. L. M. Cheung, L. Y. L. Cheng, and W. S. O, "Male genital tract antioxidant enzymes: their source, function in the female, and ability to preserve sperm DNA integrity in the golden hamster," *Journal of Andrology*, vol. 24, no. 5, pp. 704–711, 2003.
 - [32] G. S. McElroy and N. S. Chandel, "Mitochondria control acute and chronic responses to hypoxia," *Experimental Cell Research*, vol. 356, no. 2, pp. 217–222, 2017.
 - [33] K. M. Porter, B. Y. Kang, S. E. Adesina, T. C. Murphy, C. M. Hart, and R. L. Sutliff, "Chronic hypoxia promotes pulmonary artery endothelial cell proliferation through H2O2-induced 5-lipoxygenase," *PLoS One*, vol. 9, no. 6, article e98532, 2014.
 - [34] F. G. Thankam, I. S. Chandra, A. N. Kovilam et al., "Amplification of mitochondrial activity in the healing response following rotator cuff tendon injury," *Scientific Reports*, vol. 8, no. 1, article 17027, 2018.
 - [35] B. H. Jiang, G. L. Semenza, C. Bauer, and H. H. Marti, "Hypoxia-inducible factor 1 levels vary exponentially over a physiologically relevant range of O₂ tension," *The American Journal of Physiology*, vol. 271, no. 4, pp. C1172–C1180, 1996.
 - [36] H. F. Burgers, D. W. Schelshorn, W. Wagner, W. Kuschinsky, and M. H. Maurer, "Acute anoxia stimulates proliferation in adult neural stem cells from the rat brain," *Experimental Brain Research*, vol. 188, no. 1, pp. 33–43, 2008.
 - [37] S. Cao, S. Yang, C. Wu, Y. Wang, J. Jiang, and Z. Lu, "Protein expression of hypoxia-inducible factor-1 alpha and hepatocellular carcinoma: a systematic review with meta-analysis," *Clinics and Research in Hepatology and Gastroenterology*, vol. 38, no. 5, pp. 598–603, 2014.

- [38] A. J. Majmundar, W. J. Wong, and M. C. Simon, "Hypoxia-inducible factors and the response to hypoxic stress," *Molecular Cell*, vol. 40, no. 2, pp. 294–309, 2010.
- [39] H. Li, X. Li, X. Jing et al., "Hypoxia promotes maintenance of the chondrogenic phenotype in rat growth plate chondrocytes through the HIF-1 α /YAP signaling pathway," *International Journal of Molecular Medicine*, vol. 42, no. 6, pp. 3181–3192, 2018.
- [40] L. Liu, W. Liu, L. Wang, T. Zhu, J. Zhong, and N. Xie, "Hypoxia-inducible factor 1 mediates intermittent hypoxia-induced migration of human breast cancer MDA-MB-231 cells," *Oncology Letters*, vol. 14, no. 6, pp. 7715–7722, 2017.
- [41] C. H. Chen, Y. Cao, Y. F. Wu, A. J. Bais, J. S. Gao, and J. B. Tang, "Tendon healing in vivo: gene expression and production of multiple growth factors in early tendon healing period," *The Journal of Hand Surgery*, vol. 33, no. 10, pp. 1834–1842, 2008.
- [42] W. F. Mao, Y. F. Wu, Q. Q. Yang et al., "Modulation of digital flexor tendon healing by vascular endothelial growth factor gene transfection in a chicken model," *Gene Therapy*, vol. 24, no. 4, pp. 234–240, 2017.
- [43] M. Liang, H. R. Cornell, N. Zargar Baboldashti, M. S. Thompson, A. J. Carr, and P. A. Hulley, "Regulation of hypoxia-induced cell death in human tenocytes," *Advances in Orthopedics*, vol. 2012, no. 984950, 12 pages, 2012.
- [44] J. Zhang, C. Keenan, and J. H. Wang, "The effects of dexamethasone on human patellar tendon stem cells: implications for dexamethasone treatment of tendon injury," *Journal of Orthopaedic Research*, vol. 31, no. 1, pp. 105–110, 2013.
- [45] R. James, G. Kesturu, G. Balian, and A. B. Chhabra, "Tendon: biology, biomechanics, repair, growth factors, and evolving treatment options," *The Journal of Hand Surgery*, vol. 33, no. 1, pp. 102–112, 2008.
- [46] G. Riley, "The pathogenesis of tendinopathy. A molecular perspective," *Rheumatology*, vol. 43, no. 2, pp. 131–142, 2004.
- [47] Y. Bi, D. Ehrichtiou, T. M. Kilts et al., "Identification of tendon stem/progenitor cells and the role of the extracellular matrix in their niche," *Nature Medicine*, vol. 13, no. 10, pp. 1219–1227, 2007.
- [48] K. Ueda, O. Shimizu, S. Oka, M. Saito, M. Hide, and M. Matsumoto, "Distribution of tenascin-C, fibronectin and collagen types III and IV during regeneration of rat submandibular gland," *International Journal of Oral and Maxillofacial Surgery*, vol. 38, no. 1, pp. 79–84, 2009.
- [49] D. F. Webster and H. C. Burry, "The effects of hypoxia on human skin, lung and tendon cells in vitro," *British Journal of Experimental Pathology*, vol. 63, no. 1, pp. 50–55, 1982.
- [50] E. A. Makris, D. J. Responde, N. K. Paschos, J. C. Hu, and K. A. Athanasiou, "Developing functional musculoskeletal tissues through hypoxia and lysyl oxidase-induced collagen cross-linking," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 111, no. 45, pp. E4832–E4841, 2014.
- [51] S. M. Mosier, G. Pomeroy, and A. Manoli 2nd., "Pathoanatomy and etiology of posterior tibial tendon dysfunction," *Clinical Orthopaedics and Related Research*, vol. 365, pp. 12–22, 1999.
- [52] J. Yuan, G. A. Murrell, A. Q. Wei, and M. X. Wang, "Apoptosis in rotator cuff tendonopathy," *Journal of Orthopaedic Research*, vol. 20, no. 6, pp. 1372–1379, 2002.
- [53] R. T. Benson, S. M. McDonnell, H. J. Knowles, J. L. Rees, A. J. Carr, and P. A. Hulley, "Tendinopathy and tears of the rotator cuff are associated with hypoxia and apoptosis," *Journal of Bone and Joint Surgery. British Volume*, vol. 92-B, no. 3, pp. 448–453, 2010.
- [54] E. Sasabe, Y. Tatemoto, D. Li, T. Yamamoto, and T. Osaki, "Mechanism of HIF-1 α -dependent suppression of hypoxia-induced apoptosis in squamous cell carcinoma cells," *Cancer Science*, vol. 96, no. 7, pp. 394–402, 2005.