

Research Article

Protective Effect of Epigallocatechin-3-Gallate in Hydrogen Peroxide-Induced Oxidative Damage in Chicken Lymphocytes

Xiaoqing Chi , Xiaodan Ma, Zoushuai Li, Yong Zhang, Yong Wang, Lijia Yuan, Ye Wu, Wei Xu , and Songhua Hu 

Department of Veterinary Medicine, College of Animal Sci., Zhejiang University, Hangzhou, Zhejiang 310058, China

Correspondence should be addressed to Songhua Hu; songhua@zju.edu.cn

Received 15 July 2020; Revised 4 December 2020; Accepted 17 December 2020; Published 31 December 2020

Academic Editor: Ana Cipak Gasparovic

Copyright © 2020 Xiaoqing Chi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Epigallocatechin-3-gallate (EGCG) is one of the fundamental compounds in green tea. The present study was to evaluate the protective effect of EGCG in oxidative damage and apoptosis induced by hydrogen peroxide (H_2O_2) in chicken lymphocytes. Results showed that preincubation of lymphocytes with EGCG significantly decreased H_2O_2 -reduced cell viability and apoptotic cells with DNA damage, restored the H_2O_2 -dependent reduction in total antioxidant capacity (T-AOC), glutathione peroxidase (GSH-PX), superoxide dismutase (SOD), glutathione (GSH), and glutathione disulfide (GSSG), and suppressed the increase in intracellular reactive oxygen species (ROS), nitric oxide (NO), nitric oxide synthesis (NOS), malondialdehyde (MDA), lipid peroxide (LPO), and protein carbonyl (Carbonyl). In addition, preincubation of the cells with EGCG increased mitochondrial membrane potential (MMP) and reduced calcium ion ($[Ca^{2+}]_i$) load. The protective effect of EGCG in oxidative damage in lymphocytes was accompanied by mRNA expression of SOD, Heme oxygenase-1 (HO-1), Catalase (CAT), GSH-PX, nuclear factor erythroid 2-related factor 2 (Nrf2), and thioredoxin-1 (Trx-1). As EGCG had been removed before lymphocytes were challenged with H_2O_2 , the activation of genes such as Nrf2 and Trx-1 by preincubation with EGCG could be the main reason for EGCG to protect the cells from oxidative damage by H_2O_2 . Since oxidative stress is an important mechanism of biological damage and is regarded as the reasons of several pathologies, the present findings may be helpful for the use of tea products to prevent oxidative stress and maintain healthy in both humans and animals.

1. Introduction

The poultry industry has become more and more important in daily life. The consumption of global meat has increased 52% since 2001, and the proportion of chicken consumption has risen from 32% to 42% [1]. In 2014, FAO predicted that poultry production will surpass pork production and become the first in 2022 [2]. However, poultry diseases are important harmful factors that affect the animals. In the poultry industry, oxidative stress is regarded to play an important role in the pathophysiology of infectious bursal disease, Newcastle disease, avian influenza, etc. [3]. Reactive oxygen species (ROS) and free radicals generated from various stresses or physiological factors may be responsible for oxidative stress and various pathological effects, including DNA damage, mitochondrial dysfunction, and cell injury [4]. Oxidative stress can cause cell damage and excessive production of

ROS or other oxygen-free radicals (e.g., $ONOO^{\bullet-}$, $O_2^{\bullet-}$, and hydrogen peroxide) have been found to attenuate the antioxidant system in the organs such as the brain and testis, particularly in the spleen [5, 6]. It has been reported that the innate and adaptive immunity is damaged during oxidative stress [7, 8]. Suppressed immune responses caused by oxidative stress have been found in both mammals and birds [9–11]. Therefore, the control of oxidative stress in the poultry industry may improve the bird's immunity and enhance the resistance of poultry against diseases and is of significance in veterinary practice.

Some natural compounds have been found a tendency to quench redundant-free radicals and decrease oxidative stress [12]. Al-Sayed et al. reported that a proanthocyanidin-rich fraction obtained from *B. hookeri* (BHPF) had the hepato- and nephroprotective effect, and the fraction was able to reduce the rate of lipid peroxidation, enhance the antioxidant

defense status, and guard against the pathological changes of the liver and kidney caused by oxidative damage [13]. A previous study showed that oral administration of tea extract made from green tea, the leaf of a plant *Camellia sinensis* (L) O. Ktze., had an effect in oxidative stress induced by cyclophosphamide in chickens. Green tea contains large amounts of tea catechins with antioxidant properties [14]. Epigallocatechin-3-gallate (EGCG) (Figure 1), epigallocatechin (EGC), epicatechin-3-gallate (ECG), and epicatechin (EC) are the main components [15]; meanwhile, EGCG rates 50-80% of total catechins [16]. Katiyar et al. highlighted that the EGCG could be a topical agent for ultimate usefulness against ROS-associated inflammatory dermatosis, photoaging, and photocarcinogenesis induced by UVB [17].

Because of the rich natural resource of tea, it has become possible to use the leaf and its products as an antioxidant agent in poultry production. Although numerous studies have been reported on the antioxidative activities of tea, no study has been found regarding the mechanisms underlying its effect in oxidative stress in chickens.

The present study was designed to investigate the effect of EGCG against oxidative stress of chicken lymphocytes induced by hydrogen peroxide (H_2O_2) with the goal of further understanding of tea and its products as an antioxidant agent in chicken production.

2. Materials and Methods

2.1. Chemicals. Epigallocatechin-3-gallate (EGCG) (93894), 3-(4, 5-dimethyl-2-thiazolyl)-2, 5-diphenyl-2-H-tetrazolium bromide (MTT) (M2128), and 34.5-36.5% H_2O_2 (18304) were the products of Sigma-Aldrich Inc. (St. Louis, Missouri, USA); phosphate-buffered saline (PBS) (10010023) and RPMI-1640 medium (7200047) were the products of Gibco (Thermo Fisher Scientific, Waltham, USA); lymphocyte separation medium (P9120) and cell lysis buffer (R0030) were the products of Solarbio Co. Ltd. (Beijing, China); fetal bovine serum (FBS) (13011-8611) was the product of Sijiqing Co. (Hangzhou, China).

2.2. Reagent Kits. Coomassie Brilliant Blue (CBB) kit (A045-2-2) and detection kits for total antioxidant capacity (T-AOC) (A015-2-1), glutathione peroxidase (GSH-PX) (A005-1-2), nitric oxide (NO) (A013-2-1), nitric oxide synthesis (NOS) (A014-2-2), malondialdehyde (MDA) (A003-4-1), lipid peroxide (LPO) (A106-1-3), and protein carbonyl (Carbonyl) (A087-1-2) were the products of Nanjing Jiancheng Institute of Bioengineering and Technology (Nanjing, China); detection kits for glutathione (GSH), glutathione disulfide (GSSG) (G263), and DNA damage quantification kit (DK02) were the products of Dojindo Molecular Technologies (Rockville, MD, USA); assay kits for reactive oxygen species (ROS) and superoxide dismutase (SOD) (ab139476), genomic DNA extraction kit (ab156900), and intracellular calcium ion ($[Ca^{2+}]_i$) detection kit (Fluo-3/AM) (ab145254) were the products of Abcam (Cambridge, UK); apoptosis detection kit (556547) was the product of BD Biosciences (San Jose, CA, USA); mitochondrial membrane potential (MMP) detection kit (JC-1) (M8650) was the prod-




FIGURE 1: Structure of epigallocatechin-3-gallate (EGCG).

TABLE 1: Experimental design.

Groups	<i>n</i>	H_2O_2 (μ M)	EGCG (μ M)
H_2O_2 +EGCG 90 μ M	6	80	90
H_2O_2 +EGCG 45 μ M	6	80	45
H_2O_2 +EGCG 22.5 μ M	6	80	22.5
H_2O_2 (model)	6	80	0
Normal control	6	0	0

uct of Solarbio Co. (Beijing, China); RNAiso™ Plus kit (9108) and PrimeScript™ RT reagent kit (6210A) were the products of Takara (Dalian, China).

2.3. Lymphocytes. The procedures on handling animals in this experiment were approved by the Institutional Animal Care and Use Committee of Zhejiang University. Lymphocytes were isolated from chickens as described by Bi et al. [18]. Briefly, the spleens were collected from 20 d old female Sanhuang chickens (Ningbo Zhenning Stock Breeding Inc., Ningbo, China) and polished to homogenate. Cell suspension was obtained by gently pushing the homogenate through a 70 μ m sterile plastic mesh. The suspension was centrifuged at $2,000 \times g$ for 15 min at room temperature on an equivalent volume of lymphocyte separation medium (1.085-1.092 g/mL). White cells in the interface were collected and washed. Cell density was adjusted to 5×10^5 cells/mL and cultured in RPMI 1640 containing HEPES and 2 mM glutamine, 100 units of penicillin/mL, and 100 μ g of streptomycin/mL, supplemented with 10% FBS at 39°C in a humidified atmosphere containing 5% CO_2 . The viable cells were more than 95% by trypan blue exclusion test, and the lymphocyte proportion was more than 85%.

2.4. Treatment of Cells with EGCG and H_2O_2 . To each well of a 12-well culture plate, 1 mL of cell suspension (5×10^5 cells) was added first, and EGCG was then added at the final concentrations of 90, 45, 22.5, or 0 μ M. The mixtures were incubated at 39°C in a humidified atmosphere in 5% CO_2 for 6 h. Afterward, cells were washed in PBS twice to remove EGCG and incubated in a media containing H_2O_2 (80 μ M) for an additional 4 h. Cells without treatment of EGCG and H_2O_2 were used as normal control or treated with H_2O_2 only were served as models (Table 1). After centrifuging the plates at $1,000 \times g$ for 10 min, the supernatant was discarded and the cells were resuspended in PBS with a density of 5×10^5 cells/mL for determination of cell viability, intracellular

enzymes and redox parameters, apoptosis, mitochondrial membrane potential (MMP), $[Ca^{2+}]_i$ concentration, and mRNA expression of SOD, GSH-PX, heme oxygenase-1 (HO-1), catalase (CAT), nuclear factor erythroid 2-related factor 2 (Nrf2), and thioredoxin-1 (Trx-1).

2.5. Cell Viability. The cell viability was measured by the MTT method [19]. Briefly, $100\ \mu\text{L}$ of cell suspension (5×10^4 cells) was added to 96-well culture plates. Then, $50\ \mu\text{L}$ of MTT solution (2 mg/mL) was added and incubated for 4 h. After that, the plates were centrifuged at $1,000 \times g$ for 10 min; untransformed MTT in supernatant was carefully removed. The MTT formazan was dissolved by adding $150\ \mu\text{L}$ of DMSO (0.04 N HCl). The optical density (OD) was read at 570 nm in Multiskan FC (Thermo, USA). The well was set in triplicate.

2.6. Assay for Intracellular Antioxidant Enzymes and Redox Products. To measure intracellular antioxidant enzymes and redox products, lymphocytes (5×10^5 cells) were lysed by a lysis buffer to release the intracellular content for biochemical analysis. Protein concentration was determined by the Coomassie Brilliant Blue (CBB) kit with bovine serum albumin (BSA) as standard. The levels of T-AOC, GSH-PX, NO, NOS, MDA, LPO, and Carbonyl were estimated according to the manufacturers' instructions by spectrophotometry using kits obtained from Nanjing Jiancheng Institute of Bioengineering and Technology and expressed as per milligram protein (mgprot). The activities of GSH and GSSG were measured using GSSH and GSH Quantification kits acquired from Dojindo Molecular Technologies according to the manufacturers' instructions and expressed as micromole per liter ($\mu\text{mol/L}$). In brief, the assay is based on the glutathione-dependent reduction of 5,5'-dithiobis-2-nitrobenzoic acid to 5-mercapto-2-nitrobenzoic acid (λ_{max} : 415 nm) [20–28].

2.7. Assay for Intracellular ROS and SOD. To measure intracellular ROS and SOD, $500\ \mu\text{L}$ of cell suspension (2.5×10^5 cells) was mixed with $10\ \mu\text{L}$ ROS and $10\ \mu\text{L}$ SOD detection reagents in a flow tube and incubated under 37°C for 30 min in dark with periodic shakings. The ROS/SOD detection kit includes two fluorescent dyes, the green dye is for ROS detection, and the orange dye is for SOD detection. The fluorescent colors can be visualized by flow cytometry. In this study, signals resulting from ROS and SOD were reflected in the FL1 and FL2 channels, respectively, and detected by FCM (BD FACSCalibur).

2.8. Detection of Cell Damage. The ab156900 genomic DNA extraction kit was used for the extraction of genomic DNA from the cells according to previously described by Kuechler et al. [29]. Each sample was adjusted to contain $100\ \mu\text{g}$ of extracted DNA per milliliter (μmL). Oxidative attack on the deoxyribose moiety by hydroxy radical will lead to the release of free bases from DNA, generating strand breaks with various sugar modifications and abasic sites (AP). AP sites are one of the major types of damage generated by ROS. Aldehyde Reactive Probe (ARP) reagent (N' -aminooxymethyl-carbonylhydrazino-D-biotin) reacts specifically with an

TABLE 2: Primer sequences used for real-time qPCR assay.

Gene [†]	Sequence (5'-3') [‡]	Product size (bp)
SOD	CGTCATTCACTTCGAGCAGAAGG	233
	GTCTGAGACTCAGACCACATA	
HO-1	ACTTCTATGGCAGCAACT	129
	AATAGCGGGTGTAGGC	
CAT	CTGTTGCTGGAGAATCTGGGTC	160
	TGGCTATGGATGAAGGATGGAA	
GSH-PX	TTGTAAACATCAGGGGCAAA	140
	TGGGCCAAGATCTTTCTGTAA	
Nrf2	ATTGAGCAAGTTTGGGAGGA	214
	AAGACACTGTAAGTCAGGAAT	
Trx-1	GGA	152
	GCAGGACAGGCTGGAAGTCACA	
β -Actin	CGAGAAGTGCGAGGTGAACG	152
	GAGAAATTGTGCGTGACATCA	
	CCTGAACCTCTCATTGCCA	

[†]SOD: superoxide dismutase; HO-1: Heme oxygenase-1; CAT: catalase; GSH-PX: glutathione peroxidase; Nrf2: nuclear factor erythroid 2-related factor 2; Trx-1: thioredoxin-1. [‡]Shown as forward primer followed by reverse primer.

aldehyde group which is an open ring form of the AP sites. After treatment of DNA containing AP sites with ARP reagent, AP sites are tagged with biotin residues. All AP sites will be converted to biotin-tagged forms by an excess amount of ARP. Therefore, AP sites can be quantified using avidin-biotin assay. The number of AP sites was measured using a DNA damage quantification kit based on calorimetric assay as reported [30]. Briefly, $10\ \mu\text{L}$ of DNA extract and an equal volume of ARP solution was mixed in a $0.5\ \text{mL}$ vial and incubated at 37°C for 1 h. After dilution of ARP-labeled genomic DNA with $310\ \mu\text{L}$ of buffer, $60\ \mu\text{L}$ of the diluted solution was taken to mix with $60\ \mu\text{L}$ of standard ARP-DNA solution in 96-well plates. Then, $100\ \mu\text{L}$ of DNA binding solution was added and mixed well. The plate was kept at room temperature (25°C) overnight. After DNA binding solution was discarded, the plate was washed with PBS and added with diluted $150\ \mu\text{L}$ of HRP-streptavidin solution. After incubation of the plate at 37°C for 1 h, $100\ \mu\text{L}$ of substrate solution was added and incubated at 37°C for another 1 h. The optical density (OD) was read at 650 nm. The number of AP sites in the genomic DNA was calculated according to the linear calibration curve generated for each experiment using ARP-DNA standard solutions.

2.9. Apoptosis Detection. An apoptosis detection kit was used to measure apoptosis of the cells as previously described [31]. Cells were washed in PBS at $1,000 \times g$ for 10 min and resuspended in $1\ \text{mL}$ of $1 \times$ binding buffer with adjusting to a concentration of 5×10^5 cells/mL. Then, $100\ \mu\text{L}$ of the cell suspension was mixed with $7\ \mu\text{L}$ annexin V-FITC and $7\ \mu\text{L}$ propidium iodide (PI) staining solution and incubated for 20 min at 37°C in dark. Afterward, $300\ \mu\text{L}$ of $1 \times$ binding

buffer was added, and fluorescence in cells was detected by FCM. The data were analyzed using FlowJo V10 software, and apoptotic rate was expressed as the ratio of apoptotic cells to the total number of cells.

2.10. Measurement of Mitochondrial Membrane Potential (MMP). Mitochondrial membrane potential (MMP) of the lymphocytes was measured using 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethyl-imidacarbocyanine iodide (JC-1) by flow cytometry as previously described [32]. Briefly, cells (2.5×10^5 cells) were cultured in 0.5 mL of medium containing JC-1 ($5 \mu\text{g/mL}$) in dark for 30 min at 37°C with periodic shaking to ensure a beneficial suspension. After washing by centrifugation at $1,000 \times g$ for 10 min in staining buffer, the cells were resuspended in 0.5 mL of PBS and then assayed in FCM within 1 h. The data were analyzed using FlowJo V10 software. The reduction of the red/green ratio illustrates decreased MMP, which is presented by the polymer mean fluorescent intensity (MFI)/monomer MFI.

2.11. Assay for Intracellular Calcium Ion ($[\text{Ca}^{2+}]_i$). Fluo-3/AM fluorescence density was measured to determine intracellular $[\text{Ca}^{2+}]_i$ concentration as previously described [33]. In brief, $10 \mu\text{L}$ of Fluo-3/AM was added into 1 mL of cell suspensions (5×10^5 cells) and incubated at 37°C for 45 min in dark. After washing, the cells were suspended in PBS, and Fluo-3/AM fluorescent intensity was measured by flow cytometry. Data were analyzed using FlowJo V10 software and expressed as MFI of Fluo-3/AM.

2.12. Real-Time Quantitative PCR Analysis. Total RNA was extracted from lymphocytes (5×10^5 cells) using RNAiso™ Plus kit according to the manufacturer's protocol. The purity and quality of total RNA were evaluated by the ratio of absorption (260:280 nm) ranged from 1.8 to 2.0 and quantitated on a NanoPhotometer spectrophotometer (IMPLEN, CA, USA) [34]. Subsequently, $2 \mu\text{g}$ of total RNA was used to synthesize cDNA using a PrimeScript™ RT reagent kit based on the manufacturer's instructions. The primers for RT-qPCR analysis were listed in Table 2. β -Actin was used as an internal control to normalize gene expression. Primers were designed by Primer 5 software and synthesized by Takara (Takara, Dalian, China). For RT-qPCR reactions, $20 \mu\text{L}$ mixtures were made up, containing $2 \mu\text{L}$ cDNA, $10 \mu\text{L}$ SYBR Taq, $1.6 \mu\text{L}$ forward and reverse primer (1:1), $0.4 \mu\text{L}$ ROX Reference Dye, and $6 \mu\text{L}$ RNase-free water. Each sample was set with triplicate. SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1 expression of messenger RNA (mRNA) were detected by performing RT-qPCR reactions on Bio-Rad T100 (Bio-Rad Laboratories, Inc., USA). The PCR procedure included 95°C for 2 min, followed by 35 cycles of 95°C for 10 s, 58°C for 30 s, and 72°C for 30 s. Only one peak for each PCR product was analyzed by melting curve. The comparative Ct value method was used to quantify mRNA expression relative to β -actin expression using the $2^{-\Delta\Delta\text{CT}}$ method. The relative mRNA expressions of SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1 were expressed as mean \pm standard deviation (S.D.).




FIGURE 2: Effect of EGCG on cell viability. Lymphocytes (5×10^4) were treated with EGCG (0, 22.5, 45, and $90 \mu\text{M}$) for 6 h first and then cultured in media with (model) or without (normal control) H_2O_2 ($80 \mu\text{M}$) for additional 4 h. Cell viability was determined by the MTT assay using the absorbance of formazan at 570 nm. Data are presented as % of the normal control (mean \pm S.D., $n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).

2.13. Statistical Analysis. One-way ANOVA of SPSS Statistics for Mac (Version 20) was used for statistical analysis of the data, followed by Tukey's post hoc test for multiple comparisons between groups. The results were presented as the means \pm S.D. All statistical analyses were performed using the GraphPad InStat software (Version 7, GraphPad). P value < 0.05 or P value < 0.01 was considered statistically significant.

3. Results

3.1. Effect of EGCG on Cell Viability. To evaluate the effect of EGCG on cell viability, lymphocytes were cultured with EGCG first and then treated with H_2O_2 to induce oxidative stress of the cells. Cell viability was determined by the MTT assay. Figure 2 showed that the number of viable cells significantly decreased after lymphocytes were treated with H_2O_2 (model) when compared to the normal control ($P < 0.01$). However, pretreatment of the cells with EGCG increased the number of viable cells in a dose-dependent manner at the range of 22.5– $90 \mu\text{M}$ when compared to the model group.

3.2. EGCC Increased Antioxidant Capacity and Decreased Redox Products. To observe the redox state in the lymphocytes, intracellular contents released from the cells were analyzed. Figures 3(a)–3(d) showed that incubation of the lymphocytes with H_2O_2 significantly suppressed enzymatic (T-AOC and GSH-PX) and nonenzymatic (GSH and GSSG) antioxidant capacity; and preincubation with EGCG significantly enhanced the antioxidant ability when compared to




FIGURE 3: Intracellular antioxidants and redox products. Lymphocytes (5×10^5) were treated with EGCG (0, 22.5, 45, and $90 \mu\text{M}$) for 6 h first and then cultured in media with (model) or without (normal control) H_2O_2 ($80 \mu\text{M}$) for an additional 4 h. The cells were treated in lysis buffer to release the intracellular materials for analysis of enzymatic (T-AOC and GSH-PX) and nonenzymatic (GSH and GSSG) antioxidants (a–d) and redox products (NO, NOS, MDA, LPO, and Carbonyl) (e–i). Data are presented as mean \pm S.D. ($n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).




FIGURE 4: Intracellular ROS and SOD. Lymphocytes (2.5×10^5) were treated with EGCG (0, 22.5, 45, and 90 μM) for 6 h first and then cultured in media with (model) or without (normal control) H_2O_2 (80 μM) for 4 h, then dyed for ROS and SOD. Cells with increased production of SOD showed orange fluorescence and were detected with the FL2 channel. Cells with increased production of ROS demonstrated green light and were registered in the FL1 channel. (a) Flow cytometric gating percentage; (b) $H_2O_2 + EGCG$ 90 μM group; (c) $H_2O_2 + EGCG$ 45 μM group; (d) $H_2O_2 + EGCG$ 22.5 μM group; (e) H_2O_2 (model) group; (f) normal control group; (g) ROS levels represented by mean green fluorescence intensity in the indicated groups; (h) SOD levels represented by mean orange fluorescence intensity in the indicated groups. Data are presented as mean \pm S.D. ($n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).

the model group, indicating that pretreatment of lymphocytes with EGCG prevented H_2O_2 -induced suppression in antioxidant capacity in cells. Figures 3(e)–3(i) showed that redox products (NO, NOS, MDA, LPO, and Carbonyl) were significantly increased after incubation of the lymphocytes

with H_2O_2 but preincubation with EGCG significantly decreased the products, suggesting that pretreatment of the lymphocytes with EGCG could prevent the increased production of redox products due to decreased antioxidant capacity induced by H_2O_2 .




FIGURE 5: Effect of EGCG on cell DNA damage. Lymphocytes (5×10^5) were treated with EGCG (0, 22.5, 45, and 90 μ M) for 6 h first and then cultured in media with (model) or without (normal control) H₂O₂ (80 μ M) for an additional 4 h. The oxidative DNA damage in cells was estimated with apurinic/apryrimidinic sites (AP sites). The number of AP sites was measured using a DNA damage quantification kit based on calorimetric assay. Data are presented as mean \pm S.D. ($n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).

3.3. EGCC Decreased ROS and Increased SOD Generation. To observe intracellular ROS and SOD, lymphocytes were dyed with a ROS/SOD detection kit and analyzed by FCM. Figure 4 showed that intracellular ROS significantly increased and SOD numerically decreased after incubation of the lymphocytes with H₂O₂ when compared to the normal control. However, preincubation of the lymphocytes with EGCG significantly decreased ROS and increased SOD when compared to the model. The results indicated that pretreatment of the lymphocytes prevented the intracellular production of reactive oxygen species (ROS) and enhanced the capacity of superoxide dismutase (SOD) in cells in stress induced by H₂O₂.

3.4. EGCC Decreased Cell Damage and Apoptosis. In this study, the number of apurinic/apryrimidinic sites (AP sites) was used to estimate cell DNA damage, and FCM analysis was used to detect apoptotic cells (Annexin V+/PI-). Figures 5 and 6 showed that incubation of the lymphocytes with H₂O₂ significantly increased the number of the AP sites and apoptotic cells; preincubation of cells with EGCG significantly decreased the number of the AP sites and apoptotic cells. The results suggested that pretreatment with EGCG prevented cell damage and apoptosis caused by H₂O₂ treatment.

3.5. EGCC Increased Mitochondrial Membrane Potential and Decreased [Ca²⁺]_i. To estimate the integrity of the cell membrane, mitochondrial membrane potential (MMP) was measured with JC-1, and the change of [Ca²⁺]_i concentrations in

cells was estimated using Fluo-3/AM. Incubation of the lymphocytes with H₂O₂ significantly decreased MMP (Figure 7) and increased the [Ca²⁺]_i in cells (Figure 8); however, preincubation with EGCG significantly increased MMP (Figure 7) and decreased the [Ca²⁺]_i in cells (Figure 8), suggesting that pretreatment with EGCG prevented cell membrane damage resulting from H₂O₂ treatment.

3.6. EGCC Increased mRNA Expression of SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1. RT-qPCR was used to quantify the expression of antioxidant genes. Figure 9 showed that incubation of the lymphocytes with H₂O₂ depressed the expression of mRNA of SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1 in cells, while preincubation with EGCG significantly increased the expression of the antioxidant genes when compared to the model group, indicating that pretreatment of the cells with EGCG recovered the depressed expression of antioxidant genes caused by H₂O₂ treatment.

4. Discussion

The protective effect of EGCG in oxidative damage induced by H₂O₂ in lymphocytes was demonstrated. Preincubation of lymphocytes with EGCG significantly decreased H₂O₂-induced intracellular ROS, DNA damage, cell apoptosis, [Ca²⁺]_i overload, and increased MMP, which were accompanied by the increased antioxidant capacity (T-AOC, GSH-PX, SOD, GSH, and GSSG) and decreased redox products (NO, NOS, MDA, LPO, and Carbonyl). These changes were in association with mRNA expression of SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1.

Lymphocytes are important cells in the cellular and humoral immunity. As lymphocyte membranes are rich in unsaturated fatty acids, the cells are among the main target for oxidative stress and particularly easy to be attacked by ROS. In the poultry production, chickens are usually exposed to many harmful factors such as condensed population, polluted air, and contaminated feed or drinking water, which may cause overproduction of ROS. Increased ROS may induce oxidative stress, damage the structure of lymphocytes, and suppress the immunity. H₂O₂ has been usually used to induce oxidative stress in vitro. Like the process found in vivo, exogenous H₂O₂ crosses the cell membrane; destroys nucleic acids, proteins, and lipids; demolishes [Ca²⁺]_i homeostasis; and activates mitochondria signals, ultimately leading to cell apoptosis [35–38]. In the present study, exposure of lymphocytes to H₂O₂ for 4 h lead to increase intracellular ROS (Figure 4), DNA damage (Figure 5), apoptotic cells (Figure 6), decrease MMP (Figure 7), and cause [Ca²⁺]_i overload (Figure 8), resulting in significantly decreased cell viability (Figure 2). However, pretreatment of the cells with EGCG decreased intracellular ROS, DNA damage, apoptotic cells, and [Ca²⁺]_i load and improved MMP with increased cell viability when compared to model cells. Because of the damage of lymphocytes during oxidative stress, inhibited immune responses to vaccination in association with oxidative stress were observed in chickens in our previous studies [39–41].

DNA damage under oxidative stress is a result of the interaction of DNA and ROS, mainly hydroxy radical which




FIGURE 6: Effect of EGCG on cell apoptosis. Lymphocytes (5×10^5) were treated with EGCG (0, 22.5, 45, and 90 μM) for 6 h first and then cultured in media with (model) or without (normal control) H_2O_2 (80 μM) for an additional 4 h. Apoptotic cells (annexin V+/PI-) were discriminated with FCM analysis. (a) Flow cytometric gating percentage; (b) H_2O_2 + EGCG 90 μM group; (c) H_2O_2 + EGCG 45 μM group; (d) H_2O_2 + EGCG 22.5 μM group; (e) H_2O_2 (model) group; (f) normal control group; (g) the ratio of apoptotic cells to the total number of cells in the indicated groups. Values are expressed as mean \pm S.D. ($n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).

is converted from superoxide and hydrogen peroxide by the Fenton reaction. Hydroxy radicals cause a multiplicity of modifications in DNA. Oxidative attacks by hydroxy radicals on the deoxyribose moiety lead to the release of free bases from DNA, generating strand breaks with various sugar

modifications and simple abasic sites (AP sites). AP sites are one of the major types of damage generated by ROS. The redundant ROS changes physiological homeostasis and causes tissue damage [42]. Figure 5 showed that the number of AP sites was significantly increased after H_2O_2




FIGURE 7: Effect of EGCG on mitochondrial depolarization. Lymphocytes (2.5×10^5) were treated with EGCG (0, 22.5, 45, and 90 μ M) for 6 h first and then cultured in media with (model) or without (normal control) H₂O₂ (80 μ M) for an additional 4 h. Cells were incubated with JC-1 (5 μ g/mL) and assayed by FCM. Mitochondrial depolarization was represented by a reduction in the red/green fluorescence intensity ratio. (a) Flow cytometric gating percentage; (b) H₂O₂ + EGCG 90 μ M group; (c) H₂O₂ + EGCG 45 μ M group; (d) H₂O₂ + EGCG 22.5 μ M group; (e) H₂O₂ (model) group; (f) Normal Control group; (g) bar diagram represented JC-1 polymer/monomer MFI. Values are expressed as mean \pm S.D. ($n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).

treatment but decreased after pretreatment of the cells with EGCG, indicating that EGCG reduced DNA damage caused by H₂O₂.

The poultry antioxidant system comprises enzymatic and nonenzymatic defenses [43, 44]. Enzymatic defenses include T-AOC, SOD, and GSH-PX. The total antioxidant capacity

(T-AOC) may give more biologically relevant information than the individual antioxidants only [45]. SOD is the first line of cellular defense against oxidative damage, which converts O₂^{•-} into O₂ or H₂O₂, decomposes into H₂O ultimately [46]. GSH-PX is an enzyme responsible for reducing soluble hydrogen peroxide and alkyl peroxides, scavenging ROS and




FIGURE 8: Effect of EGCG on H₂O₂-induced change in the [Ca²⁺]_i levels. Lymphocytes (5×10^5) were treated with EGCG (0, 22.5, 45, and 90 μ M) for 6 h first and then cultured in media with (model) or without (normal control) H₂O₂ (80 μ M) for an additional 4 h. Afterward, the cells were incubated with Fluo3/AM ([Ca²⁺]_i probe), and Fluo-3/AM fluorescence was measured by FCM. (a) Flow cytometric gating percentage; (b) the fluorescence intensity of Fluo-3/AM reflected the [Ca²⁺]_i levels; (c) bar diagram represented Fluo-3/AM MFI. Values are expressed as mean \pm S.D. ($n = 6$). Bars with different letters are significantly different ($P < 0.05$).

its oxidation products to protect the body from lipid peroxidation [47, 48]. All these enzymes work together to eliminate active oxygen species [49]. The present study elucidated that H₂O₂ treatment significantly (T-AOC and GSH-PX) or numerically (SOD) depressed the enzymes, and pretreatment with EGCG neutralized the inhibitory effect of H₂O₂ on T-AOC, GSH-PX, and SOD as indicated in Figures 3(a) and 3(b) and 4(h). Nonenzymatic antioxidant defense includes GSH and GSSG. GSH is one of the major cellular antioxidants [50–52]. It can directly react with radicals such as superoxide and deliver electrons for the reduction of peroxides by GSH-PX [50]. The product of such reactions is GSSG, which is reduced in cells to GSH in the NADPH-dependent reaction, catalyzed by glutathione reductase [50, 52]. In the present study, GSH and GSSG were significantly decreased after the cells were treated with H₂O₂ and increased when the cells were pretreated with EGCG as indicated in

Figures 3(c) and 3(d). Therefore, enzymatic and nonenzymatic antioxidants may cooperate and provide protection against ROS attack.

Lipid peroxidation is generated by the conversion of polyunsaturated fatty acid into lipid peroxides or MDA. As a major product of LPO, MDA is supposed to be a marker of tissue damage [53, 54]. Protein carbonyl is one of the final products of protein peroxidation, which is formed by oxidation via either an increase of ROS or the attack of reactive aldehydes such as NO, NOS, MDA, and LPO formed during lipid peroxidation [55], namely, in a way that LPO aggravates protein peroxidation. Accordingly, the protein carbonyl level is considered a surrogate marker of protein peroxidation [56]. In addition, NO is likely to cross membranes and reacts with superoxide anion which forms as a by-product of respiration to produce peroxynitrite (ONOO^{•−}) and cytotoxic species that have been demonstrated to lead to lipid




FIGURE 9: Effect of EGCG on the relative mRNA expression. Lymphocytes (5×10^5) were treated with EGCG (0, 22.5, 45, and 90 μ M) for 6 h first and then cultured in media with (model) or without (normal control) H₂O₂ (80 μ M) for an additional 4 h. RT-qPCR was carried out to quantify mRNA of SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1 (a–f) and data were normalized using the $2^{-\Delta\Delta C_T}$ method. Values are expressed as % of the normal control (mean \pm S.D., $n = 6$). Bars with different letters are significantly different ($P < 0.05$ or $P < 0.01$).

peroxidation as a result of $[Ca^{2+}]_i$ dysfunction [57] and are theoretically conducive to enhance intracellular levels of ROS. Our study clearly showed a significant increase in NO, NOS, MDA, LPO, and Carbonyl post-H₂O₂ treatment, but EGCG counteracted the increase as indicated in Figures 3(e)–3(i), indicating that EGCG may effectively pro-

tect the tissue from damage by eliminating or diminishing the production of NO, NOS, MDA, LPO, or Carbonyl.

The antioxidant effect of tea in oxidative stress in chickens has been reported previously. Chi et al. observed that oral administration of tea saponins significantly increased T-AOC, SOD, CAT, GSH-PX, GSH, VC, and VE

and enhanced the immune responses to Newcastle disease and infectious bronchitis vaccines in chickens with oxidative stress induced by cyclophosphamide [41]. They recently found that oral administration of tea extract granule (TEG) significantly increased T-AOC, SOD, CAT, GSH-PX, and GSH as well as decreased Carbonyl, LPO, and MDA [58]. EGCG has a characteristic structure with a large number of hydroxyl groups. The presence of an *ortho*-trihydroxyl group in the B-ring has been shown to be important to the radical scavenging abilities of EGCG; a gallate moiety at the 3 position of the C-ring increases the radical scavenging effectiveness (Figure 1) [59]. There is evidence that at least some catechin metabolites maintain comparable antioxidant capacities to their parent compounds [60]. After administration of green tea in rats reaching roughly 400 mg of catechins, measured-free catechins in the plasma were found to account for only 20% of its additive total radical-trapping parameter (TRAP), proposed that catechin conjugates and metabolites might have contributed to the measured TRAP increase [61]. These findings showed that the biological activity of tea extract may depend on its special structure, and the metabolites combined in vivo can be also with a large state role. As EGCG is the major bioactive component with its antioxidant properties, EGCG may directly contribute to the antioxidant activity when tea and its products are orally administered.

In addition, EGCG may participate in the antioxidant defense through the thioredoxin (Trx-1) system comprising of thioredoxin and thioredoxin reductase and the nuclear factor erythroid 2-related factor 2 (Nrf2) system [62]. Under abnormal conditions, the Trx-1 system serves as the first line defense system against oxidative stress, sparing the activation of the Nrf2 system [63]. Consequently, enhanced expression of the Trx-1 in the cytoplasm by directly eliminating ROS and in the nucleus by restraining the phosphorylation of I κ B- α , restraining the activation of NF- κ B, and making an indirect effect on antioxidation [64]. The transcription factor Nrf2 is commonly involved in the transcriptional both constitutive and inducible regulation of genes encoding antioxidant proteins under stress conditions. Some natural sources can stimulate Nrf2 signaling pathways and may be the inducer of phase II detoxifying and antioxidant enzymes representing GSH-PX and SOD [38]. SOD is an antioxidant enzyme involved in the scavenging of superoxide radicals; in the antioxidant defense system of birds, GSH-PX is more important than in the case of mammals [65], and EGCG availability can be the limiting factor for the activation of GSH-PX mRNA in chickens. In addition to inducing several phase II detoxifying enzymes, Nrf2 is also involved in the de novo synthesis of various antioxidant enzymes responsible for protection against cytotoxicity caused by oxidative stress [66]. Nrf2 promotes the expression of antioxidant enzymes (e.g., HO-1 and CAT) as a preventable response to oxidative stress. As the rate-limiting enzyme of heme decomposition reaction, HO-1 can stimulate production of endogenous carbon monoxide, biliverdin, and ferrous; biliverdin reductase converts biliverdin into bilirubin, which the two substances are possible endogenous antioxidants entailed in the structure of the defense system against ROS and free radical chain

reactions (e.g., lipid peroxidation injury) [67]. Our data showed that an increased Nrf2 mRNA expression was accompanied by enhanced expression of genes SOD, CAT, GSH-PX, and HO-1 in EGCG-treated lymphocytes (Figure 8), suggesting that the activation of antioxidant genes by EGCG could mainly contribute to the reduced damage by H₂O₂ in EGCG-treated cells.

The present study showed in vitro antioxidant effect of EGCG, one of the fundamental compounds in tea. Tea and its products are common and popular in humans but rarely used in animals. Since oxidative stress is an important mechanism of biological damage and is regarded as the reasons of several pathologies, tea and its products may be useful to prevent oxidative stress and maintain healthy in both humans and animals.

5. Conclusion

The present study demonstrated the protective effect of EGCG in oxidative damage and apoptosis induced by H₂O₂ in chicken lymphocytes. Preincubation of lymphocytes with EGCG significantly decreased H₂O₂-reduced cell viability and apoptotic cells with DNA damage, restored the H₂O₂-dependent reduction in T-AOC, GSH-PX, SOD, GSH, and GSSG, and suppressed the increase in intracellular ROS, NO, NOS, MDA, LPO, and Carbonyl. In addition, preincubation of the cells with EGCG increased MMP and reduced [Ca²⁺]_i load. The protective effect of EGCG in oxidative damage in lymphocytes was accompanied by mRNA expression of SOD, HO-1, CAT, GSH-PX, Nrf2, and Trx-1. As EGCG had been removed before lymphocytes were challenged with H₂O₂, the activation of genes such as Nrf2 and Trx-1 by preincubation with EGCG could be the main reason for EGCG to protect the cells from oxidative damage by H₂O₂.

6. Innovation

The protective effect of EGCG was first demonstrated in oxidative damage and apoptosis induced by H₂O₂ in chicken lymphocytes. Since oxidative stress is an important mechanism of biological damage and is regarded as the reasons of several pathologies that affected animals, the present findings may be helpful for the use of tea products to prevent oxidative stress and maintain healthy in both humans and animals.

Data Availability

The data are available and acquired from the authors.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Xiaoqing Chi and Songhua Hu conceived and designed the experiments. Xiaoqing Chi, Xiaodan Ma, and Zoushuai Li performed the experiments. Xiaoqing Chi, Yong Zhang, and Ye Wu analyzed the data. Xiaoqing Chi, Yong Wang,

and Lijia Yuan contributed reagents and materials. Xiaoqing Chi and Songhua Hu wrote the paper.

Acknowledgments

The authors are grateful to the students in the Laboratory of Traditional Chinese Veterinary Medicine (TCVM Lab) for their assistance in the present study. The National Key R&D Program of China (Project No. 2017YFD0502200) was acknowledged.

References

- [1] N. Alloui and O. Bennoune, "Poultry production in Algeria: current situation and future prospects," *World's Poultry Science Journal*, vol. 69, no. 3, pp. 613–620, 2013.
- [2] P. Stevenson, D. Battaglia, C. Bullon, and A. Carita, "Review of animal welfare legislation in the beef, pork, and poultry industries," *FAO Investment Centre-Directions in Investment (FAO) eng*, no. 10, 2014.
- [3] F. J. Hoerr, "Clinical aspects of immunosuppression in poultry," *Avian Diseases*, vol. 54, no. 1, pp. 2–15, 2010.
- [4] J. Zielonka and B. Kalyanaraman, "ROS-generating mitochondrial DNA mutations can regulate tumor cell metastasis—a critical commentary," *Free Radical Biology & Medicine*, vol. 45, no. 9, pp. 1217–1219, 2008.
- [5] R. Ortiz-Butrón, V. Blas-Valdivia, M. Franco-Colin, M. Pineda-Reynoso, and E. Cano-Europa, "An increase of oxidative stress markers and the alteration of the antioxidant enzymatic system are associated with spleen damage caused by methimazole-induced hypothyroidism," *Drug & Chemical Toxicology*, vol. 34, no. 2, pp. 180–188, 2011.
- [6] G. Caputa, S. Zhao, A. E. G. Criado, D. S. Ory, J. G. Duncan, and J. E. Schaffer, "RNASET2 is required for ROS propagation during oxidative stress-mediated cell death," *Cell Death & Differentiation*, vol. 23, no. 2, pp. 347–357, 2016.
- [7] M. Maes, Z. Fišar, M. Medina, G. Scapagnini, G. Nowak, and M. Berk, "New drug targets in depression: inflammatory, cell-mediated immune, oxidative and nitrosative stress, mitochondrial, antioxidant, and neuroprogressive pathways. And new drug candidates—Nrf2 activators and GSK-3 inhibitors," *Inflammopharmacology*, vol. 20, no. 3, pp. 127–150, 2012.
- [8] J. C. Greene, A. J. Whitworth, L. A. Andrews, T. J. Parker, and L. J. Pallanck, "Genetic and genomic studies of *Drosophila parkin* mutants implicate oxidative stress and innate immune responses in pathogenesis," *Human Molecular Genetics*, vol. 14, no. 6, pp. 799–811, 2005.
- [9] Z. Zhang, Q. Wang, J. Zhang, S. Li, X. L. Wang, and S. W. Xu, "Effects of oxidative stress on immunosuppression induced by selenium deficiency in chickens," *Biological Trace Element Research*, vol. 149, no. 3, article 9439, pp. 352–361, 2012.
- [10] S. K. Katiyar, "Grape seed proanthocyanidines and skin cancer prevention: inhibition of oxidative stress and protection of immune system," *Molecular Nutrition & Food Research*, vol. 52, no. 1, pp. S71–S76, 2008.
- [11] S. I. Lee and K. S. Kang, "Omega-3 fatty acids modulate cyclophosphamide induced markers of immunosuppression and oxidative stress in pigs," *Scientific Reports*, vol. 9, no. 1, pp. 2684–2692, 2019.
- [12] L. R. Nissen, L. Månsson, G. Bertelsen, T. Huynh-Ba, and L. H. Skibsted, "Protection of dehydrated chicken meat by natural antioxidants as evaluated by electron spin resonance spectrometry," *Journal of Agricultural & Food Chemistry*, vol. 48, no. 11, pp. 5548–5556, 2000.
- [13] E. al-Sayed, M. M. Abdel-Daim, O. E. Kilany, M. Karonen, and J. Sinkkonen, "Protective role of polyphenols from *Bauhinia hookeri* against carbon tetrachloride-induced hepato- and nephrotoxicity in mice," *Renal Failure*, vol. 37, no. 7, pp. 1198–1207, 2015.
- [14] S. Tang, J. P. Kerry, D. Sheehan, and J. D. Buckley, "A comparative study of tea catechins and α -tocopherol as antioxidants in cooked beef and chicken meat," *European Food Research & Technology*, vol. 213, no. 4–5, pp. 286–289, 2001.
- [15] E. Al-Sayed and M. M. Abdel-Daim, "Analgesic and anti-inflammatory activities of epicatechin gallate from *Bauhinia hookeri*," *Drug Development Research*, vol. 79, no. 4, pp. 157–164, 2018.
- [16] R. Bigelow and J. A. Cardelli, "The green tea catechins, (–)-Epigallocatechin-3-gallate (EGCG) and (–)-Epicatechin-3-gallate (ECG), inhibit HGF/Met signaling in immortalized and tumorigenic breast epithelial cells," *Oncogene*, vol. 25, no. 13, pp. 1922–1930, 2006.
- [17] S. K. Katiyar, M. S. Matsui, C. A. Elmets, and H. Mukhtar, "Polyphenolic antioxidant (–)-epigallocatechin-3-gallate from green tea reduces UVB-induced inflammatory responses and infiltration of leukocytes in human skin," *Photochemistry and Photobiology*, vol. 69, no. 2, pp. 148–153, 1999.
- [18] S. Bi, X. Ma, Y. Wang et al., "Protective effect of ginsenoside rg1 on oxidative damage induced by hydrogen peroxide in chicken splenic lymphocytes," *Oxidative Medicine and Cellular Longevity*, vol. 2019, no. 3, Article ID 8465030, 13 pages, 2019.
- [19] P. Wang, S. M. Henning, and D. Heber, "Limitations of MTT and MTS-based assays for measurement of antiproliferative activity of green tea polyphenols," *PLoS One*, vol. 5, no. 4, article e10202, pp. 10202–10212, 2010.
- [20] F. Liu, X. Li, T. Lin et al., "The cyclophosphamide metabolite, acrolein, induces cytoskeletal changes and oxidative stress in Sertoli cells," *Molecular Biology Reports*, vol. 39, no. 1, pp. 493–500, 2012.
- [21] R. Oztürk-Urek, L. A. Bozkaya, and L. Tarhan, "The effects of some antioxidant vitamin- and trace element-supplemented diets on activities of SOD, CAT, GSH-Px and LPO levels in chicken tissues," *Cell biochemistry and function*, vol. 19, no. 2, pp. 125–132, 2001.
- [22] L. Musallam, C. Éthier, P. S. Haddad, F. Denizeau, and M. Bilodeau, "Resistance to Fas-induced apoptosis in hepatocytes: role of GSH depletion by cell isolation and culture," *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 283, no. 3, pp. G709–G718, 2002.
- [23] U. I. Walther, B. Wilhelm, S. Walther, H. Mückter, and B. Fichtl, "Zinc toxicity in various lung cell lines is mediated by glutathione and GSSG reductase activity," *Biological Trace Element Research*, vol. 78, no. 1–3, pp. 163–178, 2000.
- [24] Y. Zhao, W. Jiang, B. Li et al., "Artesunate enhances radiosensitivity of human non-small cell lung cancer A549 cells via increasing NO production to induce cell cycle arrest at G₂/M phase," *International Immunopharmacology*, vol. 11, no. 12, pp. 2039–2046, 2011.
- [25] M. Gilchrist, S. D. McCauley, and A. D. Befus, "Expression, localization, and regulation of NOS in human mast cell lines: effects on leukotriene production," *Blood*, vol. 104, no. 2, pp. 462–469, 2004.

- [26] D. Del Rio, A. J. Stewart, and N. Pellegrini, "A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress," *Nutrition, Metabolism and Cardiovascular Diseases*, vol. 15, no. 4, pp. 316–328, 2005.
- [27] A. Mezzetti, G. Zuliani, F. Romano et al., "Vitamin E and lipid peroxide plasma levels predict the risk of cardiovascular events in a group of healthy very old people," *Journal of the American Geriatrics Society*, vol. 49, no. 5, pp. 533–537, 2001.
- [28] K. B. Pandey and S. I. Rizvi, "Protection of protein carbonyl formation by quercetin in erythrocytes subjected to oxidative stress," *Medicinal Chemistry Research*, vol. 19, no. 2, pp. 186–192, 2010.
- [29] E. C. K  chler, P. N. Tannure, P. Falagan-Lotsch, T. S. Lopes, J. M. Granjeiro, and L. M. F. Amorim, "Buccal cells DNA extraction to obtain high quality human genomic DNA suitable for polymorphism genotyping by PCR-RFLP and real-time PCR," *Journal of Applied Oral Science*, vol. 20, no. 4, pp. 467–471, 2012.
- [30] W. Li, Y. Luo, F. Zhang et al., "Ischemic preconditioning in the rat brain enhances the repair of endogenous oxidative DNA damage by activating the base-excision repair pathway," *Journal of Cerebral Blood Flow & Metabolism*, vol. 26, no. 2, pp. 181–198, 2006.
- [31] J. Barber, C. West, A. E. Kiltie, S. A. Roberts, and D. Scott, "Detection of individual differences in radiation-induced apoptosis of peripheral blood lymphocytes in normal individuals, ataxia telangiectasia homozygotes and heterozygotes, and breast cancer patients after radiotherapy," *Radiation Research*, vol. 153, no. 5, pp. 570–578, 2000.
- [32] H. Tsai, S. Chang, C. Chou, T. P. Weng, C. C. Hsu, and J. S. Wang, "Exercise training alleviates hypoxia-induced mitochondrial dysfunction in the lymphocytes of sedentary males," *Scientific Reports*, vol. 6, no. 1, pp. 35170–35183, 2016.
- [33] R. E. Dolmetsch, R. S. Lewis, C. C. Goodnow, and J. I. Healy, "Differential activation of transcription factors induced by Ca^{2+} response amplitude and duration," *Nature*, vol. 386, no. 6627, pp. 855–858, 1997.
- [34] X. Cui, Y. Wang, B. Maqbool et al., "Early IgG response to foot and mouth disease vaccine formulated with a vegetable oil adjuvant," *Vaccines*, vol. 7, no. 4, pp. 143–153, 2019.
- [35] R. Duan, X. Xing, Y. Qi et al., "Taxane-derived compounds protect SK-N-SH cells against oxidative stress injury induced by H_2O_2 ," *Neurological Research*, vol. 39, no. 7, pp. 632–639, 2017.
- [36] W. Zhu, J. Xu, Y. Ge et al., "Epigallocatechin-3-gallate (EGCG) protects skin cells from ionizing radiation via heme oxygenase-1 (HO-1) overexpression," *Journal of Radiation Research*, vol. 55, no. 6, pp. 1056–1065, 2014.
- [37] L. Wang, Y. Chen, P. Sternberg, and J. Cai, "Essential roles of the PI3 kinase/Akt pathway in regulating Nrf2-dependent antioxidant functions in the RPE," *Investigative Ophthalmology & Visual Science*, vol. 49, no. 4, pp. 1671–1678, 2008.
- [38] V. R. Muthusamy, S. Kannan, K. Sadhaasivam et al., "Acute exercise stress activates Nrf2/ARE signaling and promotes antioxidant mechanisms in the myocardium," *Free Radical Biology & Medicine*, vol. 52, no. 2, pp. 366–376, 2012.
- [39] J. Yu, F. S. Shi, and S. Hu, "Improved immune responses to a bivalent vaccine of Newcastle disease and avian influenza in chickens by ginseng stem-leaf saponins," *Veterinary Immunology and Immunopathology*, vol. 167, no. 3–4, pp. 147–155, 2015.
- [40] S. Bi, X. Chi, Y. Zhang et al., "Ginsenoside Rg1 enhanced immune responses to infectious bursal disease vaccine in chickens with oxidative stress induced by cyclophosphamide," *Poultry Science*, vol. 97, no. 8, pp. 2698–2707, 2018.
- [41] X. Chi, S. Bi, W. Xu, Y. Zhang, S. Liang, and S. Hu, "Oral administration of tea saponins to relieve oxidative stress and immune suppression in chickens," *Poultry Science*, vol. 96, no. 9, pp. 3058–3067, 2017.
- [42] A. Djordjevic, S. Spasic, A. Jovanovic-Galovic, R. Djordjevic, and G. Grubor-Lajsic, "Oxidative stress in diabetic pregnancy: SOD, CAT and GSH-Px activity and lipid peroxidation products," *The Journal of Maternal-Fetal & Neonatal Medicine*, vol. 16, no. 6, pp. 367–372, 2004.
- [43] P. F. Surai, R. C. Noble, and B. K. Speake, "Relationship between vitamin E content and susceptibility to lipid peroxidation in tissues of the newly hatched chick," *British Poultry Science*, vol. 40, no. 3, pp. 406–410, 1999.
- [44] P. F. Surai, "Effect of selenium and vitamin E content of the maternal diet on the antioxidant system of the yolk and the developing chick," *British Poultry Science*, vol. 41, no. 2, pp. 235–243, 2000.
- [45] A. Ghiselli, M. Serafini, F. Natella, and C. Scaccini, "Total antioxidant capacity as a tool to assess redox status: critical view and experimental data," *Free Radical Biology & Medicine*, vol. 29, no. 11, pp. 1106–1114, 2000.
- [46] I. K. Hwang, W. S. Eum, K. Y. Yoo et al., "Copper chaperone for Cu,Zn-SOD supplement potentiates the Cu,Zn-SOD function of neuroprotective effects against ischemic neuronal damage in the gerbil hippocampus," *Free Radical Biology & Medicine*, vol. 39, no. 3, pp. 392–402, 2005.
- [47] F. N. Bebe and M. Panemangalore, "Exposure to low doses of endosulfan and chlorpyrifos modifies endogenous antioxidants in tissues of rats," *Journal of Environmental Science and Health, Part B*, vol. 38, no. 3, pp. 349–363, 2003.
- [48] J. Cui, J. Yuan, and Z. Zhang, "Anti-oxidation activity of the crude polysaccharides isolated from *Polygonum Cillinerve* (Nakai) Ohwi in immunosuppressed mice," *Journal of Ethnopharmacology*, vol. 132, no. 2, pp. 512–517, 2010.
- [49] S. Ince, I. Kucukkurt, H. H. Demirel, D. A. Acaroz, E. Akbel, and I. H. Cigerci, "Protective effects of boron on cyclophosphamide induced lipid peroxidation and genotoxicity in rats," *Chemosphere*, vol. 108, no. 1, pp. 197–204, 2014.
- [50] I. Rebrin, S. Zicker, K. J. Wedekind, I. Paetau-Robinson, L. Packer, and R. S. Sohal, "Effect of antioxidant-enriched diets on glutathione redox status in tissue homogenates and mitochondria of the senescence-accelerated mouse," *Free Radical Biology & Medicine*, vol. 39, no. 4, pp. 549–557, 2005.
- [51] D. P. Jones, "[11] Redox potential of GSH/GSSG couple: assay and biological significance," *Methods in Enzymology*, vol. 348, no. 11, pp. 93–112, 2002.
- [52] O. Zitka, S. Skalickova, J. Gumulec et al., "Redox status expressed as GSH:GSSG ratio as a marker for oxidative stress in paediatric tumour patients," *Oncology Letters*, vol. 4, no. 6, pp. 1247–1253, 2012.
- [53] X. Y. Zhang, W. G. Li, Y. J. Wu, and M. T. Gao, "Amelioration of doxorubicin-induced myocardial oxidative stress and immunosuppression by grape seed proanthocyanidins in tumour-bearing mice," *Journal of Pharmacy & Pharmacology*, vol. 57, no. 8, pp. 1043–1051, 2005.
- [54] S. Eroglu, D. Pandir, F. G. Uzun, and H. Bas, "Protective role of vitamins C and E in diclorvos-induced oxidative stress in

- human erythrocytes in vitro,” *Biological Research*, vol. 46, no. 1, pp. 33–38, 2013.
- [55] A. Haribabu, V. S. Reddy, C. Pallavi et al., “Evaluation of protein oxidation and its association with lipid peroxidation and thyrotropin levels in overt and subclinical hypothyroidism,” *Endocrine*, vol. 44, no. 1, pp. 152–157, 2013.
- [56] R. Negi, D. Pande, K. Karki, A. Kumar, R. S. Khanna, and H. D. Khanna, “Association of oxidative DNA damage, protein oxidation and antioxidant function with oxidative stress induced cellular injury in pre-eclamptic/eclamptic mothers during fetal circulation,” *Chemico-Biological Interactions*, vol. 208, no. 11, pp. 77–83, 2014.
- [57] R. Balez and L. Ooi, “Getting to NO Alzheimer’s disease: neuroprotection versus neurotoxicity mediated by nitric oxide,” *Oxidative Medicine and Cellular Longevity*, vol. 2016, no. 6157, Article ID 3806157, 8 pages, 2016.
- [58] X. Chi, Y. Zhang, X. Ma et al., “Antioxidative stress of oral administration of tea extract granule in chickens,” *Poultry Science*, vol. 99, no. 4, pp. 1956–1966, 2020.
- [59] F. Nanjo, M. Mori, K. Goto, and Y. Hara, “Radical scavenging activity of tea catechins and their related compounds,” *Bioscience, Biotechnology, and Biochemistry*, vol. 63, no. 9, pp. 1621–1623, 1999.
- [60] K. B. Pandey and S. I. Rizvi, “Plant polyphenols as dietary antioxidants in human health and disease,” *Oxidative Medicine and Cellular Longevity*, vol. 2, no. 5, Article ID 897484, 278 pages, 2009.
- [61] P. Pietta, P. Simonetti, C. Gardana, and P. Mauri, “Trolox equivalent antioxidant capacity (TEAC) of Ginkgo biloba flavonol and Camellia sinensis catechin metabolites,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 23, no. 1, pp. 223–226, 2000.
- [62] R. Dong, D. Wang, X. Wang et al., “Epigallocatechin-3-gallate enhances key enzymatic activities of hepatic thioredoxin and glutathione systems in selenium-optimal mice but activates hepatic Nrf2 responses in selenium-deficient mice,” *Redox Biology*, vol. 10, no. 9, pp. 221–232, 2016.
- [63] M. Cebula, E. E. Schmidt, and E. S. J. Arnér, “TrxR1 as a potent regulator of the Nrf2-Keap1 response system,” *Antioxidants & redox signaling*, vol. 23, no. 10, pp. 823–853, 2015.
- [64] K. Hirota, M. Murata, Y. Sachi et al., “Distinct roles of thioredoxin in the cytoplasm and in the nucleus,” *Journal of Biological Chemistry*, vol. 274, no. 39, pp. 27891–27897, 1999.
- [65] J. Huang, D. Li, H. Zhao et al., “The selenium deficiency disease exudative diathesis in chicks is associated with downregulation of seven common selenoprotein genes in liver and muscle,” *Journal of Nutrition*, vol. 141, no. 9, pp. 1605–1610, 2011.
- [66] T. Ishii, K. Itoh, S. Takahashi et al., “Transcription factor Nrf2 coordinately regulates a group of oxidative stress-inducible genes in macrophages,” *Journal of Biological Chemistry*, vol. 275, no. 21, pp. 16023–16029, 2000.
- [67] B. Whittle and C. Varga, “New light on the anti-colitic actions of therapeutic aminosalicylates: the role of heme oxygenase,” *Pharmacological Reports*, vol. 62, no. 3, pp. 548–556, 2010.