Research Article

Electroacupuncture Pretreatment Attenuates Intestinal Injury after Autogenous Orthotopic Liver Transplantation in Rats via the JAK/STAT Pathway

Lili Jia , Wenli Yu , Hongli Yu, and Yiqi Weng

Department of Anesthesiology, Tianjin First Center Hospital, Tianjin 300192, China

Correspondence should be addressed to Wenli Yu; yzxwenliyu@163.com

Received 18 April 2020; Revised 5 June 2020; Accepted 15 July 2020; Published 4 August 2020

Guest Editor: Weifeng Yao

Copyright © 2020 Lili Jia et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Liver transplantation induces self-injury and affects remote organs, such as the lung, kidney, and intestine. Postoperative intestinal dysfunction has been associated with prolonged hospitalization and affects a patient's health and quality of life. Electroacupuncture (EA) has been proven effective in multiple organ protection. However, the potential mechanism underlying the protective effects of EA on intestinal injury after liver transplantation remains unclear. **Methods.** After establishing an autogenous orthotopic liver transplantation (AOLT) model, we studied the effects of EA pretreatment on intestinal injury after AOLT. We used the JAK2-specific inhibitor AG490 to explore the underlying mechanism. Histological analysis and apoptosis assays were used to evaluate intestinal injury. Oxidative stress index and inflammatory response were also measured after AOLT. Furthermore, we detected the phosphorylation levels of JAK2, STAT1, and STAT3 by Western blot. **Results.** We found that pretreatment with EA alleviated intestinal injury after AOLT, as shown by HE staining and TUNEL methods. EA pretreatment inhibited the expressions of p-JAK2, p-STAT1, and p-STAT3 in the intestines after AOLT. Upon treatment with JAK2-specific inhibitor AG490, intestinal injury was balanced. **Conclusion.** The data indicated EA pretreatment alleviated intestinal injury after AOLT by inhibiting the JAK/STAT signaling pathway. These results provide basic evidence to support the potential therapeutic efficacy of EA.

1. Introduction

Liver transplantation is the most effective method to address end-stage liver diseases. During the past few decades, remote organ injury has been widely studied during liver transplantation. Liver transplantation not only induced self-injury but also affected remote organs, such as the lung, kidney, and intestine [1–4]. With the development of surgical techniques, postoperative mortality and morbidity significantly reduced in recent years. However, postoperative intestinal dysfunction remains associated with prolonged hospitalization and affects a patient's health and quality of life. Therefore, numerous studies have been conducted on the prevention of intestinal injury, from clinical trials to animal studies. Based on Nozato et al.'s study [5], we established an autogenous orthotopic liver transplantation (AOLT) model in rats, which can well imitate the clinical surgery process. Although

previous studies have explored various methods and drugs in ischemia organ injury, reported side effects have limited these studies from applying to clinical practice.

Acupuncture is a traditional Chinese medical treatment and is simple to perform, safe, and reliable. Electroacupuncture (EA) has been proven effective in multiple organ protection [6–8]. A previous study showed that excessive inflammation and oxidative stress response play vital roles in intestinal injury after liver ischemia-reperfusion [9]. Another study showed that EA protects against liver injury after liver ischemia-reperfusion [10]. Zusanli (ST36) is a widely used acupoint in humans to exert anti-inflammatory effects during acupuncture for treating pain or ischemia-reperfusion conditions [11, 12]. Herein, we used the ST36 acupoint to assess its protective effect against intestinal injury after AOLT in rats. Although EA has been increasingly used to treat various diseases, the mechanism remains unknown.

The Janus kinase/signal transducer and activator of the transcription (JAK/STAT) signaling pathway is involved in a wide range of distinct cellular processes, including inflammation, apoptosis, cell-cycle control, and development [13]. The JAK/STAT pathway comprises a family of receptor-associated cytosolic tyrosine kinases (JAKs) that phosphorylate tyrosine residues on bound transcription factors (STATs). JAK-mediated tyrosine phosphorylation of STAT family members enables the translocation of these transcription factors to the nucleus and leads to an augmentation of gene transcription [14, 15]. However, it is unclear whether the JAK/STAT signaling pathway is involved in EA pretreatment to prevent intestinal injury after AOLT. Based on the protective effect of EA on various tissues, our study is aimed at exploring the effect of EA on intestinal injury induced by AOLT in rats and at investigating the underlying mechanisms.

2. Methods and Materials

2.1. Animals. A total of 40 adult male Sprague-Dawley rats (220–250 g) were purchased from the People's Liberation Army Military Academy of Medical Sciences Laboratory Animal Center. The animals were housed in 12 h light-dark cycles with controlled room temperature and were fed with regular rat chow and water ad libitum but were fasted 12 h before experiments. Animals were randomly assigned into five groups: group A, sham operated (sham group); group B, autogenous orthotopic liver transplantation (AOLT group); group C, pretreated with EA (ST36, 1–2 mA, 2–100 Hz, 30 min) for 3 days+AOLT (EA+AOLT group); group D, pretreated with EA for 3 days+sham operated (EA+sham group); and group E, pretreated with EA for 3 days+AG490 (5 mg/kg, i.p., Selleck, USA) 30 min before establishing the AOLT model (EA+AOLT+AG490 group). The dosage was determined from a previous study [1]. The experimental procedures were conducted following the Guide for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee.

2.2. Animal Model. A rat AOLT model was established using a previously reported method [5]. Anesthesia was induced by inhalation of 3.0–4.0% isoflurane and maintained 1.5–2.0% isoflurane inhalation. During surgery, all rats were free to breathe O₂ and were prone on a heating blanket. Rats in the sham group underwent laparotomy without performing AOLT as control. Total liver ischemia was induced by clamping the hepatic artery, the portal vein, suprahepatic vena cava (SHVC), and intrahepatic vena cava (IHVC). After clamping, the portal vein and IHVC were cannulated with a polyethylene tube, and the liver was perfused through the portal vein with 500 mL of heparinized cold saline (Jiangsu Wanbang Biochemical Pharmaceutical Co., Ltd., China) (2.5 IU/mL) to wash out all blood from the liver. The tubes were removed after perfusion, and the openings of tube insertion were closed. The SHVC and IHVC were then declamped immediately after repairing the vessels. After 45 min, the clamps were removed to allow the return of blood flow to the liver. After 6 h, the animals were euthanized by administering CO₂ inhalation. Blood samples and intestinal tissue biopsies were

taken. The tissues were fresh frozen for histological and biochemical evaluation. (Figure 1).

2.3. EA Pretreatment. Previous studies have shown that treatment with diluted 100 Hz and 2 Hz alternating frequency (2/100 Hz) EA stimulation of the ST36 acupoint can alleviate the inflammatory response caused by Complete Freund's adjuvant (CFA) [16, 17]. Zusanli acupoint is located 5 mm beneath the capitulum fibulae and lateral posterior to the knee joint. We performed EA pretreatment as described in a previous study [18]. Two acupuncture needles (diameter, 0.25 mm; length, 30 mm) were inserted into bilateral ST36 acupoints. Then, the two acupuncture needles were connected to the output of a Master-9 Pulse Stimulator (AMPI, Israel). EA parameters were set as follows: bidirectional symmetric square wave. The intensity was adjusted to induce moderate muscle contraction of the hind limb (1–2 mA), and the EA groups were subjected to EA with alternating frequencies and pulse waves. EA was applied 30 min/day from day 1 to day 3 (Figure 1).

2.4. Histological Examination. Formalin-fixed intestinal tissue was dehydrated, embedded in paraffin, and sliced into 4 μ m thick sections, which were stained with hematoxylin and eosin (HE). The damage of intestinal mucosal cells was analyzed by the Chiu score method [19]: 0—normal mucosa, 1—the tip of the hair on the top of the subcutaneous gap increases, 2—moderate separation of the epithelium and lamina, 3—when there are a large number of villi on both sides of the top of the villi with partial villi, 4—villous damage with a large number of inherent layer of capillary exposed, and 5—inherent layer damage of bleeding and ulcers.

2.5. Apoptosis Assay. To detect the apoptotic positive cells, terminal deoxynucleotidyl transferase-mediated digoxigenin-labeled UTP nick end labeling assay (TUNEL) was performed using an ApopTag peroxidase in situ cell death detection kit (Roche, Basel, Switzerland). Briefly, the 4 μ m thick paraffin sections were deparaffinized, treated with 0.1% TritonX-100, and incubated at 37°C for 8 min in a humidified atmosphere. PBS wash was performed 3 times. Then, the sections were further incubated with a mixture of marker and enzyme solution at 37°C for 1 h. PBS wash was performed 3 times, and the samples were observed under a fluorescence microscope (Olympus, Japan) at 450–550 nm. Five high-power fields were randomly selected to determine the percentage of apoptotic cells.

2.6. Enzyme-Linked Immunosorbent Assay (ELISA). Blood samples were collected from all rats. The serum levels of tumor necrosis factor- α (TNF- α), high-mobility group box-1 (HMGB-1), intestinal fatty acid-binding protein (IFABP), and endotoxin (LPS) were examined using an ELISA kit (Shanghai Baiwo Co., Ltd., China) according to the manufacturer's instructions [20].

2.7. Wet/Dry (W/D) Ratio of the Intestine. To test the edema of intestinal after AOLT 6 h following reperfusion, 5 cm of ileum tissue at 2 cm from the back of the blind was harvested, cleaned by removing blood and water, and weighed. Then, the sample was incubated at 70°C for 24 h and weighed.

FIGURE 1: The autogenous orthotopic liver transplantation (AOLT) model in rats and electroacupuncture pretreatment on rats.

The *W/D* ratio was calculated as the ratio of the wet weight to the dry weight of the intestine.

2.8. Oxidative Stress. The levels of malondialdehyde (MDA) and superoxide dismutase (SOD) were determined using assay kits (Nanjing Institute Co., Ltd.) according to the manufacturer's instructions [20]. The tissue MDA level was determined by the Esterbauer and Cheeseman method based on its reaction with thiobarbituric acid at 90–100°C; absorbance was measured at 532 nm. MDA reacts with thiobarbituric acid (TBA) and produces a pink pigment, which has a maximum absorption at 532 nm. The value of each sample was obtained from the standard curve and was expressed as $\mu\text{mol/g}$ tissue. SOD activity was measured according to the Paoletti and Mocali method. In this assay, superoxide anion was generated from molecular oxygen in the presence of EDTA, manganese II chloride, and mercaptoethanol. Nicotinamide adenine dinucleotide phosphate oxidation was linked to the availability of superoxide anions in the medium.

2.9. Immunohistochemistry. Immunohistochemical staining of the intestinal tissue was performed on formalin-fixed paraffin sections using a microwave-based technique. The 4 μm thick sections of the fixed intestines were dewaxed with xylene, hydrated in graded concentrations of ethanol, and treated with 0.3% hydrogen peroxide for 10 min to quench endogenous peroxidase. The sections were heated in a microwave oven in sodium citrate (pH = 6.0) at 95–99°C for 15 min and then cooled at room temperature. The sections were incubated in 5% blocking serum for 30 min and then in primary antibodies (Cleaved Caspase-3: 1:1 000, from Cell Signaling Technology, Beverly, MA, USA) at 4°C overnight. They were subsequently incubated with biotinylated secondary antibodies for 30 min and finally counterstained with hematoxylin.

2.10. Western Blot. The protein lysates were prepared from frozen tissues in ice-cold RIPA buffer (Sigma-Aldrich). The extracted protein was separated in a 10% sodium dodecyl sulfate- (SDS-) PAGE and then electrophoretically transferred to a nitrocellulose membrane (Hybond, Amersham Biosciences, Little Chalfont, UK). Membranes were blocked with 5% nonfat milk powder in TBS for 1 h at room temperature. These membranes were subjected to immunoblot analysis with antibodies to JAK2, p-JAK2, STAT1, p-STAT1, STAT3, and p-STAT3. The protein antibody immune

complexes were detected with horseradish peroxidase-conjugated secondary antibodies and enhanced chemiluminescence reagents (Pierce Biotechnology, Rockford, IL). GAPDH was used as an internal control to calculate the ratio of optical density, and values were compared with those of sham controls.

2.11. Statistical Analysis. All values are given as the mean $\pm$ SD. Statistical analysis was carried out using GraphPad Prism 8.0 software (GraphPad Software, USA). One-way analysis of variance was used to compare the measurement data between groups. $P < 0.05$ was considered significant.

3. Results

3.1. EA Pretreatment Significantly Decreases Intestinal Histology Injury. To examine the protective effect of EA pretreatment against intestinal injury after AOLT, each rat was analyzed by histopathology, and the intestinal mucous damage score was calculated. A normal histological structure was observed in the sham group by HE staining. Severe disruption of structural integrity in brush border, including loss of mucus, villi, and widespread necrotic area, was observed in the AOLT group. However, these damages were improved in the EA pretreatment group. The intestinal mucous damage score significantly increased in the AOLT group compared with the sham group. EA pretreatment combined with AG490 improved intestinal damage; no significant differences were observed between the EA+sham group and the EA+AOLT+AG490 group (Figures 2).

3.2. EA Pretreatment Alleviates *W/D* Ratio of the Intestine. To test intestinal edema after AOLT, we calculated the *W/D* ratio changes in all groups. As shown in Table 1, EA pretreatment significantly reduced the *W/D* ratio caused by AOLT. When compared with the AOLT group, the intestinal *W/D* ratio decreased in the EA+sham and EA+AOLT+AG490 groups. The EA+sham group did not show a significant difference from the sham group (Table 1, Figure 3).

3.3. EA Pretreatment Decreases Intestinal Inflammatory Reaction and Oxidative Stress. As aforementioned, EA application was shown to protect against intestinal injury following AOLT. However, the underlying mechanisms remain unknown. Severe intestinal damage was found to be accompanied by oxidative stress and inflammatory reaction

FIGURE 2: EA pretreatment significantly decreases intestinal histology injury (magnification, $\times 200$). Representative microphotographs were taken from the intestine of the sham (a), AOLT (b), EA+AOLT (c), EA+sham (d), and EA+AOLT+AG490 (e) groups at the time point of 6 h after AOLT in rats. (f) Histopathological scoring was calculated in each group: (A–E) the sham, AOLT, EA+AOLT, EA+sham, and EA+AOLT+AG490 groups. Data were represented as the mean $\pm$ SD ($n = 8$, per group). * $P < 0.05$ vs. the sham group. # $P < 0.05$ vs. the AOLT group.

TABLE 1: W/D ratio of experiment rats ($n = 8$, $\bar{x} \pm s$).

Groups	W/D ratio
Sham	3.89 ± 0.81
AOLT	$9.35 \pm 0.13^*$
EA+AOLT	$5.85 \pm 0.55^{* \#}$
EA+sham	$3.70 \pm 0.57^{\#}$
EA+AOLT+AG490	$4.45 \pm 0.41^{\#}$

Data presented are the mean $\pm$ SD. * $P < 0.05$ vs. the sham group. # $P < 0.05$ vs. the AOLT group.

activation. The levels of TNF- α , HMGB-1, iFABP, LPS, and MDA increased markedly in the AOLT group. However, the application of EA effectively balanced oxidative stress and inflammatory reactions. The levels of TNF- α , HMGB-1, iFABP, LPS, and MDA decreased, whereas the level of SOD increased in EA+AOLT group. In the EA+sham group, EA had no effect on oxidative stress and inflammatory response, which were comparable with the results observed in the sham group. Results from the EA+AOLT+AG490 group were also similar to those from the EA+sham group (Figures 4 and 3).

3.4. EA Pretreatment Inhibits Intestinal P-JAK2, P-STAT1, and P-STAT3 Protein Expressions. The JAK/STAT pathway

FIGURE 3: Effects of EA on the plasma level of MDA, SOD, and the W/D ratio following AOLT-induced intestinal injury. The levels of plasma MDA, SOD, and the W/D ratio were measured at 6 h following AOLT. (A–E) The sham, AOLT, EA+AOLT, EA+sham, and EA+AOLT+AG490 groups. Data were represented as the mean $\pm$ SD ($n = 8$, per group). * $P < 0.05$ vs. the sham group. # $P < 0.05$ vs. the AOLT group.

FIGURE 4: Effects of EA on the plasma level of TNF- α , HMGB-1, IFABP, and LPS following AOLT induced intestinal injury. The levels of plasma TNF- α , HMGB-1, IFABP, and LPS were measured at 6 h following AOLT. (A–E) The sham, AOLT, EA+AOLT, EA+sham, and EA+AOLT+AG490 groups. Data were represented as the mean $\pm$ SD ($n = 8$, per group). * $P < 0.05$ vs. sham group. # $P < 0.05$ vs. the AOLT group.

plays a vital role in regulating the immune response. Western blot analysis was performed to demonstrate the effect of EA on JAK/STAT pathway proteins. JAK2, STAT1, and STAT3 proteins were mainly expressed in intestinal cells. The expressions of p-JAK2, p-STAT1, and p-STAT3 proteins significantly increased in intestines subjected to AOLT ($P < 0.05$ vs. the sham group). However, EA application decreased the expression of p-JAK2, p-STAT1, and p-STAT3 proteins ($P < 0.05$ vs. the AOLT group). Treatment with AG490 reduced the expressions of p-JAK2, p-STAT1, and p-STAT3 proteins ($P < 0.05$ vs. the AOLT group). These results demonstrated that EA inhibited the JAK/STAT signaling pathway in intestinal injury after AOLT. The JAK/STAT signaling pathway was active in the AOLT group (Figure 5).

3.5. EA Pretreatment Decreases the Expression of Cleaved Caspase-3 in the Intestine. From the immunohistochemical staining, intense staining for active Cleaved Caspase-3 in

the cytoplasm of epithelial cells of rats subjected to AOLT was observed. However, very few Cleaved Caspase-3 immunoreactive cells were observed in the sham and EA+sham groups. These results again showed the tissue-protective effect of EA against intestinal injury after AOLT (Figure 6). When pretreated with EA and AG490, the EA+sham and EA+AOLT+AG490 groups did not show a significant difference from the sham group.

3.6. EA Pretreatment Attenuates Intestinal Epithelial Cell Apoptosis. TUNEL assay was used to evaluate the apoptosis of tubular epithelial cells induced by AOLT. A large number of apoptotic epithelial cells were visible in the intestines subjected to AOLT ($P < 0.05$ vs. the sham group). EA application was associated with the occurrence of apoptosis of epithelial cells, which was less than that observed with the AOLT group ($P < 0.05$). In the EA+sham group, EA treatment had no effect on apoptotic epithelial cells, comparable with the sham

FIGURE 5: EA inhibited the phosphorylations of JAK2, STAT1, and STAT3. Representative Western blots for the phosphorylations of JAK2, STAT1, and STAT3 (a) of the intestine were detected after 6 h of AOLT; (A–E) The sham, AOLT, EA+AOLT, EA+sham, and EA+AOLT+AG490 groups. Densitometry analysis of Western blots for the ratio of p-JAK2/JAK2 (b), p-STAT1/STAT1 (c), and p-STAT3/STAT3 (d). Data were represented as the mean $\pm$ SD ($n = 8$, per group). * $P < 0.05$ vs. the sham group. # $P < 0.05$ vs. the AOLT group.

group. The EA+AOLT+AG490 group had similar results as the sham group (Figure 7).

4. Discussion

We found that AOLT-induced intestinal injury in rats. We determined whether EA affected the results. Pretreatment with EA alleviated intestinal injury after AOLT, as shown by HE staining and TUNEL methods. Compared with the sham group, the EA+sham group did not produce adverse reactions as normal rats. We explored the potential mechanisms underlying EA pretreatment. EA pretreatment inhibited the expressions of p-JAK2, p-STAT1, and p-STAT3 in intestines after AOLT. When treated with JAK2-specific inhibitor AG490, intestinal injury was balanced. This indicated that EA pretreatment alleviated intestinal injury after AOLT by inhibiting the JAK/STAT signaling pathway and provides basic evidence supporting its potential therapeutic efficacy (Figure 8).

AOLT blocks the return of venous blood in the lower extremities, which induces a systemic response and a release of harmful substances that may damage remote organs, including the intestines [21–24]. After AOLT in rats, blood flow is blocked, and intestinal injury undergoes two processes: ischemia and secondary reperfusion injury [20]. Due to intestinal hyperemia and congestion during liver transplantation, intestinal peristalsis and barriers are often impaired. Intestinal endogenous endotoxin and bacteria enter

the blood or lymphatic system and are transferred to other organs, which may lead to multiple organs dysfunction and systemic inflammatory response [25]. Intestinal injury is a complex, multifactorial, and pathophysiological process that involves dysfunction of bacterial translocation, absorption, and production of reactive oxygen species, cytokines, nitric oxide, and initiates multiorgan dysfunction syndrome [4, 26]. A recent study found that intestinal barrier destruction was widely observed during liver transplantation [27]. Other studies have also explored the possible mechanism of intestinal injury caused by liver transplantation. TLR4/NF- κ B signaling pathway activation-induced cell apoptosis was involved in intestinal injury during liver transplantation [28]. In our study, we found that the intestinal injury score increased after AOLT. Our previous study showed that AOLT induced remote organ injury after 6 h reperfusion, and the inflammatory response was obvious [29]. Therefore, in the present study, we selected 6 h after reperfusion to measure the intestinal injuries caused by AOLT.

Numerous strategies have been designed to reduce intestinal injury after AOLT [30]. Although some drugs have been successfully used to reduce intestinal injury in animal models, few can be used in clinical settings or may not be available during operations. Acupuncture has been used to control weight [31], reduce epilepsy [32], improve learning and memory disorders [33], and reduce pain [34–37]. EA is a complementary alternative medicine approach and involves applying an electrical current to acupuncture points.

FIGURE 6: The effect of EA on AOLT induced Cleaved Caspase-3 of intestinal cells (magnification, $\times 200$). Representative microphotographs were taken from the intestine of the sham (a), AOLT (b), EA+AOLT (c), EA+sham (d), and EA+AOLT+AG490 (e) groups at the time point of 6 h after AOLT. (f) Quantification of Cleaved Caspase-3 positive cells were counted in each group: (A–E) the sham, AOLT, EA+AOLT, EA+sham, and EA+AOLT+AG490 groups.

Zusanli is an important acupuncture point commonly used in acupuncture practices to promote blood circulation and can also cure peripheral soft tissue inflammation and ischemia-reperfusion [6, 38–41]. Indeed, several studies have demonstrated that EA significantly attenuated apoptosis in the brain, heart, and intestinal ischemia-reperfusion [42–44]. In our study, inflammatory reaction markers (TNF- α , HMGB-1) and oxidative stress factors (MDA) increased significantly in the AOLT group. However, in the EA+AOLT group, apoptosis of epithelial cells resulting from the intestinal injury was significantly decreased. Caspase-3, which participates in the apoptosis pathway in the intestinal mucosa, was also markedly reduced in the EA+AOLT group, which confirmed the decreased disruption of the structural integrity of intestinal mucosa, as observed by HE staining. This study evaluated the protective and antiapoptotic effects of EA on intestinal architecture and apoptosis in intestinal injury.

JAK/STAT is involved in many pathophysiological processes in the body [45, 46]. JAK2 is a member of the Janus kinase family and is involved in immune, hematopoietic,

neural, and other signal transduction systems [47]. A recent study showed that EA has a protective effect on focal cerebral ischemia by inhibiting JAK2 phosphorylated activation in rats [48]. AG490 is a specific inhibitor of JAK2 which can inhibit the activation of JAK2 and downregulate the phosphorylation of STATs. The JAK/STAT signaling pathway is involved in ischemia-reperfusion processes. Our previous study showed that the JAK/STAT signaling pathway was inhibited by propofol in the hippocampus [29]. To confirm the hypothesis that the JAK/STAT signaling pathway participated in regulating apoptotic process in our model, AG490 was given in the EA+AOLT group. Our study showed that EA+AG490 significantly improved intestinal apoptosis and reduced the expression of Cleaved Caspase-3 protein following AOLT. Our data also showed significantly decreased expressions of p-JAK2 in the EA and EA+AG490 groups, accompanied by the downregulation of STAT1 and STAT3 phosphorylation. Our results indicated that EA significantly protected intestinal injury after AOLT in rats by inhibiting the activation of the JAK/STAT signaling pathway.

FIGURE 7: The effect of EA on AOLT induced apoptosis of intestinal cells (magnification, $\times 200$). Representative microphotographs were taken from the intestine of the sham (a), AOLT (b), EA+AOLT (c), EA+sham (d), and EA+AOLT+AG490 (e) groups at the time point of 6 h after AOLT. Apoptosis was evaluated by TUNEL staining. Quantification of TUNEL-positive cells was counted following AOLT (f). * $P < 0.05$ vs. the sham group. # $P < 0.05$ vs. the AOLT group.

Taken together, intestinal injury contributed to the damage of intestinal function and histological structure and enhanced the apoptosis of intestinal epithelial cells and the expression of protein Cleaved Caspase-3. We demonstrated that EA pretreatment attenuated intestinal injury by inhibiting the JAK/STAT signaling pathway in a rat AOLT model. Zusanli is a classic and preferred acupoint

for treating gastrointestinal diseases and is widely used in clinical practice. Its therapeutic effect is recognized by many medical workers and patients. EA is a branch of acupuncture and moxibustion therapy. Compared with classic acupuncture, it has many advantages, such as good therapeutic effects, a wide range, ease of stimulation control, and continuous needle movements [49]. However, there

FIGURE 8: The possible mechanism of EA on AOLT induced apoptosis of intestine (a). The possible mechanism involved in the JAK/STAT pathway (b). EA pretreatment may have protective effects on AOLT-induced intestine injury by inhibiting JAK/STAT pathway.

are several limitations to our study. This study was based on our previous study; we could have used gene sequencing technology to explore the involved mechanism. Then, we would explore the mechanism in vivo and investigate whether this mechanism functioned at the cellular level. Finally, we need to apply this technique in clinical practice. To make progress on this front, our research team has begun a single-center study on the protective effect of EA pretreatment on postoperative organs.

In conclusion, our study showed that EA reduced intestinal injury against AOLT, at least in part through its inhibitory effects on the injury-induced activation of the JAK/STAT signaling pathway. If extrapolated to a clinical setting, EA has the potential to serve as a clinical strategy for preventing perioperative intestinal injury after AOLT.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

Authors' Contributions

Lili Jia designed and wrote this paper, Hongli Yu and Yiqi Weng provided the technical assistance, and Wenli Yu offered the guidance of this study.

Acknowledgments

The paper was financially supported by the Natural Science Foundation of Tianjin (18JCYBJC27500 and 17JCYBJC28000), the Tianjin Clinical Key Discipline Project (Anesthesiology), and 2019 Tianjin First Central Hospital Academy Project (2019CF09).

References

- [1] Y. Si, H. Bao, L. Han et al., "Dexmedetomidine protects against renal ischemia and reperfusion injury by inhibiting the JAK/STAT signaling activation," *Journal of Translational Medicine*, vol. 11, no. 1, p. 141, 2013.
- [2] J. Shen, G. Fu, L. Jiang, J. Xu, L. Li, and G. Fu, "Effect of dexmedetomidine pretreatment on lung injury following intestinal ischemia-reperfusion," *Experimental and Therapeutic Medicine*, vol. 6, no. 6, pp. 1359–1364, 2013.
- [3] X. K. Zhang, X. P. Zhou, Q. Zhang, and F. Zhu, "The preventive effects of dexmedetomidine against intestinal ischemia-reperfusion injury in Wistar rats," *Iranian Journal of Basic Medical Sciences*, vol. 18, no. 6, pp. 604–609, 2015.
- [4] C. Nastos, K. Kalimeris, N. Papoutsidakis et al., "Global consequences of liver ischemia/reperfusion injury," *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 906965, 13 pages, 2014.
- [5] E. Nozato, M. Shiraishi, T. Miyaguni et al., "In situ perfusion of the liver under portosystemic shunt in rats: a stable model of cold ischemia for orthotopic liver transplantation," *Transplantation Proceedings*, vol. 30, no. 7, pp. 3718–3720, 1998.
- [6] Y. Geng, D. Chen, J. Zhou, H. Jiang, and H. Zhang, "Role of cholinergic anti-inflammatory pathway in treatment of intestinal ischemia-reperfusion injury by electroacupuncture at Zusanli," *Evidence-based Complementary and Alternative Medicine*, vol. 2017, Article ID 6471984, 5 pages, 2017.
- [7] M. Cai, J. H. Lee, and E. J. Yang, "Electroacupuncture attenuates cognition impairment via anti-neuroinflammation in an Alzheimer's disease animal model," *Journal of Neuroinflammation*, vol. 16, no. 1, p. 264, 2019.
- [8] Y. Li, C. Yin, X. Li et al., "Electroacupuncture alleviates paclitaxel-induced peripheral neuropathic pain in rats via suppressing TLR4 signaling and TRPV1 upregulation in sensory neurons," *International Journal of Molecular Sciences*, vol. 20, no. 23, p. 5917, 2019.
- [9] N. Yun, J. W. Kang, and S. M. Lee, "Protective effects of chlorogenic acid against ischemia/reperfusion injury in rat liver: molecular evidence of its antioxidant and anti-inflammatory properties," *The Journal of Nutritional Biochemistry*, vol. 23, no. 10, pp. 1249–1255, 2012.
- [10] Y. Li, Y. Chen, X. Zhang et al., "Protective effect of electroacupuncture on liver ischemia-reperfusion injury in rats," *Experimental and Therapeutic Medicine*, vol. 16, no. 2, pp. 1373–1380, 2018.
- [11] M. V. R. Scognamiglio-Szabó, G. H. Bechara, S. H. Ferreira, and F. Q. Cunha, "Effect of various acupuncture treatment protocols upon sepsis in Wistar rats," *Annals of the New York Academy of Sciences*, vol. 1026, no. 1, pp. 251–256, 2004.
- [12] W. Geng, L. Cai, K. Han et al., "Electroacupuncture pretreatment alleviates cerebral ischemia-reperfusion injury by increasing GSK-3 β phosphorylation level via adenosine A1 receptor," *BioMed Research International*, vol. 2020, Article ID 6848450, 9 pages, 2020.
- [13] J. J. O'Shea, M. Gadina, and R. D. Schreiber, "Cytokine signaling in 2002," *Cell*, vol. 109, no. 2, pp. S121–S131, 2002.
- [14] S. Negoro, K. Kunisada, E. Tone et al., "Activation of JAK/STAT pathway transduces cytoprotective signal in rat acute myocardial infarction," *Cardiovascular Research*, vol. 47, no. 4, pp. 797–805, 2000.
- [15] A. Das, F. N. Salloum, D. Durrant, R. Ockaili, and R. C. Kukreja, "Rapamycin protects against myocardial ischemia-reperfusion injury through JAK2-STAT3 signaling pathway," *Journal of Molecular and Cellular Cardiology*, vol. 53, no. 6, pp. 858–869, 2012.
- [16] J. Q. Fang, J. Y. du, Y. Liang, and J. F. Fang, "Intervention of electroacupuncture on spinal p38 MAPK/ATF-2/VR-1 pathway in treating inflammatory pain induced by CFA in rats," *Molecular Pain*, vol. 9, 2013.
- [17] S. Katsuyama, H. Mizoguchi, H. Kuwahata et al., "Involvement of peripheral cannabinoid and opioid receptors in β -caryophyllene-induced antinociception," *European Journal of Pain*, vol. 17, no. 5, pp. 664–675, 2013.
- [18] J. Zhou, Y. Jin, R. Ma et al., "Electroacupuncture alleviates experimental chronic inflammatory pain by inhibiting calcium voltage-gated channel-mediated inflammation," *Evidence-based Complementary and Alternative Medicine*, vol. 2020, Article ID 7061972, 10 pages, 2020.
- [19] C. J. Chiu, A. H. McArdle, R. Brown, H. J. Scott, and F. N. Gurd, "Intestinal mucosal lesion in low-flow states. I. A morphological, hemodynamic, and metabolic reappraisal," *Archives of Surgery*, vol. 101, no. 4, pp. 478–483, 1970.
- [20] L. M. Gonzalez, A. J. Moeser, and A. T. Blikslager, "Animal models of ischemia-reperfusion-induced intestinal injury: progress and promise for translational research," *American Journal of Physiology. Gastrointestinal and Liver Physiology*, vol. 308, no. 2, pp. G63–G75, 2015.
- [21] B. Erenoglu, H. S. Gokturk, T. Kucukkartallar et al., "Mechanical intestinal cleansing and antibiotic prophylaxis for preventing bacterial translocation during the Pringle maneuver in rabbits," *Surgery Today*, vol. 41, no. 6, pp. 824–828, 2011.
- [22] D. L. Liu, B. Jeppsson, C. H. Hakansson, and R. Odselius, "Multiple-system organ damage resulting from prolonged hepatic inflow interruption," *Archives of Surgery*, vol. 131, no. 4, pp. 442–447, 1996.
- [23] P. A. Clavien, H. Petrowsky, M. L. DeOliveira, and R. Graf, "Strategies for safer liver surgery and partial liver transplantation," *The New England Journal of Medicine*, vol. 356, no. 15, pp. 1545–1559, 2007.
- [24] S. A. W. G. Dello, K. W. Reisinger, R. M. van Dam et al., "Total intermittent Pringle maneuver during liver resection can induce intestinal epithelial cell damage and endotoxemia," *PLoS One*, vol. 7, no. 1, article e30539, 2012.
- [25] M. Ge, X. Chi, A. Zhang et al., "Intestinal NF-E2-related factor-2 expression and antioxidant activity changes in rats undergoing orthotopic liver autotransplantation," *Oncology Letters*, vol. 6, no. 5, pp. 1307–1312, 2013.
- [26] D. L. Carden and D. N. Granger, "Pathophysiology of ischaemia-reperfusion injury," *The Journal of Pathology*, vol. 190, no. 3, pp. 255–266, 2000.
- [27] X. Tian, Z. Yang, F. Luo, and S. Zheng, "Gut microbial balance and liver transplantation: alteration, management, and prediction," *Frontiers in Medicine*, vol. 12, no. 2, pp. 123–129, 2018.
- [28] D.-D. Yuan, X.-J. Chi, Y. Jin et al., "Intestinal injury following liver transplantation was mediated by TLR4/NF- κ B activation-

- induced cell apoptosis," *Molecular Medicine Reports*, vol. 13, no. 2, pp. 1525–1532, 2016.
- [29] L. Jia, F. Wang, X. Gu et al., "Propofol postconditioning attenuates hippocampus ischemia-reperfusion injury via modulating JAK2/STAT3 pathway in rats after autogenous orthotopic liver transplantation," *Brain Research*, vol. 1657, no. 15, pp. 202–207, 2017.
- [30] R. Bahde and H. U. Spiegel, "Hepatic ischaemia-reperfusion injury from bench to bedside," *The British Journal of Surgery*, vol. 97, no. 10, pp. 1461–1475, 2010.
- [31] Y. Li, P. Adamek, H. Zhang et al., "The cancer chemotherapeutic paclitaxel increases human and rodent sensory neuron responses to TRPV1 by activation of TLR4," *The Journal of Neuroscience*, vol. 35, no. 39, pp. 13487–13500, 2015.
- [32] P. Flood, J. M. Sonner, D. Gong, and K. M. Coates, "Isoflurane hyperalgesia is modulated by nicotinic inhibition," *Anesthesiology*, vol. 97, no. 1, pp. 192–198, 2002.
- [33] P. M. Cornett, J. A. Matta, and G. P. Ahern, "General anesthetics sensitize the capsaicin receptor transient receptor potential V1," *Molecular Pharmacology*, vol. 74, no. 5, pp. 1261–1268, 2008.
- [34] S. J. Raithel, M. R. Sapio, M. J. Iadarola, and A. J. Mannes, "Thermal A- δ nociceptors, identified by transcriptomics, express higher levels of anesthesia-sensitive receptors than thermal C-fibers and are more suppressible by low-dose isoflurane," *Anesthesia and Analgesia*, vol. 127, no. 1, pp. 263–266, 2018.
- [35] M. J. Caterina, M. A. Schumacher, M. Tominaga, T. A. Rosen, J. D. Levine, and D. Julius, "The capsaicin receptor: a heat-activated ion channel in the pain pathway," *Nature*, vol. 389, no. 6653, pp. 816–824, 1997.
- [36] C.-Y. Chen, C.-N. Lin, R.-S. Chern, Y.-C. Tsai, Y.-H. Chang, and C.-H. Chien, "Neuronal activity stimulated by liquid substrates injection at zusanli (ST36) acupoint: the possible mechanism of aquapuncture," *Evidence-based Complementary and Alternative Medicine*, vol. 2014, Article ID 627342, 7 pages, 2014.
- [37] S. I. Choi, J. Y. Lim, S. Yoo, H. Kim, and S. W. Hwang, "Emerging role of spinal cord TRPV1 in pain exacerbation," *Neural Plasticity*, vol. 2016, Article ID 5954890, 10 pages, 2016.
- [38] A. Villegas-Bastida, R. Torres-Rosas, L. A. Arriaga-Pizano, J. Flores-Estrada, A. Gustavo-Acosta, and M. A. Moreno-Eutimio, "Electrical stimulation at the ST36 acupoint protects against sepsis lethality and reduces serum TNF levels through vagus nerve- and catecholamine-dependent mechanisms," *Evidence-based Complementary and Alternative Medicine*, vol. 2014, Article ID 451674, 8 pages, 2014.
- [39] S. Hu, M. H. du, H. M. Luo et al., "Electroacupuncture at Zusanli (ST36) prevents intestinal barrier and remote organ dysfunction following gut ischemia through activating the cholinergic anti-inflammatory-dependent mechanism," *Evidence-based Complementary and Alternative Medicine*, vol. 2013, Article ID 592127, 10 pages, 2013.
- [40] M.-H. Du, H. M. Luo, Y. J. Tian et al., "Electroacupuncture ST36 prevents postoperative intra-abdominal adhesions formation," *The Journal of Surgical Research*, vol. 195, no. 1, pp. 89–98, 2015.
- [41] L. Zhang, H. Wang, Z. Huang et al., "Inhibiting effect of electroacupuncture at Zusanli on early inflammatory factor levels formed by postoperative abdominal adhesions," *Evidence-based Complementary and Alternative Medicine*, vol. 2014, Article ID 950326, 5 pages, 2014.
- [42] Y. Geng, D. Chen, J. Zhou et al., "Synergistic effects of electroacupuncture and mesenchymal stem cells on intestinal ischemia/reperfusion injury in rats," *Inflammation*, vol. 39, no. 4, pp. 1414–1420, 2016.
- [43] R. Torres-Rosas, G. Yehia, G. Peña et al., "Dopamine mediates vagal modulation of the immune system by electroacupuncture," *Nature Medicine*, vol. 20, no. 3, pp. 291–295, 2014.
- [44] D. Grech, Z. Li, P. Morcillo et al., "Intraoperative low-frequency electroacupuncture under general anesthesia improves postoperative recovery in a randomized trial," *Journal of Acupuncture and Meridian Studies*, vol. 9, no. 5, pp. 234–241, 2016.
- [45] J. J. O'Shea, M. Gadina, and R. D. Schreiber, "Cytokine signaling in 2002: new surprises in the Jak/Stat pathway," *Cell*, vol. 109, no. 2, pp. S121–S131, 2002.
- [46] K. Imada and W. J. Leonard, "The JAK-STAT pathway," *Molecular Immunology*, vol. 37, no. 1-2, pp. 1–11, 2000.
- [47] L. A. Wang, H. Bi, H. R. Ji, Y. P. Wang, and Z. M. Du, "Effect of lumbrokinase on JAK-STAT pathway in rat brain during cerebral ischemia and reperfusion," *Zhongguo Yao Xue Za Zhi*, vol. 44, no. 24, pp. 1862–1865, 2009.
- [48] R. Liu, N. Xu, W. Yi, K. Huang, and M. Su, "Electroacupuncture effect on neurological behavior and tyrosine kinase-JAK 2 in rats with focal cerebral ischemia," *Journal of Traditional Chinese Medicine*, vol. 32, no. 3, pp. 465–470, 2012.
- [49] Y. X. Gen, D. Cheng, J. Lu, M. Q. Chen, H. D. Zhang, and J. Zhou, "Effects of electroacupuncture at Zusanli (ST36) on mucosal barrier repair in rats with intestinal ischemia reperfusion injury," *Journal of Emergency in Traditional Chinese Medicine*, vol. 25, no. 7, pp. 1300–1302, 2016.