Research Article

Folic Acid Protects Melanocytes from Oxidative Stress via Activation of Nrf2 and Inhibition of HMGB1

Pengran Du , Shaolong Zhang , Shuli Li , Yuqi Yang , Pan Kang , Jiayi Chen ,
Tianwen Gao, Chunying Li, Qian Zhang , and Weigang Zhang

Department of Dermatology, Xijing Hospital, Fourth Military Medical University, No. 127 of West Changle Road, Xi'an, Shaanxi 710032, China

Correspondence should be addressed to Qian Zhang; qian815032543@126.com and Weigang Zhang; zhangwg@fmmu.edu.cn

Received 22 September 2021; Revised 13 November 2021; Accepted 13 November 2021; Published 7 December 2021

Academic Editor: Zhouwei Wu

Copyright © 2021 Pengran Du et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Vitiligo is a cutaneous depigmentation disease due to loss of epidermal melanocytes. Accumulating evidence has indicated that oxidative stress plays a vital role in vitiligo via directly destructing melanocytes and triggering inflammatory response that ultimately undermines melanocytes. Folic acid (FA), an oxidized form of folate with high bioavailability, exhibits potent antioxidant properties and shows therapeutic potential in multiple oxidative stress-related diseases. However, whether FA safeguards melanocytes from oxidative damages remains unknown. In this study, we first found that FA relieved melanocytes from H_2O_2 -induced abnormal growth and apoptosis. Furthermore, FA enhanced the activity of antioxidative enzymes and remarkably reduced intracellular ROS levels in melanocytes. Subsequently, FA effectively activated nuclear factor E2-related factor 2 (Nrf2) pathway, and Nrf2 knockdown blocked the protective effects of FA on H_2O_2 -treated melanocytes. Additionally, FA inhibited the production of proinflammatory HMGB1 in melanocytes under oxidative stress. Taken together, our findings support the protective effects of FA on human melanocytes against oxidative injury via the activation of Nrf2 and the inhibition of HMGB1, thus indicating FA as a potential therapeutic agent for the treatment of vitiligo.

1. Introduction

Vitiligo is an acquired depigmentation dermatosis with an estimated prevalence of 1% worldwide [1]. To date, the reason for the loss of epidermal melanocytes as the key event in vitiligo is still unclear, which hinders the development of effective therapeutic strategies [2]. Nevertheless, recent studies have indicated the involvement of oxidative stress in the pathogenesis of vitiligo [3]. Along with the deficiency of antioxidant system of melanocytes, especially nuclear factor E2-related factor 2 (Nrf2) pathway that is a master regulator of antioxidative response [4, 5], reactive oxygen species (ROS) such as hydrogen peroxide (H_2O_2) is excessively accumulated in vitiligo [6]. Accordingly, melanocytes are susceptible to ROS-induced apoptosis in vitiligo [3, 5]. Moreover, ROS facilitates the release of high-mobility group protein B1 (HMGB1) that belongs to damage-associated molecular patterns (DAMPs) with strong proinflammatory effects from

melanocytes [7–10]. HMGB1 subsequently causes the production of chemokines in adjacent keratinocytes and the maturation of dendritic cells (DCs) in a paracrine way, which ultimately promotes the formation of cytotoxic T cells (CTLs) that undermine melanocytes in vitiligo [7]. Therefore, oxidative stress plays a vital role in vitiligo via not only direct oxidative damages on melanocytes but also triggering cutaneous T cell response that targets melanocytes.

Folic acid (FA), a synthetic oxidized form of folate with high bioavailability, is well known for its protective effect against neural tube defects [11]. FA digested at intestine is ultimately converted to 5-methyltetrahydrofolate with the help of methylenetetrahydrofolate reductase (MTHFR) and then involved in the remethylation of homocysteine (Hcy) to methionine [12]. In 2014, our group reported an association between single-nucleotide polymorphisms (SNPs) in *MTHFR* gene and vitiligo susceptibility [13]. Furthermore, Hcy was observed to accumulate in vitiligo and induce the

apoptosis of melanocytes via endoplasmic reticulum stress [14]. These findings indicate that the disruption of folate metabolism plays a role in the pathogenesis of vitiligo. Notably, some recent studies have found that FA possesses excellent antioxidant property and is effective in maintaining cellular redox status [15, 16]. However, the effect of FA on melanocytes under oxidative stress in vitiligo has not been investigated before.

Herein, we initially observed that FA did protect melanocytes from oxidative damages. Furthermore, the influence of FA on the antioxidant response of melanocytes was evaluated. In addition, the effect of FA on the status of HMGB1 in melanocytes under oxidative stress was also investigated in the present study.

2. Materials and Methods

2.1. Cell Culture and Treatment. The immortalized normal human epidermal melanocyte cell line PIG1 (a gift from Dr. Caroline Le Poole, Loyola University, Chicago, USA) was maintained in Medium 254 (M254500, Gibco, USA), supplemented with human melanocyte growth supplement (Gibco) and 5% fetal bovine serum at 37°C amid 5% CO₂. Oxidative stress was induced by adding H₂O₂ (7722841, Sigma-Aldrich, USA) at 800 or 500 μ M into the culture medium for 48 h. To determine the protective effect of FA (F8758, Sigma-Aldrich, USA) on melanocytes against oxidative damages, FA and H₂O₂ were added simultaneously into the culture medium of PIG1 cells at indicated concentrations.

2.2. CCK8 Assay. Cell viability was evaluated by using a CCK8 kit (C008, 7Sea biotech, China) according to manufacturer's manual. Generally, PIG1 cells were seeded into 96-well plates at the density of 2×10^4 cells per well before further treatments as indicated. Next, 10 μ l of CCK8 solution was added to each well, and the cells were further incubated at 37°C for 2 h. The optical density (OD) was measured at 450 nm by Model 680 Microplate Reader (BioRad, USA). All experiments were performed in triplicate.

2.3. Apoptosis Assay. Cell apoptosis was measured using Annexin V-FITC/PI kit (A005, 7 Seabiotec, China). Briefly, the treated PIG1 cells were stained according to manufacturer's instructions and then detected by flow cytometry (Beckman Coulter, USA) and analyzed with Expo32 software (BeckmanCoulter, USA).

2.4. Western Blot Assay. Whole cell lysates were extracted with RIPA Buffer (P0013C, Beyotime, China). Nuclear/cytoplasmic fractionation was separated using the Nuclear and Cytoplasmic Protein Extraction Kit (P0028, Beyotime, China) according to manufacturer's instructions. Protein concentration was measured using the BCA Protein Assay kit (23225, ThermoScientific, USA). Equal amounts of protein were separated by SDS-PAGE (Bio-Rad, USA) and transferred to polyvinylidene difluoride membranes (Millipore, USA). Membranes were blocked in 5% nonfat milk for 2 h and incubated overnight at 4°C in primary antibodies diluted with PBS containing 1% BSA, including mouse anti-

human β -actin (8H10D10) (1:5000, 3700, Cell Signaling Technology, USA), rabbit anti-human LaminA/C (1:1000, 10298-1-AP, Proteintech, China), rabbit anti-human Tubulin (1:2000, 11224-1-AP, Proteintech, China), rabbit anti-human Bcl-2 (D55G8) (1:1000, 4223, Cell Signaling Technology, USA), rabbit anti-human Bax (D2E11) (1:1000, 5023, Cell Signaling Technology, USA), rabbit anti-human cleaved Caspase-3 (Asp175) (5A1E) (1:1000, 9664, Cell Signaling Technology, USA), rabbit anti-human Caspase3 (1:1000, Lot. 9662, Cell Signaling Technology), rabbit anti-human Nrf2 (1:500, ab62352, Abcam, USA), rabbit anti-human Nrf2 (phospho S40) (1:5000, ab76026, Abcam, USA), mouse anti-human HO-1 (1:500, ab13248, Abcam, USA), rabbit anti-human SOD2 (1:2000, ab13533, Abcam, USA), and rabbit anti-human HMGB1 (1:500, ab18256, Abcam, USA). After washing, the membranes were incubated at room temperature for 2 h in corresponding peroxidase-conjugated secondary antibodies diluted with PBS containing 1% BSA, including goat anti-rabbit IgG antibody (1:5000, AP132P, Sigma-Aldrich, United States) and goat anti-mouse IgG antibody (1:5000, AP124P, Sigma-Aldrich, United States). At last, the bands were detected by an enhanced chemiluminescence reagent (Millipore, USA) under Western blotting detection system (Bio-Rad, USA).

2.5. SOD Activity Assay. After cells were lysed, the total protein was extracted to detect the activity of SOD by using Total Superoxide Dismutase Assay Kit with WST-8 (S0101M, Beyotime, China) following manufacturer's instructions.

2.6. Measurement of Intracellular ROS. The intracellular ROS was measured using a fluorogenic probe for ROS (CM-H2DCFDA) (C6827, Invitrogen, USA) following the protocol reported previously [17]. Briefly, PIG1 cells were seeded into 6-well plates with a density of 5×10^5 cells. After indicated treatments, 10 μ M of DCFH-DA was added for 30 min, and then, cells were collected for detection of fluorescence intensity of DCF via flow cytometry (Beckman Coulter, USA).

2.7. Immunofluorescence Assay. PIG1 cells were grown in single-layer glass slides (801002, NEST Biotechnology, China) at a density of 5000 cells per dish. After indicated treatments, cells were washed by PBS and fixed with 4% paraformaldehyde. Cells were permeabilized for 15 min in PBS supplemented with 0.1% Triton X-100 at room temperature and labeled with the primary antibody anti-Nrf2 (1:200, ab62352, Abcam, USA) at 4°C overnight and corresponding secondary antibody Cy3-tagged goat anti-rabbit IgG (1:1000, ab6939, Abcam, USA) for 1 h at room temperature. At last, cells were incubated with the nuclear dye 40, 6-diamidino-2-phenylindole (DAPI) (1:1000, 62247, Thermo Fisher Scientific, USA) for 10 min at room temperature in the dark. The fluorescence was detected by using FV-1000/ES laser confocal microscopy (Olympus, Japan).

2.8. Enzyme-Linked Immunosorbent Assay (ELISA). ELISA was performed using the Human HMGB1 ELISA Kit (Shino-Test, Japan) according to manufacturer's

instructions. The absorbance (A450) was measured with a plate reader (Bio-Rad).

2.9. Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR). Total RNA was extracted using Trizol Reagent (15596018, Invitrogen, United States) and then reversely transcribed to cDNA by a PrimeScript RT reagent kit (AK4301, TaKaRa, Japan). The qRT-PCR assay was performed using SYBR PremixEx Taq II (AKA1008, TaKaRa, Japan) with the real-time PCR Detection System (Bio-Rad, United States). The relative mRNA expression was normalized to the β -actin gene. The primers used in this study were as follows: HMGB1, forward: AGCCCTCTTCATGTTC CGAAGTGT, reverse: TCATGTCAACACCTGCAGTCC CTT; β -actin, forward: TCATGAAGTGTGACGTGGACA TC, reverse: CAGGAGGAGCAATGATCTTGATCT.

2.10. RNA Interference. PIG1 cells were seeded at 2×10^5 cells per well for 24 h before transfection. Cells were transfected with Nrf2 shRNA or irrelevant shRNA control (GenePharma, China) with Lipofectamine 3000 (Invitrogen) following manufacturer's protocol. The sequences of the shRNAs used in the present study were as follows: shRNA-Nrf2: 5'-GGTTGCCACATTCCCAAATC-3'; shRNA-Control: 5'-TTCTCCGAACGTGTCACGT-3'.

2.11. Statistical Analysis. Data analysis was performed using GraphPad Prism version 6.0 software (GraphPad Software, San Diego, CA). The two-tailed Student's *t*-test or one-way analysis of variance (ANOVA) was used in our analyses. *P* values less than 0.05 were considered significant. Data represent as mean $\pm$ SD for at least three independent experiments.

3. Results

3.1. FA Protects Melanocytes from Oxidative Stress. CCK8 assay was initially performed to evaluate suitable doses of FA for use in subsequent experiments. As a result, FA at the concentrations of 10 to 500 μ M had no significant influence on the growth of PIG1 cells over an incubation course of three days, but high doses (1000 and 3000 μ M) of FA showed toxicity on the third day (Figure 1(a)). To investigate the protective effect of FA on melanocytes under oxidative stress, PIG1 cells were treated with FA at a concentration gradient of 25, 50, and 100 μ M, respectively, and costimulated with 800 μ M H_2O_2 for 24 h. It turned out that H_2O_2 -induced morphologic changes of shortened or disappeared dendrites in PIG1 cells were significantly rescued by the pretreatment with FA at relatively higher doses of 50 or 100 μ M (Figure 1(b)). Consistently, FA reversed the inhibited viability of PIG1 cells caused by H_2O_2 in a dose-dependent manner (Figure 1(c)).

We went on to investigate whether FA protects melanocytes from apoptosis under oxidative stress. Flow cytometry analysis showed that the proportion of apoptotic PIG1 cells was markedly increased after the treatment with H_2O_2 , whereas the cotreatment with FA at the concentrations of 50 or 100 μ M significantly rescued PIG1 cells from H_2O_2 -

induced apoptosis (Figures 2(a) and 2(b)). Moreover, the upregulated protein levels of proapoptotic Bax and cleaved Caspase-3 and the downregulated protein level of Bcl-2 caused by H_2O_2 were all reversed by cotreated FA (100 μ M) in PIG1 cells (Figure 2(c)). Altogether, these results support that FA protects melanocytes from oxidative damages.

3.2. FA Potentiates Antioxidant Response in Melanocytes under Oxidative Stress. Next, the influence of FA on the antioxidant system of melanocytes was evaluated. The accumulation of ROS in PIG1 cells treated with H_2O_2 was attenuated by the cotreatment with FA at the concentrations of 50 μ M or 100 μ M (Figures 3(a) and 3(b)). Meanwhile, the repressed activity of superoxide dismutase (SOD), a key antioxidant enzyme that scavenges ROS [17, 18], was significantly reversed by FA at 100 μ M in PIG1 cells treated with H_2O_2 (Figure 3(c)).

Since Nrf2 plays a central role in the antioxidant system of melanocytes [3, 5], we then evaluated the effect of FA on the activation of Nrf2 pathway. As shown by our Western blot analysis, the cotreatment of FA promotes the expressions of Nrf2, phosphorylated Nrf2 (p-Nrf2) that acts as a transcriptional factor as well as SOD2 and heme oxygenase-1 (HO-1) that are antioxidant proteins transcriptionally regulated by p-Nrf2 in PIG1 cells treated with H_2O_2 (Figure 3(d)). Subsequent detection of nuclear and cytosolic Nrf2 separately disclosed that FA induced the translocation of Nrf2 from cytoplasm to nucleus in H_2O_2 -treated PIG1 cells (Figure 3(e)), which was further supported by immunofluorescence assay (Figure 3(f)). Collectively, our findings indicate that FA enhances antioxidant response in melanocytes under oxidative stress.

3.3. FA Prevents Melanocytes from Oxidative Damages via Activating Nrf2 Pathway. To decide whether the protective effect of FA on melanocytes under oxidative stress is dependent on the activation of Nrf2, Nrf2 was silenced by transfecting PIG1 cells with short hairpin RNA (shRNA) of Nrf2 (interference efficiency shown in Figure 4(a)). It was found that FA failed to eliminate intracellular ROS in H_2O_2 -treated PIG1 cells as long as Nrf2 was deficient (Figures 4(b) and 4(c)). Moreover, Nrf2 knockdown abolished the protection of FA against H_2O_2 -induced apoptosis on PIG1 cells (Figures 4(d) and 4(e)). Further Western blot assay observed that Nrf2 knockdown abrogated FA-induced upregulation of HO-1 and SOD2 in H_2O_2 -treated PIG1 cells (Figure 4(f)). Altogether, these findings demonstrate that the activation of Nrf2 pathway mediates the protection of FA on melanocytes against oxidative damages.

3.4. FA Reduces the Release of HMGB1 from Melanocytes under Oxidative Stress. Given the proinflammatory role of HMGB1 secreted by melanocytes under oxidative stress in the immune pathogenesis of vitiligo as described before, the effect of FA on the status of HMGB1 in ROS-triggered melanocytes was examined at last. Consistent with our previous report [7], H_2O_2 prominently increased the mRNA and protein levels of HMGB1 in PIG1 cells; both of which,

FIGURE 1: FA attenuated H_2O_2 -induced oxidative damages in PIG1 cells. (a) The growth of PIG1 cells treated with different concentrations of FA over three days was determined by CCK8 assay. (b, c) PIG cells were treated with different concentrations of FA and 800 μM H_2O_2 simultaneously for 48 h. The morphological features of melanocytes were detected by microscope (b). Each field shown is a representative image of at least nine similar fields from three independent experiments. Scale bar = 200 μm . Cell viability was determined by CCK8 assay (c). All data are presented as the mean $\pm$ SD across three independent experiments. * $P < 0.05$, ** $P < 0.01$; ns: not significant.

however, were repressed by the addition of FA into the culture (Figures 5(a) and 5(b)). Additionally, the cotreatment with FA successfully prevented the release of HMGB1 from H_2O_2 -treated PIG1 cells (Figure 5(c)). Therefore, FA is of promising potential in inhibiting oxidative stress-induced inflammatory response mediated by HMGB1 in vitiligo.

4. Discussion

Oxidative stress is convincingly a key pathogenetic factor of vitiligo and contributes to its onset and progression [19]. Excessive ROS production and insufficient antioxidant response jointly lead to the destruction of melanocytes in

FIGURE 2: FA protected PIG1 cells from H_2O_2 -induced apoptosis. PIG1 cells were exposed to 800 μ M H_2O_2 and FA at indicated concentrations for 48 h simultaneously. (a, b) The percentage of apoptotic cells was determined by flow cytometry assay. Bar graphs represent the mean values of flow cytometry data. (c) Effects of FA on H_2O_2 -induced expressions of Bcl-2, Bax, Caspase-3, and cleaved Caspase-3 were determined via Western blot. β -Actin was detected as loading control. Bar graphs represent the quantification of Bcl-2/Bax ratio and cleaved/full length Caspase-3 ratio via gray intensity analysis across three replicate experiments. All data are presented as the mean $\pm$ SD across three independent experiments. * P < 0.05, *** P < 0.001; ns: not significant.

FIGURE 3: FA potentiated the antioxidant response in H_2O_2 -treated PIG1 cells. PIG1 cells were exposed to 800 μM H_2O_2 and FA at indicated concentrations for 48 h simultaneously. (a, b) Intracellular ROS level was determined by flow cytometry assay. Bar graphs represent the mean values of the fluorescence intensity of ROS level. (c) The activity of antioxidant enzyme SOD was examined via using a specialized kit. (d) Effects of FA on H_2O_2 -induced expressions of Nrf2, p-Nrf2, SOD2, and HO-1 were determined via Western blot. β -Actin was detected as loading control. (e, f) The nuclear/cytoplasmic distribution of Nrf2 was detected via Western blot (e) and immunofluorescence (f), respectively. Lamin A/C and tubulin were detected as loading controls in western blot assay. Nrf2 was stained with Cy3 (red) and nuclei were counterstained with DAPI (blue) in fluorescence assay. Scale bar = 10 μm . All data are presented as the mean $\pm$ SD across three independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; ns: not significant.

FIGURE 4: Continued.

FIGURE 4: Nrf2 knockdown abolished the protective effects of FA on PIG1 cells against oxidative damages induced by H_2O_2 . PIG1 cells were transfected with the shRNA against Nrf2 or control shRNA for 24 h and then treated with H_2O_2 and FA at indicated concentrations for 48 h. (a) The interference efficiency of Nrf2 shRNA was evaluated via Western blot. β -Actin was detected as loading control. (b, c) Intracellular ROS level was determined by flow cytometry assay. Bar graphs represent the mean values of the fluorescence intensity of ROS level. (d, e) The percentage of apoptotic cells was determined by flow cytometry assay. Bar graphs represent the mean values of flow cytometry data. (f) The expressions of Nrf2, p-Nrf2, SOD2, and HO-1 were determined via Western blot. β -Actin was detected as loading control. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; ns: not significant.

vitiligo [18], which prompts us to seek for measures that rebalance the redox homeostasis of melanocytes. Previous studies have shown that FA keeps the survival of astrocytes and purkinje neurons by decreasing ROS level [20, 21]. Meanwhile, rats treated with FA could obtain high levels of

antioxidative enzymes like SOD and low ROS levels [16], indicating that FA potentiates antioxidant system both *in vitro* and *in vivo*. The current study further showed that FA was able to protect melanocytes from oxidative injury by lowering intracellular ROS levels and upregulating HO-

FIGURE 5: FA inhibited the production of HMGB1 in PIG1 cells treated with H₂O₂. (a) The mRNA level of HMGB1 normalized to β -actin in PIG1 cells treated with H₂O₂ and FA at indicated concentrations for 24 h was determined via qRT-PCR. (b) The protein level of HMGB1 in PIG1 cells treated with H₂O₂ and FA at indicated concentrations for 48 h was determined via Western blot. β -Actin was detected as loading control. (c) The secretion level of HMGB1 from PIG1 cells treated with H₂O₂ and FA at indicated concentrations for 72 h was determined via ELISA. All data are presented as the mean $\pm$ SD across three independent experiments. * $P < 0.05$.

1 and SOD2, both of which were reported deficient in vitiligo melanocytes [3, 5]. Further studies using H₂O₂-induced vitiligo animal model [22] are needed to confirm the protective effect of FA on melanocytes from H₂O₂-triggered oxidation *in vivo*.

Apoptosis is the main form of cell death for melanocytes under oxidative stress [18]. Bcl-2 family proteins including prosurvival Bcl-2 and proapoptotic BAX are the key regulators of apoptosis pathway, in which Caspase-3 acts as a classical executor protein [23]. Herein, FA exhibited potent antiapoptotic ability in human melanocytes under oxidative stress. We previously found that FA protected melanocytes from Hcy-induced apoptosis [14]. Interestingly, Li et al. and Zhou et al. observed that FA inhibited aging-induced apoptosis in astrocytes and neurocytes, respectively, suggesting that FA could suppress apoptosis in various conditions [15, 24]. Additionally, our study showed the upregulation of Bax, the downregulation of Bcl2, and the inhibition of Caspase-3 activation in melanocytes undergoing oxidative stress following FA treatment, indicating that the antiapoptotic capacity of FA may result from decreasing cleaved Caspase-3 through modulating the ratio of Bax to Bcl-2 expression in oxidative injured melanocytes.

Nrf2 is a vital antioxidant regulator that functions as a potent transcriptional activator [4, 25]. Under physiological conditions, Nrf2 is localized primarily in the cytoplasm and bound by Kelch-like ECH-associated protein 1 (Keap1)

that hinders the activation of Nrf2 [26]. In response to oxidative stress, Nrf2 is disengaged from its binding to Keap1 and translocated into the nucleus, where Nrf2 potentiates the transcription of antioxidant response elements (AREs) and induces a battery of antioxidant proteins [27]. Nrf2 pathway plays a crucial role in protecting human melanocytes from oxidative stress as demonstrated by our previous studies [4, 5, 28, 29]. Melanocytes in vitiligo are deficient of the activity of Nrf2 and thus more vulnerable to oxidative stress, and reintroduction of Nrf2 pathway is expectedly accompanied with improved survival of melanocytes under oxidative stress [5]. The activation of Nrf2 pathway is characterized by the phosphorylation of Nrf2, the nuclear translocation of Nrf2, and the expression of antioxidant proteins including SOD2 and HO-1 [30], all of which were observed in FA-treated melanocytes by our study. Further knockdown of Nrf2 abolished the protection of FA on melanocytes against oxidative damages, supporting that the antioxidant effect of FA is dependent on Nrf2 pathway in melanocytes. In parallel with our results, Cao et al. found that FA promoted the translocation of Nrf2 from cytoplasm to nucleus and elevated HO-1 expression in rats with spontaneous hypertension [31]. Consistently, the deficiency of folate in fish gills was accompanied with decreased expression of Nrf2 [32]. All of these findings indicate FA as a strong activator of Nrf2 that can be applied extensively.

Apart from directly killing melanocytes, oxidative stress is known to ignite inflammatory response that ultimately undermines melanocytes in vitiligo [3]. One of the most important mediators of inflammation in vitiligo is HMGB1 released from melanocytes primed by ROS [7]. As a proinflammatory DAMP molecule, HMGB1 acts on its receptors, including several toll-like receptors and receptor for advanced glycation end products (RAGE), and initiates the second wave of protein kinase cascade activation, which generates a positive feedback mechanism to maintain and amplify inflammatory response [33]. Accordingly, the therapies targeting HMGB1 directly or disrupting its binding to proinflammatory receptors may be effective in alleviating inflammation [34]. Sun et al. found that FA-derived drugs disrupted the binding of HMGB1 to receptors and suppressed HMGB1-induced TNF release in human macrophages [35]. Beyond this previous report, our study showed that FA dramatically reduced the mRNA level, protein level, and secretion level of HMGB1 in H₂O₂-treated melanocytes, implicating that FA could inhibit oxidative stress-triggered cutaneous inflammation mediated by HMGB1 in vitiligo. In addition, Dong et al. has demonstrated that oxidative damages mediated by HMGB1 could be alleviated with the activation of Nrf2 that further inhibits the expression of HMGB1 [36]. Therefore, the FA-mediated inhibition of HMGB1 in melanocytes observed in the present study was probably mediated by Nrf2 pathway activated by FA.

The serum level of folate is generally normal in vitiligo patients [37], but recent clinical investigations have confirmed the deficiency of folate in the patients given the phototherapy of narrow-band ultraviolet B (NB-UVB) [38], which is one of the most effective treatments currently applied to vitiligo [39]. Accordingly, taking supplementary FA has been recommended for vitiligo patients on phototherapy, especially for women of childbearing age [40]. Although the results of previous studies are mixed regards to the effect of FA supplementation in vitiligo patients [41], the present study demonstrates FA as an effective agent that protects melanocytes from oxidative damages via its antioxidant and anti-inflammatory capability. Further clinical studies with larger sample size are thus encouraged to evaluate the therapeutic potential of FA for vitiligo.

Abbreviations

Nrf2:	Nuclear factor E2-related factor 2
ROS:	Reactive oxygen species
HMGB1:	High-mobility group protein B1
DAMP:	Damage-associated molecular pattern
DC:	Dendritic cell
CTL:	Cytotoxic T cell
FA:	Folic acid
MTHFR:	Methylenetetrahydrofolate reductase
Hcy:	Homocysteine
SNP:	Single-nucleotide polymorphism
p-Nrf2:	Phosphorylated Nrf2
SOD2:	Superoxide dismutase 2
HO-1:	Heme oxygenase-1

shRNA:	Short hairpin RNA
Keap1:	Kelch-like ECH-associated protein 1
ARE:	Antioxidant response element
RAGE:	Receptor for advanced glycation end products
NB-UVB:	Narrow-band ultraviolet B
Bax:	B cell lymphoma 2-associated X protein
Bcl-2:	B cell lymphoma-2
OD:	Optical density
H ₂ O ₂ :	Hydrogen peroxide
PI:	Propidium iodide
SDS-PAGE:	Sodium dodecyl sulfate-poly-acrylamide gel electrophoresis
PBS:	Phosphate buffer solution
DAPI, 4':	6-Diamidino-2-phenylindole
ANOVA:	One-way analysis of variance.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflict of interest.

Authors' Contributions

PD, SZ, and WZ conceived the project and designed the experiments. PD, SZ, and YY performed the experiments. SL, YY, PK, and JC analyzed the data. PD, SL, and PK prepared the manuscript. WZ, QZ, TG, and CL revised the manuscript. All authors read and approved the final manuscript. Pengran Du and Shaolong Zhang contributed equally to this work.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (nos. 81903207, 81903209, 81930087, 91742201, and 81625020).

References

- [1] M. L. Frisoli, K. Essien, and J. E. Harris, "Vitiligo: mechanisms of pathogenesis and treatment," *Annual Review of Immunology*, vol. 38, no. 1, pp. 621–648, 2020.
- [2] B. Khaitan and T. Sindhuja, "Autoimmunity in vitiligo: therapeutic implications and opportunities," *Autoimmunity Reviews*, no. article 102932, 2021.
- [3] J. Chen, S. Li, and C. Li, "Mechanisms of melanocyte death in vitiligo," *Medicinal Research Reviews*, vol. 41, no. 2, pp. 1138–1166, 2021.
- [4] Z. Jian, K. Li, L. Liu et al., "Heme oxygenase-1 protects human melanocytes from H₂O₂-induced oxidative stress via the Nrf2-ARE pathway," *The Journal of Investigative Dermatology*, vol. 131, no. 7, pp. 1420–1427, 2011.
- [5] Z. Jian, K. Li, P. Song et al., "Impaired activation of the Nrf2-ARE signaling pathway undermines H₂O₂-induced oxidative stress response: a possible mechanism for melanocyte

- degeneration in vitiligo,” *The Journal of Investigative Dermatology*, vol. 134, no. 8, pp. 2221–2230, 2014.
- [6] Y. Yang, S. Li, G. Zhu et al., “A similar local immune and oxidative stress phenotype in vitiligo and halo nevus,” *Journal of Dermatological Science*, vol. 87, no. 1, pp. 50–59, 2017.
 - [7] T. Cui, W. Zhang, S. Li et al., “Oxidative stress-induced HMGB1 release from melanocytes: a paracrine mechanism underlying the cutaneous inflammation in vitiligo,” *The Journal of Investigative Dermatology*, vol. 139, no. 10, pp. 2174–2184.e4, 2019.
 - [8] K. Mou, W. Liu, Y. Miao, F. Cao, and P. Li, “HMGB1 deficiency reduces H₂O₂-induced oxidative damage in human melanocytes via the Nrf2 pathway,” *Journal of Cellular and Molecular Medicine*, vol. 22, no. 12, pp. 6148–6156, 2018.
 - [9] H. Yanai, T. Ban, Z. Wang et al., “HMGB proteins function as universal sentinels for nucleic-acid-mediated innate immune responses,” *Nature*, vol. 462, no. 7269, pp. 99–103, 2009.
 - [10] M. Becatti, “Oxidative stress and high-mobility group box 1 (HMGB1) protein release in vitiligo,” *The British Journal of Dermatology*, vol. 176, no. 6, pp. 1436–1437, 2017.
 - [11] M. Viswanathan, K. Treiman, J. Kish-Doto, J. C. Middleton, E. J. L. Coker-Schwimmer, and W. K. Nicholson, “Folic acid supplementation for the prevention of neural tube defects,” *JAMA*, vol. 317, no. 2, pp. 190–203, 2017.
 - [12] M. Mattson, I. Kruman, and W. Duan, “Folic acid and homocysteine in age-related disease,” *Ageing Research Reviews*, vol. 1, no. 1, pp. 95–111, 2002.
 - [13] J. Chen, Q. Shi, X. Wang et al., “Genetic polymorphisms in the methylenetetrahydrofolate reductase gene (MTHFR) and risk of vitiligo in Han Chinese populations: a genotype–phenotype correlation study,” *The British Journal of Dermatology*, vol. 170, no. 5, pp. 1092–1099, 2014.
 - [14] J. Chen, T. Zhuang, J. Chen et al., “Homocysteine induces melanocytes apoptosis via PERK-eIF2 α -CHOP pathway in vitiligo,” *Clinical Science*, vol. 134, no. 10, pp. 1127–1141, 2020.
 - [15] Z. Li, D. Zhou, D. Zhang et al., “Folic acid inhibits aging-induced telomere attrition and apoptosis in astrocytes in vivo and in vitro,” *Cerebral Cortex*, 2021.
 - [16] T. Nikolic, D. Petrovic, S. Matic et al., “The influence of folic acid-induced acute kidney injury on cardiac function and redox status in rats,” *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 393, no. 1, pp. 99–109, 2020.
 - [17] G. Bodega, M. Alique, L. Puebla, J. Carracedo, and R. M. Ramírez, “Microvesicles: ROS scavengers and ROS producers,” *Journal of Extracellular Vesicles*, vol. 8, no. 1, article 1626654, 2019.
 - [18] H. Xie, F. Zhou, L. Liu et al., “Vitiligo: how do oxidative stress-induced autoantigens trigger autoimmunity?,” *Journal of Dermatological Science*, vol. 81, no. 1, pp. 3–9, 2016.
 - [19] S. Li, G. Zhu, Y. Yang et al., “Oxidative stress drives CD8⁺ T-cell skin trafficking in patients with vitiligo through CXCL16 upregulation by activating the unfolded protein response in keratinocytes,” *The Journal of Allergy and Clinical Immunology*, vol. 140, no. 1, pp. 177–189.e9, 2017.
 - [20] H. Koohpeyma, I. Goudarzi, M. Elahdadi Salmani, T. Lashkarbolouki, and M. Shabani, “Postnatal administration of homocysteine induces cerebellar damage in rats: protective effect of folic acid,” *Neurotoxicity Research*, vol. 35, no. 3, pp. 724–738, 2019.
 - [21] W. Li, Y. Ma, Z. Li et al., “Folic acid decreases astrocyte apoptosis by preventing oxidative stress-induced telomere attrition,” *International Journal of Molecular Sciences*, vol. 21, 2020.
 - [22] M. Sun, X. Xu, Y. du et al., “Biomimetic melanosomes promote orientation-selective delivery and melanocyte pigmentation in the H₂O₂-induced vitiligo mouse model,” *ACS Nano*, 2021.
 - [23] R. Singh, A. Letai, and K. Sarosiek, “Regulation of apoptosis in health and disease: the balancing act of BCL-2 family proteins,” *Nature Reviews. Molecular Cell Biology*, vol. 20, no. 3, pp. 175–193, 2019.
 - [24] D. Zhou, X. Lv, Y. Wang et al., “Folic acid alleviates age-related cognitive decline and inhibits apoptosis of neurocytes in senescence-accelerated mouse prone 8: deoxythymidine triphosphate biosynthesis as a potential mechanism,” *The Journal of Nutritional Biochemistry*, vol. 97, article 108796, 2021.
 - [25] A. Cuadrado, A. Rojo, G. Wells et al., “Therapeutic targeting of the NRF2 and KEAP1 partnership in chronic diseases,” *Nature Reviews. Drug Discovery*, vol. 18, no. 4, pp. 295–317, 2019.
 - [26] L. Qiu, Z. Song, and V. Setaluri, “Oxidative stress and vitiligo: the Nrf2-ARE signaling connection,” *The Journal of Investigative Dermatology*, vol. 134, no. 8, pp. 2074–2076, 2014.
 - [27] K. Okazaki, H. Anzawa, Z. Liu et al., “Enhancer remodeling promotes tumor-initiating activity in NRF2-activated non-small cell lung cancers,” *Nature Communications*, vol. 11, no. 1, p. 5911, 2020.
 - [28] Y. Chang, S. Li, W. Guo et al., “Simvastatin protects human melanocytes from H₂O₂-induced oxidative stress by activating Nrf2,” *The Journal of Investigative Dermatology*, vol. 137, no. 6, pp. 1286–1296, 2017.
 - [29] Z. Jian, L. Tang, X. Yi et al., “Aspirin induces Nrf2-mediated transcriptional activation of haem oxygenase-1 in protection of human melanocytes from H₂O₂-induced oxidative stress,” *Journal of Cellular and Molecular Medicine*, vol. 20, no. 7, pp. 1307–1318, 2016.
 - [30] I. Bellezza, I. Giambanco, A. Minelli, and R. Donato, “Nrf2-Keap1 signaling in oxidative and reductive stress,” *Biochimica et biophysica acta. Molecular cell research*, vol. 1865, no. 5, pp. 721–733, 2018.
 - [31] P. Cao, W. Zhang, G. Wang et al., “Low dose of folic acid can ameliorate hyperhomocysteinemia-induced cardiac fibrosis and diastolic dysfunction in spontaneously hypertensive rats,” *International Heart Journal*, vol. 62, no. 3, pp. 627–635, 2021.
 - [32] L. Shi, L. Feng, W. Jiang et al., “Folic acid deficiency impairs the gill health status associated with the NF- κ B, MLCK and Nrf2 signaling pathways in the gills of young grass carp (*Ctenopharyngodon idella*),” *Fish & Shellfish Immunology*, vol. 47, no. 1, pp. 289–301, 2015.
 - [33] H. Yang, H. Wang, and U. Andersson, “Targeting inflammation driven by HMGB1,” *Frontiers in Immunology*, vol. 11, p. 484, 2020.
 - [34] H. Xu, N. Wandersee, Y. Guo et al., “Sickle cell disease increases high mobility group box 1: a novel mechanism of inflammation,” *Blood*, vol. 124, no. 26, pp. 3978–3981, 2014.
 - [35] S. Sun, M. He, Y. Wang, H. Yang, and Y. al-Abed, “Folic acid derived-P5779 mimetics regulate DAMP-mediated inflammation through disruption of HMGB1:TLR4:MD-2 axes,” *PLoS One*, vol. 13, no. 2, article e0193028, 2018.
 - [36] W. Dong, Y. Liu, Z. Lv et al., “Lung endothelial barrier protection by resveratrol involves inhibition of HMGB1 release and HMGB1-induced mitochondrial oxidative damage via an Nrf2-dependent mechanism,” *Free Radical Biology & Medicine*, vol. 88, Part B, pp. 404–416, 2015.

- [37] T. Tsai, C. Kuo, and Y. Huang, "Serum homocysteine, folate, and vitamin B₁₂ levels in patients with vitiligo and their potential roles as disease activity biomarkers: a systematic review and meta-analysis," *Journal of the American Academy of Dermatology*, vol. 80, no. 3, pp. 646–654.e5, 2019.
- [38] A. Shirzadian Kebria, M. Hosseini, and S. Khafri, "The effect of narrowband ultraviolet B phototherapy on serum folate level," *Caspian Journal of Internal Medicine*, vol. 12, no. 2, pp. 180–183, 2021.
- [39] S. Elshafy Khashaba, R. Elkot, and A. Ibrahim, "Efficacy of NB-UVB, microneedling with triamcinolone acetonide, and a combination of both modalities in the treatment of vitiligo: a comparative study," *Journal of the American Academy of Dermatology*, vol. 79, no. 2, pp. 365–367, 2018.
- [40] M. Zhang, G. Goyert, and H. Lim, "Folate and phototherapy: what should we inform our patients?," *Journal of the American Academy of Dermatology*, vol. 77, no. 5, pp. 958–964, 2017.
- [41] A. Shakhbazova, H. Wu, C. J. Chambers, and R. K. Sivamani, "A systematic review of nutrition, supplement, and herbal-based adjunctive therapies for vitiligo," *Journal of alternative and complementary medicine*, vol. 27, no. 4, pp. 294–311, 2021.