

Research Article

Parameters of Oxidative Stress, Vitamin D, Osteopontin, and Melatonin in Patients with Lip, Oral Cavity, and Pharyngeal Cancer

Jarosław Nuskiewicz ¹, Jolanta Czuczejko ^{2,3}, Marta Maruszak ³, Marta Pawłowska ¹,
Alina Woźniak ¹, Bogdan Małkowski ⁴, and Karolina Szewczyk-Golec ¹

¹Department of Medical Biology and Biochemistry, Faculty of Medicine, Ludwik Rydygier Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University in Toruń, 24 Karłowicza St., 85-092 Bydgoszcz, Poland

²Department of Psychiatry, Faculty of Medicine, Ludwik Rydygier Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University in Toruń, 9 M. Curie Skłodowskiej St., 85-094 Bydgoszcz, Poland

³Department of Nuclear Medicine, Oncology Centre Prof. Franciszek Łukaszczyk Memorial Hospital, Bydgoszcz, 2 Dr I. Romanowskiej St., 85-796 Bydgoszcz, Poland

⁴Department of Diagnostic Imaging, Faculty of Health Sciences, Ludwik Rydygier Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University in Toruń, 2 Dr I. Romanowskiej St., 85-796 Bydgoszcz, Poland

Correspondence should be addressed to Jarosław Nuskiewicz; jnuskiewicz@cm.umk.pl

Received 14 September 2021; Accepted 7 October 2021; Published 20 October 2021

Academic Editor: Alessandra Durazzo

Copyright © 2021 Jarosław Nuskiewicz et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Lip, oral cavity, and pharyngeal cancers (LOCP) constitute a group of rare neoplasms with unfavorable prognosis. So far, not much is known about the role of vitamin D and oxidative stress in the pathogenesis of LOCP in the European population. The aim of the study was to determine the concentrations of vitamin D, osteopontin, melatonin, and malondialdehyde (MDA) as markers of oxidative stress and/or inflammation, as well as the activities of antioxidant enzymes in the course of LOCP. The vitamin D, melatonin, and osteopontin concentrations in blood serum, the MDA levels in erythrocytes and blood plasma, and the activities of superoxide dismutase (SOD-1), catalase (CAT), and glutathione peroxidase (GPx) in erythrocytes were measured in blood samples taken from 25 LOCP patients of middle age (YCG), 20 LOCP elderly patients (OCG), and 25 healthy middle-aged volunteers. In both cancer groups, decreases in vitamin D and CAT, as well as increases in osteopontin and blood plasma MDA, were observed. An increase in GPx activity in YCG and a decrease in melatonin level in OCG were found. The results indicate the vitamin D deficiency and disturbed oxidant-antioxidant homeostasis in LOCP patients. Osteopontin seems to be associated with LOCP carcinogenesis and requires further research.

1. Introduction

Lip, oral cavity, and pharyngeal cancers (LOCP) belong to the most common head and neck cancers worldwide. Moreover, scientific analyses indicate that the incidence of this type of neoplasm will increase in the future. According to data, in 2012, 529,500 new cases of LOCP were detected worldwide, which corresponds to 3.6% of all cancers [1, 2]. Mortality in 2012 from this group of neoplasms was estimated at 292,300 cases, which corresponds to 3.6% of deaths

due to neoplastic diseases [1, 2]. Projections for 2035 show a 62% increase in the number of cases to around 856,000 cases annually [1]. Cancers of lip, oral cavity, and pharynx are considered together because they are characterized by similar risk factors. Neoplasms belonging to this group affect male much more often than female, and the age group 50-70 years is particularly vulnerable [3–5]. In addition, this type of cancer is especially common in south-central Asia [1]. The main risk factors for the development of LOCP cancer include smoking [6], alcohol consumption [7, 8],

infections caused by Epstein-Barr virus (EBV) [9], and *human papillomavirus* (HPV) [10]. Early diagnosis and treatment initiation significantly increases patient survival; unfortunately, most cases are detected in the advanced stage of the disease, which lowers the 5-year survival rate to about 40% [3].

Pathogenesis of lip, oral cavity, and pharyngeal cancers is still not clear and is believed to be multifactorial in origin. Few studies indicate the participation of extracellular matrix and fibroblast changes, immune system, and oxidative stress in the pathogenesis of oral submucous fibrosis, leading to cancer of the oral cavity [11]. Additionally, molecular pathogenesis of head and neck cancer is associated with deletion in region located at chromosome 9p21–22 containing p16 tumor suppressor gene [12]. An inherent element of carcinogenesis and neoplastic disease is the increased generation of reactive oxygen species (ROS) [13–18]. Moreover, cancer cells synthesize and secrete cytokines that modulate inflammation and significantly increases ROS generation [19]. The disturbance of homeostasis by ROS generated in exceeding of physiological capacity of adaptation leads to oxidative stress [20–22]. Although the disease has a specific localization, systemic symptoms of oxidative stress are observed in patients [23, 24]. ROS are a group of chemical molecules which are characterized by the presence of nonpair electrons and high chemical reactivity [25]. The most important ROS include superoxide anion (O_2^-), hydrogen peroxide (H_2O_2), hydroxyl radical (OH^-), and singlet oxygen (1O_2) [20, 25]. Due to high chemical reactivity, ROS modify proteins, lipids, and genetic material [26, 27]. The effect of lipid peroxidation is damage to cell membranes, and the main markers of this process is malondialdehyde (MDA) and 4-hydroxynonenal [28]. Antioxidants play an important role in maintaining the redox balance [29]. Endogenous antioxidants include enzymes such as superoxide dismutases (SODs), catalase (CAT), and glutathione peroxidases (GPxs) [30]. The antioxidant defense is also constituted by small endo- and exogenous biomolecules such as vitamins A, C, and E, melatonin, and glutathione (GSH) [31, 32]. The role of vitamin D as an antioxidant remains ambiguous due to inconclusive research results [33].

Vitamin D is a biomolecule with pleiotropic properties. Calcitriol (1,25-dihydroxycholecalciferol) plays the most important role among the group of compounds called vitamin D [34]. Chemical compounds belonging to this group can be absorbed with food, most often in the form of cholecalciferol and ergocalciferol [35, 36]. Another source of vitamin D is the endogenous synthesis under the influence of ultraviolet radiation (UV) and hydroxylases found in the liver and kidney. The substrate for this process is 7-dehydrocholesterol [37]. Despite endogenous synthesis and the presence of vitamin D in food products, vitamin D deficiencies affect a significant part of the population worldwide [38–40]. Vitamin D is involved in the regulation of calcium-phosphate homeostasis, which is of particular importance for the functioning of the skeletal system [41]. Calcitriol, acting through the vitamin D receptor (VDR), reduces oxidative stress by increasing the level of SODs, GPxs, and GSH expression [33, 42]. Moreover, it was observed that vitamin

D reduced the secretion of proinflammatory cytokines, decreasing the level of oxidative stress [43]. The role of vitamin D and its derivatives in cancer is still under investigation. The results of the studies conducted so far are not unequivocal. Some researchers point to a significant role of vitamin D deficiency on cancer mortality, while no effect on morbidity [44–46]. On the contrary, some studies do not link cancer with vitamin D levels [47].

Research indicates a positive correlation between the concentration of vitamin D and osteopontin [48, 49]. Osteopontin is a glycoprotein secreted by osteoblasts and osteoclasts involved in shaping the correct bone structure [50]. The presence of this glycoprotein is not limited to the skeletal system. Osteopontin was found in many tissues and body fluids such as brain astrocytes, kidney, smooth muscle, saliva, and milk [50–52]. Tumor cells of lung, gastric, prostate, ovarian, and colorectal cancer were also found to secrete osteopontin [50]. In the course of neoplastic diseases, an increase in the concentration of osteopontin was observed along with an increase in the level of proinflammatory cytokines [53–55]. Osteopontin was found to be a modulator of the immune response [54]. The relationship between osteopontin and oxidative stress has not been analyzed frequently. The results of the research indicate that the concentration of osteopontin positively correlates with the markers of increased oxidative stress [56–59].

Melatonin (N-acetyl-5-methoxytryptamine) is a hormone synthesized and secreted by pinealocytes in the circadian rhythm [60, 61]. Gastrointestinal tract, lymphocytes, ovaries, skin, and retina are sources of extrapineal melatonin independent of the circadian rhythms [62]. The melatonin molecule contains an indole ring that neutralizes ROS directly [63, 64]. Moreover, research indicates that melatonin decreases the level of ROS by activating the silent information regulator 1 (SIRT1) pathway [65]. Melatonin may also indirectly affect the oxidant-antioxidant balance, stimulating the expression of genes encoding antioxidant enzymes, such as SODs and GPxs [66]. In addition to its direct and indirect action, melatonin inactivates ROS through its metabolites, namely, N1-acetyl-N2-formyl-5-methoxykynuramine (AFMK) and N1-acetyl-5-methoxykynuramine (AMK) [67]. The pleiotropic role of melatonin as endo- and paracrine hormone has been analyzed in carcinogenesis [68, 69]. Scientists indicated an oncostatic role of melatonin in breast, ovarian, prostate, oral, gastric, and colorectal tumors [70]. One of the mechanisms of the oncostatic action of melatonin seems to be based on the reduction of ROS levels [71, 72]. Moreover, melatonin was found to hinder angiogenesis and increase apoptosis of cancer cells [73]. However, little is known about the role of melatonin in LOCP cancer.

So far, only a few studies on the activity of antioxidant enzymes and lipid peroxidation markers in patients with LOCP have been conducted. The results described in the literature are not unequivocal. A decrease in SOD, CAT, and GPx activities with an increase in MDA concentration has been most frequently reported [74–76]. Still, according to some other research, no changes in the activity of antioxidant enzymes in the course of LOCP have been observed

[77]. The relationship between the activity of antioxidant enzymes and vitamin D and osteopontin and melatonin has not been studied. Examining the mechanisms related to vitamin D, osteopontin, melatonin, and oxidative stress in the course of LOCP seems to be important for a better understanding of the pathophysiology of this type of cancer, as well as for finding new methods of treatment and prevention. Thus, the aim of this study was to determine the activities of selected antioxidant enzymes, as well as the concentrations of vitamin D, osteopontin, melatonin, and MDA in the course of lip, oral cavity, and pharyngeal cancer.

2. Materials and Methods

2.1. Participants. The study involved 45 patients diagnosed with *carcinoma in situ* of lip, oral cavity, or pharynx according to the International Classification of Diseases–11th Revision (ICD-11)–2E60.0 [78]. The patients were divided into two groups according to their age, namely, younger cancer group (YCG) and older cancer group (OCG). The classification of patients into age groups was based on the United Nations report, which stated that old age begins after the age of 65 [79]. The participants were treated at the Oncology Center, Prof. Franciszek Łukaszyk Memorial Hospital, Bydgoszcz, Poland. The patients were referred for planning radiotherapy using positron emission tomography-computed tomography (PET/CT) after FDG ([¹⁸F]-fluorodeoxyglucose) administration. The patients with G1 squamous cell carcinoma, G2 squamous cell carcinoma, and nonkeratinizing G2 squamous cell carcinoma in histopathological analysis were included in the study. The patients with other grade and type of tumor were excluded from the study. The control group consisted of 25 healthy volunteers. The criteria of exclusion from the control group were associated conditions known to be caused by or to result in oxidative stress or involving disruption of the oxidant-antioxidant equilibrium (cancer, diabetes, cardiovascular, and infectious diseases). A survey was conducted among the people qualified for the study. The questions concerned tobacco addiction and vitamin D supplementation. The characteristics of the study and control groups are presented in Table 1. The study was approved by the Bioethics Committee of the Nicolaus Copernicus University in Toruń functioning at Collegium Medicum in Bydgoszcz, Poland (consent no. KB 221/2018).

2.2. Study Design. The patients were eligible for the study on the day of planning for radiotherapy. Blood samples were collected by qualified medical personnel in the morning (between 8:00 AM and 9:00 AM) after overnight fasting from median cubital vein just prior to the administration of the radiopharmaceutical. Every blood sample was collected into two polypropylene tubes. First tube (vol. 6 mL) contained a clotting activator to obtain blood serum, and another tube (vol. 10 mL) was covered with K₂EDTA to obtain blood plasma. The tubes were immediately transported under reduced temperature condition to the laboratory for centrifugation (6,000 g for 10 min at 4°C). After centrifugation, blood serum and plasma were separated

and stored at -80°C for further analysis. The blood morphotic elements remaining after centrifugation were washed three times with a phosphate-buffered saline (PBS) at a ratio of 1:3 and each time centrifuged (6,000 g for 10 min at 4°C) to remove leukocytes and thrombocytes. The red blood cells obtained in this method were mixed with the PBS solution to obtain erythrocytic suspension with a 50% hematocrite index.

2.3. Biochemical Analysis. The activity of selected antioxidant enzymes was determined in erythrocytic suspension with the use of spectrophotometric methods. Activity of Zn/Cu-superoxide dismutase (SOD-1; EC 1.15.1.1) was assayed according to the Misra and Fridovich method [80]. Analysis was based on the inhibition of adrenaline oxidation to adrenochrome in alkaline solution at 37°C, which induced a change in the absorbance at 480 nm. Activity of SOD-1 was expressed in IU/g Hb. CAT (EC 1.11.1.6) activity was determined with the use of the Beers and Sizer method [81] by measuring the decrease in the absorbance at 240 nm of a solution of hydrogen peroxide decomposed by the enzyme at 37°C. CAT activity was expressed in IU/g Hb. Activity of cytosolic glutathione peroxidase (GPx; EC 1.11.1.9) was assessed using the method of Paglia and Valentine [82]. The principle of the method for measuring GPx activity is based on the ability of the enzyme to reduce hydrogen peroxide with a simultaneous oxidation of GSH as a coenzyme at 37°C, measured at 340 nm. Activity of GPx was expressed in IU/g Hb. Erythrocytic and plasma MDA concentrations were determined with the method of Buege and Aust [83] in the modification of Esterbauer and Cheeseman [84]. The MDA concentration was expressed as the concentration of thiobarbituric acid-reactive substances (TBARS), measured at 532 nm at room temperature. The MDA concentration in erythrocytes was expressed in nmol/g Hb and in blood plasma in nmol/mL. Hemoglobin (Hb) concentration was evaluated using the Drabkin method [85]. Hemoglobin and selected hemoglobin derivatives under the influence of potassium ferricyanide are oxidized to methemoglobin. The absorbance is measured at 540 nm at room temperature.

Serum concentrations of melatonin, vitamin D, and osteopontin were determined with commercially available enzyme immune assay kits. The kits were used accordingly: an enzyme-linked immunosorbent assay kit for melatonin (Cloud-Clone Corp., Houston, TX, USA), a competitive enzyme-linked immunosorbent assay kit for 25(OH)-vitamin D (Immundiagnostik AG, Bensheim, Germany), and a sandwich enzyme-linked immunosorbent assay kit for human osteopontin (BioVendor, Brno, Czech Republic). The measurements were made according to manufacturer's instructions. The enzyme immune assay kits used in the study contain the reagents necessary for the study, standard concentration analytes, blank, and control samples. The principle of the assay is to bind the antigen by specific anti-human monoclonal antibodies that coat the wells of microplates found in the kits. The antigen concentration was determined from the calibration curve. The concentrations of melatonin, vitamin D, and osteopontin were expressed in pg/mL, ng/mL, and nmol/L, respectively.

TABLE 1: Anthropometric and clinical characteristic of patients with lip, oral cavity, or pharyngeal cancer and healthy volunteers (control group). Each value is mean \pm S.E.M. YCG: younger cancer group; OCG: older cancer group, * $p < 0.05$ vs. OCG.

Parameter	YCG	OCG	Control group
<i>n</i> (male/female)	25 (15/10)	20 (14/6)	25 (11/14)
Age [yrs]	58.24 \pm 1.29*	69.7 \pm 1.49	55.36 \pm 1.17*
Body mass [kg]	72.53 \pm 3.95	71.39 \pm 3.52	71.02 \pm 2.22
Height [cm]	168.92 \pm 1.84	168.55 \pm 1.52	169.88 \pm 1.72
BMI [kg/m ²]	25.09 \pm 0.99	25.00 \pm 1.02	24.50 \pm 0.47
Current smoker (y/n)	7/18	5/15	4/21
Vitamin D supplementation (y/n)	5/20	4/16	8/17

2.4. Statistical Analysis. Statistical analysis was performed using the Statistica 13.3 (TIBCO Software Inc.). The results were presented as means \pm S.E.M. Statistical analysis included Student's *t*-test for independent samples, for the comparison of study group and control group, Shapiro-Wilk test to test hypothesis of normal distribution, Levene's test to assess homogeneity of variances. Pearson's correlation coefficient was used to quantify the relationship between the parameters measured. The level of significance was set at $p < 0.05$.

3. Results

Anthropometric and clinical characteristic of patients with lip, oral cavity, or pharyngeal cancer and healthy group were presented in Table 1. No significant differences were found between YCG and control group. There was a statistically significant difference in the age of the patients between the YCG, control group, and OCG.

The SOD-1 activity was similar in all groups and amounted to 738 \pm 21 IU/g Hb in YCG, 735 \pm 19 IU/g Hb in OCG, and 755 \pm 18 IU/g Hb in control group. The statistically lower CAT activity was observed in YCG (59.34 \pm 2.6810⁴ \times IU/g Hb) and OCG (58.11 \pm 2.4710⁴ \times IU/g Hb) groups compared to the control group (70.19 \pm 1.8710⁴ \times IU/g Hb). The activity of GPx in YCG was 8.54 \pm 0.75 IU/g Hb and was significantly higher compared to the control group (6.45 \pm 0.58 IU/g Hb). In OCG, the mean GPx activity was 7.44 \pm 0.76 IU/g Hb. The results concerning the activity of antioxidant enzymes are presented in Figure 1.

There were no statistically significant differences in the concentration of erythrocytic MDA in YCG (27.52 \pm 4.23 nmol/g Hb), OCG (31.74 \pm 6.21 nmol/g Hb), and control group (25.66 \pm 1.98 nmol/g Hb). Significantly higher concentrations of MDA were observed in the plasma of YCG and OCG patients, amounting to 0.55 \pm 0.02 and 0.56 \pm 0.01 nmol/mL, respectively. In the control group, the plasma MDA level was 0.42 \pm 0.02 nmol/mL. The melatonin level in OCG was 62.12 \pm 4.70 pg/mL and was significantly lower than in the control group (84.33 \pm 6.54 pg/mL). The concentration of melatonin in YCG was 72.83 \pm 5.80 pg/mL, and no statistically significant differences were observed in relation to the other study groups. Figure 2 shows the concentration of MDA and melatonin in the graphs.

The concentration of 25(OH)-vitamin D in the serum of the healthy people was 82.57 \pm 4.28 ng/mL. Considerably lower values were observed in YCG and OCG, amounting to 63.55 \pm 7.36 ng/mL and 48.42 \pm 7.09 ng/mL, respectively. The levels of osteopontin in the LOCP patient groups were significantly higher compared to the healthy group. The concentration of osteopontin in YCG and OCG was 16.04 \pm 2.69 nmol/L and 14.98 \pm 2.48 nmol/L, respectively, while in the control group, it was 9.78 \pm 0.72 nmol/L. Figure 3 shows the graphs of 25(OH)-vitamin D and osteopontin concentration in the study groups.

The obtained data were also tested for the presence of correlations. In YCG, statistically significant negative correlations were observed between GPx and body mass ($r = -0.51$, $p = 0.009$), GPx and BMI ($r = -0.52$, $p = 0.007$), and CAT and vitamin D ($r = -0.42$, $p = 0.036$), whereas a significant positive correlation was found between plasma MDA and osteopontin ($r = 0.53$, $p = 0.007$) (see Figure 4). In the control group, a positive correlation between SOD-1 and erythrocytic MDA ($r = 0.44$, $p = 0.026$) and a negative correlation between body mass and erythrocytic MDA ($r = -0.39$, $p = 0.049$) were observed (Figure 5). No statistically significant correlations were found in OCG.

4. Discussion

As mentioned earlier, LOCP are relatively rare compared to other neoplastic diseases. Due to the small number of patients with such a diagnosis, there are few studies in which the mechanisms of antioxidant defense and the concentration of melatonin, osteopontin, and vitamin D were analyzed [74–77, 86–99]. Figure 6 presents putative mechanisms linking oxidative stress, antioxidant enzymes, vitamin D, osteopontin, and melatonin in LOCP carcinogenesis.

In the present study, specific modifications in the oxidant-antioxidant homeostasis, including a decrease in CAT and increases in GPx (in YCG) and blood plasma MDA, were found in the LOCP patients when compared to the healthy people. Moreover, GPx activity was found to negatively correlate with body mass and BMI in the YCG. Surprisingly, no statistically significant differences in SOD-1 activity were observed. This result is in contrast with the findings of other studies. In the study conducted by Gurudath et al. [86], a decrease in SOD-1 activity in patients with cancer of oral cavity was indicated. The study group




FIGURE 1: Activity of antioxidant enzymes in the erythrocytes of patients with lip, oral cavity, or pharyngeal cancer depending on age and in the healthy group. (a) Zn/Cu-superoxide dismutase (SOD-1) activity, (b) catalase (CAT) activity, (c) cytosolic glutathione peroxidase (GPx) activity. YCG: younger cancer group—mean age 58.24 ± 1.29 yrs; OCG: older cancer group—mean age 69.70 ± 1.49 yrs. Data are presented as the means \pm S.E.M. * $p < 0.05$ vs. control group.

consisted of 25 patients with oral cancer. Only smokers and tobacco chewers were included. Average age of the examined subjects was 53 yrs. The control group consisted of 25 healthy people; no information was provided on smoking or chewing tobacco in this group. Also, in the study of Subapriya et al. [87], the activity of SOD in group of 12 patients with oral precancerous lesions or oral cancer was tested. All subjects included in the study were between 35 and 60 yrs old and smoked or chewed tobacco. The cancer patients showed significantly lower SOD activity compared to the control group. Sabitha and Shyamaladevi [76] analyzed 12 blood samples obtained from patients with stage III oral can-

cer. The authors did not provide the age or information about the addictions of people participating in the study. Also, in that study, a significant decrease in SOD activity was observed in the course of cancer. Similarly, in the research by Manoharan et al. [75] decreased SOD activity was observed. A group of 46 men with oral cancer aged 40 to 60 years old was examined. The control group was free from smoking and chewing tobacco; no such information was provided in the context of the study group. Decreased activity of SOD in red blood cells was also reported by Patel et al. [77]. The age range for oral cancer patients was 22-75 years with a median of 45 years. A group of 126 patients, 113




FIGURE 2: Concentration of malondialdehyde (MDA) and melatonin in patients with lip, oral cavity, or pharyngeal cancer depending on age and in the healthy group. (a) Erythrocytic MDA concentration, (b) plasma MDA concentration, (c) melatonin concentration. YCG: younger cancer group—mean age 58.24 ± 1.29 yrs; OCG: older cancer group—mean age 69.70 ± 1.49 yrs. Data are presented as the means \pm S.E.M. * $p < 0.05$, vs. control group.

of whom smoked or chewed tobacco, was tested. Different results were observed in the study of Huo et al. [74]. In their study, a group of 25 patients of both sexes aged 40 to 45 diagnosed with oral squamous cell carcinoma were investigated. Only smokers and tobacco chewers were included in the study. A healthy control group was free of tobacco chewing and smoking habits. The activity of SOD and CAT, as well as the level of erythrocytic MDA, was tested. SOD activity was higher in the group with neoplastic disease. In the case of observed in the present study lower CAT activity in the LOCP patients, similar results were reported by Huo et al. [74], Subapriya et al. [87], Sabitha and Shyamaladevi [76], and Manoharan et al. [75]. On the contrary, Patel

et al. [77] did not observe any statistically significant changes in catalase activity in oral cancer patients. In the present study, we noted significantly higher GPx activity in the middle-aged LOCP patients than in the control group. The different results were described by Gurudath et al. [86], Subapriya et al. [87], Sabitha and Shyamaladevi [76], and Manoharan et al. [75]. The authors of these studies observed that GPx activity decreased in the course of cancer. In our study, erythrocytic MDA showed no statistically significant variability between the studied groups, unlike MDA level in blood plasma, which was higher in the cancer patients. In the other studies, increases in the level of MDA in plasma or serum and red blood cells in patients with cancers of




FIGURE 3: Concentration of 25(OH)-vitamin D and osteopontin in the blood serum of patients with lip, oral cavity, or pharyngeal cancer depending on age and in the healthy group. (a) 25(OH)-vitamin D concentration, (b) osteopontin concentration. YCG: younger cancer group—mean age 58.24 ± 1.29 yrs; OCG: older cancer group—mean age 69.70 ± 1.49 yrs. Data are presented as the means \pm S.E.M. * $p < 0.05$ vs. control group.




FIGURE 4: Statistically significant correlations in the younger cancer group (mean age 58.24 ± 1.29 yrs) between body mass and glutathione peroxidase (GPx) activity, body mass index (BMI) and GPx, osteopontin and plasma malondialdehyde (MDA) level, and vitamin D concentration and catalase (CAT) activity. The regression line is marked with a solid line, while the confidence intervals of 0.95 are marked with a dashed line.

the oral cavity and pharynx were unanimously indicated [74–76, 88–90].

Considering the differences between the studies, it is worth noting that the analyzed studies were conducted on small groups of patients of the Asian population (mainly

India) [74–77, 86–90]. Additionally, in the study groups, a significant proportion of patients were smokers or chewing tobacco. The relationship between tobacco addiction and ROS generation, which leads to an increase in the level of oxidative stress, was confirmed in numerous studies [10,




FIGURE 5: Statistically significant correlations in the healthy, control group between erythrocyte malondialdehyde (MDA) level and body mass, erythrocyte MDA concentration, and Zn/Cu-superoxide dismutase (SOD-1) activity. The regression line is marked with a solid line, while the confidence intervals of 0.95 are marked with a dashed line.




FIGURE 6: Putative mechanisms linking oxidative stress, antioxidant enzymes, vitamin D, osteopontin, and melatonin in lip, oral cavity, and pharyngeal cancer. Abbreviations used: LSD1: lysine-specific demethylase; NFκB: nuclear factor kappa-light-chain-enhancer of activated B cells; PI3K/AKT/mTOR1: phosphoinositide 3-kinase/protein kinase B/mechanistic target of rapamycin; SIRT1: silent information regulator 1.

101], so this factor could significantly influence the obtained results. The age of the patients is also known to have an impact on the oxidant-antioxidant balance of the organism [102, 103]. The discussed studies were carried out on patients from the age of 40, whereas in the present study, the patients were older [102, 103]. It is worth mentioning that the activity of SOD-1 is dependent on the zinc level in the diet, whereas GPx is an enzyme dependent on selenium. Deficiencies resulting from an unbalanced diet may reduce the activity of SOD-1 and/or GPx. The lower activity of GPx in OCG compared to YCG may be the evidence of selenium deficiency in the diet of the elderly patients with LOCP cancer. Hydrogen peroxide (H_2O_2) is a substrate for both CAT and GPxs [104, 105]. In the present study, lower CAT and higher GPx activities were observed in the cancer patients compared to the healthy control group. The lower activity of CAT might be compensated by the increase in GPx activity. Thus, the glutathione-related antioxidant defense seems to be predominant in the patients. In summary, the results of the present study point to the increased ROS generation and reduction of antioxidant defense mechanisms, which are characteristic of neoplastic diseases [18]. Increased levels of lipid peroxidation and MDA could be a consequence of the disturbance of oxidant-antioxidant homeostasis and might be involved in the carcinogenesis.

Research by Liu et al. [91] indicates the important role of melatonin as a ROS scavenger in oral cancer. The research was conducted on human umbilical vein endothelial cells (HUVECs) and six human oral cancer cell lines, including SCC25, SCC9, Tca8113, Cal27, FaDu, and human normal oral keratinocytes (hNOKs). The addition of melatonin (1 mM) to the culture medium significantly reduced the level of ROS in the Cal27 and FaDu cells. Concurrently, melatonin reduced the proliferation and induced the apoptosis of oral cancer cells. Observed inactivation of ROS-reliant Akt signaling significantly decreased the mobility of cancer cells. Inhibition of angiogenesis and reduction in tumor mass were also found. Yang et al. [92] analyzed oral squamous cell carcinoma (OSCC) tissue arrays. The reduction of lysine-specific demethylase (LSD1) expression under the influence of melatonin (0.1 g/mL) was described. Lower LSD-1 expression significantly reduced tumor cell proliferation. Human nasopharyngeal carcinoma (HONE-1), NPC-39, and NPC-BM cell line incubated in a solution containing melatonin (50 ng/mL) were investigated by Ho et al. [93]. Presence of melatonin inhibited TPA-induced cell motility by regulating the matrix metalloproteinase-9 (MMP-9) expression in nasopharyngeal neoplasm cells. Many scientists point to the protective role of melatonin in oral and nasopharyngeal cavity diseases mainly by reducing oxidative stress [106–109]. However, in the present study, a statistically significant lower melatonin concentration in patients with lip, oral cavity, or pharynx cancer compared to the healthy group was only observed in OCG. During the aging, the synthesis and secretion of melatonin are reduced [110]. It could indicate that melatonin deficiency is not particularly involved in the pathogenesis of LOCP [106–109].

Vitamin D deficiency was found to correlate with mortality in the course of neoplastic diseases [44–46]. Calcitriol

modulates immune response of the tumor microenvironment through the inactivation of the NF κ B (nuclear factor kappa-light-chain-enhancer of activated B cells) pathway [46]. The reference value for vitamin D in blood plasma is 75–125 nmol/L [111]. In our study, we observed that in the group of cancer patients, the vitamin D concentration was below the normal level. It is in accordance with the results of another research. Vitamin D deficiency was found to correlate with mortality in the course of neoplastic diseases [44–46]. The role of vitamin D in oral squamous cell carcinoma was investigated by Verma et al. [94]. Female C57BL/6 mice exposed to 4-nitroquinoline-1-oxide (4NQO) carcinogen were used. The animals were supplemented with vitamin D in the dose of 25–10,000 IU. The inhibition of tumor growth was observed. Beneficial effect in lowering oral mucositis in patients with head and neck cancer was also described by Bakr et al. [95]. The study involved 45 patients treated with radiotherapy divided into three groups. Two groups received topical oral vitamin D gel. Before the intervention, the levels of vitamin D in the blood serum of the patients were found to be deficient. The applied treatment not only reduced oral mucositis but also increased the level of vitamin D in the blood serum of patients. Also, Anand et al. [96] described vitamin D deficiency in oral cavity cancer patients. Moreover, the VDR overexpression in the course of oral cavity neoplasms was indicated in that study. Additionally, vitamin D appeared to play a special role in maintaining oral cavity health [112]. The surprising result of the present study is the negative correlation between CAT activity and vitamin D in the cancer group, suggesting that some regulatory mechanisms might be involved in the course of the disease. Undoubtedly, the role of vitamin D and its deficiency in LOCP cancer requires further research [112].

A significantly higher concentration of osteopontin was observed in our study in the LOCP patients. Also, Jeyasivanesan et al. [97] observed elevated osteopontin levels in patients with oral squamous cell carcinoma. Significant expression of osteopontin in salivary gland tumors was demonstrated in the study by Darling et al. [98]. Muramatsu et al. [99] analysed the effect of osteopontin levels on the invasiveness of oral cavity neoplasms. The study, performed on human oral squamous cell carcinoma cell lines, namely, HSC2, HSC3, HSC4, SAS, KB, and BSC-OF, revealed that high levels of osteopontin may increase the probability of metastasis. However, the mechanisms that link osteopontin and oral carcinomas have been not fully described yet. Presumably, osteopontin binds to integrin A4 β 1 and CD44 receptors, activating phosphoinositide 3-kinase/protein kinase B/mechanistic target of rapamycin (PI3K/AKT/mTOR1) pathway. In next step, mTOR1 regulates estrogen-related receptor alpha (ERR α), which binds to DNA and active transcription of osteopontin [113]. Overexpression of osteopontin was associated with increased angiogenesis, cancer cell proliferation, mobility, survival, invasion, and metastasis [113]. The positive correlation between osteopontin and lipid peroxidation, found in the present study, points to the association of the protein with deleterious oxidative processes in plasma membranes. Undoubtedly, these relations should be under further investigation.

The present study has some limitations. The small number of participants is a limiting factor. However, to the best of the authors' knowledge, no study has been conducted with the participation of patients with lip, oral cavity, and pharyngeal cancer of the European population, in which the mechanisms of antioxidant defense and the role of melatonin, vitamin D, and osteopontin were simultaneously analyzed.

5. Conclusions

The obtained results indicate a disruption of oxidant-antioxidant homeostasis in the lip, oral cavity, and pharyngeal cancer patients. Impaired antioxidant enzymatic defense and increased lipid peroxidation, correlated with high levels of osteopontin, were determined in this type of cancer. According to the results of the conducted study, melatonin seems not to be involved in the pathogenesis of the analyzed group of neoplasms. However, vitamin D deficiency in the LOCP patients was found. The role of elevated osteopontin in the pathogenesis of lip, oral cavity, and pharyngeal carcinoma requires further research. The simultaneous testing of vitamin D and osteopontin levels seems to be particularly noteworthy.

Data Availability

Data are available on request due to privacy/ethical restrictions.

Conflicts of Interest

The authors declare no conflict of interest.

Authors' Contributions

J.N., K.S.-G., and J.C. are responsible for the conceptualization; J.N., J.C., M.M., and K.S.-G. for the methodology; J.N. and K.S.-G. for the formal analysis; J.N., K.S.-G., J.C., M.M., and M.P. for the investigation; J.N. for the data curation; J.N. for the writing—original draft preparation; J.N. and K.S.-G. for the writing—review and editing; J.N. for the visualization; A.W. and B.M. for the supervision; J.N. and K.S.-G. for the project administration; and A.W. for the funding acquisition. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

This research was funded by the Faculty of Medicine, Ludwik Rydygier Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University, grant number MN-SDL-7/WL/2019, and Nicolaus Copernicus University grant number IDUB limit 1702-GRANTS_4NCU_STUDENTS_IDUB 2020-1-NZ-NU SZKIEWICZ.

References

- [1] K. D. Shield, J. Ferlay, A. Jemal et al., "The global incidence of lip, oral cavity, and pharyngeal cancers by subsite in 2012," *CA: a Cancer Journal for Clinicians*, vol. 67, no. 1, pp. 51–64, 2017.
- [2] *World Health Organization International Agency for Research on Cancer Global Cancer Observatory* <https://gco.iarc.fr/>.
- [3] D. Hashim, E. Genden, M. Posner, M. Hashibe, and P. Boffetta, "Head and neck cancer prevention: from primary prevention to impact of clinicians on reducing burden," *Annals of Oncology*, vol. 30, no. 5, pp. 744–756, 2019.
- [4] K. Saika and T. Matsuda, "International comparison of lip, oral cavity and pharynx cancer incidence," *Japanese Journal of Clinical Oncology*, vol. 50, no. 4, pp. 479–480, 2020.
- [5] B. Gupta, N. W. Johnson, and N. Kumar, "Global epidemiology of head and neck cancers: a continuing challenge," *Oncology*, vol. 91, no. 1, pp. 13–23, 2016.
- [6] T. D. Ellington, S. J. Henley, V. Senkomago et al., "Trends in incidence of cancers of the oral cavity and pharynx — United States 2007–2016," *MMWR. Morbidity and Mortality Weekly Report*, vol. 69, no. 15, pp. 433–438, 2020.
- [7] N. Schwartz, D. Nishri, S. Chin Cheong, N. Giesbrecht, and J. Klein-Geltink, "Is there an association between trends in alcohol consumption and cancer mortality? Findings from a multicountry analysis," *European Journal of Cancer Prevention*, vol. 28, no. 1, pp. 45–53, 2019.
- [8] L. C. S. Thuler, R. F. de Menezes, and A. Bergmann, "Cancer cases attributable to alcohol consumption in Brazil," *Alcohol*, vol. 54, pp. 23–26, 2016.
- [9] Y.-P. Chen, A. T. C. Chan, Q.-T. Le, P. Blanchard, Y. Sun, and J. Ma, "Nasopharyngeal carcinoma," *The Lancet*, vol. 394, no. 10192, pp. 64–80, 2019.
- [10] B. C. Leonard, E. D. Lee, N. E. Bhola et al., "ATR inhibition sensitizes HPV– and HPV+ head and neck squamous cell carcinoma to cisplatin," *Oral Oncology*, vol. 95, pp. 35–42, 2019.
- [11] V. Rai, S. Bose, S. Saha, and C. Chakraborty, "Evaluation of oxidative stress and the microenvironment in oral submucous fibrosis," *Heliyon*, vol. 5, no. 4, article e01502, 2019.
- [12] H. Ram, J. Sarkar, H. Kumar, R. Konwar, M. L. B. Bhatt, and S. Mohammad, "Oral cancer: risk factors and molecular pathogenesis," *Journal of Maxillofacial and Oral Surgery*, vol. 10, no. 2, pp. 132–137, 2011.
- [13] S. Reuter, S. C. Gupta, M. M. Chaturvedi, and B. B. Aggarwal, "Oxidative stress, inflammation, and cancer: how are they linked?," *Free Radical Biology & Medicine*, vol. 49, no. 11, pp. 1603–1616, 2010.
- [14] J. G. Gill, E. Piskounova, and S. J. Morrison, "Cancer, oxidative stress, and metastasis," *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 81, pp. 163–175, 2016.
- [15] I. I. C. Chio and D. A. Tuveson, "ROS in cancer: the burning question," *Trends in Molecular Medicine*, vol. 23, no. 5, pp. 411–429, 2017.
- [16] S. Chikara, L. D. Nagaprashantha, J. Singhal, D. Horne, S. Awasthi, and S. S. Singhal, "Oxidative stress and dietary phytochemicals: role in cancer chemoprevention and treatment," *Cancer Letters*, vol. 413, pp. 122–134, 2018.
- [17] Y.-H. Lin, "MicroRNA networks modulate oxidative stress in cancer," *International Journal of Molecular Sciences*, vol. 20, no. 18, p. 4497, 2019.
- [18] S. Sajadimajd and M. Khazaei, "Oxidative stress and cancer: the role of Nrf2," *Current Cancer Drug Targets*, vol. 18, no. 6, pp. 538–557, 2018.

- [19] M. Murata, "Inflammation and cancer," *Environmental Health and Preventive Medicine*, vol. 23, no. 1, p. 50, 2018.
- [20] A. V. Kudryavtseva, G. S. Krasnov, A. A. Dmitriev et al., "Mitochondrial dysfunction and oxidative stress in aging and cancer," *Oncotarget*, vol. 7, no. 29, pp. 44879–44905, 2016.
- [21] B. L. Tan, M. E. Norhaizan, and W.-P.-P. Liew, "Nutrients and oxidative stress: friend or foe?," *Oxidative Medicine and Cellular Longevity*, vol. 2018, Article ID 9719584, 2018.
- [22] S. K. Saha, S. B. Lee, J. Won et al., "Correlation between oxidative stress, nutrition, and cancer initiation," *International Journal of Molecular Sciences*, vol. 18, no. 7, p. 1544, 2017.
- [23] P. Poprac, K. Jomova, M. Simunkova, V. Kollar, C. J. Rhodes, and M. Valko, "Targeting free radicals in oxidative stress-related human diseases," *Trends in Pharmacological Sciences*, vol. 38, no. 7, pp. 592–607, 2017.
- [24] J. E. Klaunig, "Oxidative stress and cancer," *Current Pharmaceutical Design*, vol. 24, no. 40, pp. 4771–4778, 2019.
- [25] V. Sosa, T. Moliné, R. Somoza, R. Paciucci, H. Kondoh, and M. E. LLeonart, "Oxidative stress and cancer: an overview," *Ageing Research Reviews*, vol. 12, no. 1, pp. 376–390, 2013.
- [26] L.-D. Kajarabille, "Programmed cell-death by ferroptosis: antioxidants as mitigators," *International Journal of Molecular Sciences*, vol. 20, no. 19, p. 4968, 2019.
- [27] R. K. Gupta, A. K. Patel, N. Shah et al., "Oxidative stress and antioxidants in disease and cancer: a review," *Asian Pacific Journal of Cancer Prevention*, vol. 15, no. 11, pp. 4405–4409, 2014.
- [28] A. Ayala, M. F. Muñoz, and S. Argüelles, "Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal," *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 360438, 2014.
- [29] H. Sies, "Oxidative stress: a concept in redox biology and medicine," *Redox Biology*, vol. 4, pp. 180–183, 2015.
- [30] X. G. Lei, J.-H. Zhu, W.-H. Cheng et al., "Paradoxical roles of antioxidant enzymes: basic mechanisms and health implications," *Physiological Reviews*, vol. 96, no. 1, pp. 307–364, 2016.
- [31] G. Murdaca, A. Tonacci, S. Negrini et al., "Emerging role of vitamin D in autoimmune diseases: an update on evidence and therapeutic implications," *Autoimmunity Reviews*, vol. 18, no. 9, p. 102350, 2019.
- [32] W. A. C. K. Koekkoek and A. R. H. van Zanten, "Antioxidant vitamins and trace elements in critical illness," *Nutrition in Clinical Practice*, vol. 31, no. 4, pp. 457–474, 2016.
- [33] G. Hajiluan, M. Abbasalizad Farhangi, G. Nameni, P. Shahabi, and M. Megari-Abbasi, "Oxidative stress-induced cognitive impairment in obesity can be reversed by vitamin D administration in rats," *Nutritional Neuroscience*, vol. 21, no. 10, pp. 744–752, 2018.
- [34] D.-H. Kim, C. A. Meza, H. Clarke, J.-S. Kim, and R. C. Hickner, "Vitamin D and endothelial function," *Nutrients*, vol. 12, no. 2, p. 575, 2020.
- [35] L. R. Wilson, L. Tripkovic, K. H. Hart, and S. A. Lanham-New, "Vitamin D deficiency as a public health issue: using vitamin D₂ or vitamin D₃ in future fortification strategies," *The Proceedings of the Nutrition Society*, vol. 76, no. 3, pp. 392–399, 2017.
- [36] G. Cardwell, J. Bornman, A. James, and L. Black, "A review of mushrooms as a potential source of dietary vitamin D," *Nutrients*, vol. 10, no. 10, p. 1498, 2018.
- [37] A. Hanel and C. Carlberg, "Vitamin D and evolution: pharmacologic implications," *Biochemical Pharmacology*, vol. 173, p. 113595, 2020.
- [38] S.-W. Chang and H.-C. Lee, "Vitamin D and health - the missing vitamin in humans," *Pediatrics and Neonatology*, vol. 60, no. 3, pp. 237–244, 2019.
- [39] M. Sosa Henríquez and M. J. Gómez de Tejada Romero, "Cholecalciferol or calcifediol in the management of vitamin D deficiency," *Nutrients*, vol. 12, no. 6, p. 1617, 2020.
- [40] Y. Zhang, F. Fang, J. Tang et al., "Association between vitamin D supplementation and mortality: systematic review and meta-analysis," *BMJ*, vol. 366, p. 14673, 2019.
- [41] G. A. S. Casseb, M. P. Kaster, and A. L. S. Rodrigues, "Potential role of vitamin D for the management of depression and anxiety," *CNS Drugs*, vol. 33, no. 7, pp. 619–637, 2019.
- [42] S. J. Wimalawansa, "Vitamin D deficiency: effects on oxidative stress, epigenetics, gene regulation, and aging," *Biology*, vol. 8, no. 2, pp. 15–30, 2019.
- [43] M. Olszowiec-Chlebna, A. Koniarek-Maniecka, A. Brzozowska, A. Błaż, B. Rychlik, and I. Stelmach, "Vitamin D inhibits pro-inflammatory cytokines in the airways of cystic fibrosis patients infected by *Pseudomonas aeruginosa* - pilot study," *Italian Journal of Pediatrics*, vol. 45, no. 1, p. 41, 2019.
- [44] N. Keum, D. H. Lee, D. C. Greenwood, J. E. Manson, and E. Giovannucci, "Vitamin D supplementation and total cancer incidence and mortality: a meta-analysis of randomized controlled trials," *Annals of Oncology*, vol. 30, no. 5, pp. 733–743, 2019.
- [45] S. Minisola, F. Ferrone, V. Danese et al., "Controversies surrounding vitamin D: focus on supplementation and cancer," *International Journal of Environmental Research and Public Health*, vol. 16, no. 2, p. 189, 2019.
- [46] M. A. Zmijewski, "Vitamin D and human health," *International Journal of Molecular Sciences*, vol. 20, no. 1, p. 145, 2019.
- [47] B. Goulão, F. Stewart, J. A. Ford, G. MacLennan, and A. Avenell, "Cancer and vitamin D supplementation: a systematic review and meta-analysis," *The American Journal of Clinical Nutrition*, vol. 107, no. 4, pp. 652–663, 2018.
- [48] B. Enkhjargal, J. Malaguit, W. M. Ho et al., "Vitamin D attenuates cerebral artery remodeling through VDR/AMPK/eNOS dimer phosphorylation pathway after subarachnoid hemorrhage in rats," *Journal of Cerebral Blood Flow and Metabolism*, vol. 39, no. 2, pp. 272–284, 2019.
- [49] J. van de Peppel and J. P. T. M. van Leeuwen, "Vitamin D and gene networks in human osteoblasts," *Frontiers in Physiology*, vol. 5, pp. 1–10, 2014.
- [50] M. A. Icer and M. Gezmen-Karadag, "The multiple functions and mechanisms of osteopontin," *Clinical Biochemistry*, vol. 59, pp. 17–24, 2018.
- [51] L. Sayegh, G. E.-H. Fuleihan, and A. H. Nassar, "Vitamin D in endometriosis: a causative or confounding factor?," *Metabolism*, vol. 63, no. 1, pp. 32–41, 2014.
- [52] B. Enkhjargal, D. W. McBride, A. Manaenko et al., "Intra-nasal administration of vitamin D attenuates blood-brain barrier disruption through endogenous upregulation of osteopontin and activation of CD44/P-gp glycosylation signaling after subarachnoid hemorrhage in rats," *Journal of*

- Cerebral Blood Flow and Metabolism*, vol. 37, no. 7, pp. 2555–2566, 2017.
- [53] A.-S. Lamort, I. Giopanou, I. Psallidas, and G. T. Stathopoulos, “Osteopontin as a link between inflammation and cancer: the thorax in the spotlight,” *Cell*, vol. 8, no. 8, p. 815, 2019.
- [54] L. M. Castello, D. Raineri, L. Salmi et al., “Osteopontin at the crossroads of inflammation and tumor progression,” *Mediators of Inflammation*, vol. 2017, Article ID 4049098, 22 pages, 2017.
- [55] H. Miki, A. Okito, M. Akiyama, T. Ono, N. Tachikawa, and K. Nakahama, “Genetic and epigenetic regulation of osteopontin by cyclic adenosine 3' 5'-monophosphate in osteoblasts,” *Gene*, vol. 763, p. 145059, 2020.
- [56] P. Georgiadou, E. Iliodromitis, C. Varounis et al., “Relationship between plasma osteopontin and oxidative stress in patients with coronary artery disease,” *Expert Opinion on Therapeutic Targets*, vol. 12, no. 8, pp. 917–920, 2008.
- [57] M. Kocak, “The effect of antithyroid drugs on osteopontin and oxidative stress in graves' disease,” *Acta Endocrinologica*, vol. 15, no. 2, pp. 221–224, 2019.
- [58] J. Trostel, L. D. Truong, C. Roncal-Jimenez et al., “Different effects of global osteopontin and macrophage osteopontin in glomerular injury,” *American Journal of Physiology-Renal Physiology*, vol. 315, no. 4, pp. F759–F768, 2018.
- [59] T. Wolak, H. Kim, Y. Ren, J. Kim, N. D. Vaziri, and S. B. Nicholas, “Osteopontin modulates angiotensin II-induced inflammation, oxidative stress, and fibrosis of the kidney,” *Kidney International*, vol. 76, no. 1, pp. 32–43, 2009.
- [60] A. T. Slominski, R. Hardeland, M. A. Zmijewski, R. M. Slominski, R. J. Reiter, and R. Paus, “Melatonin: a cutaneous perspective on its production, metabolism, and functions,” *The Journal of Investigative Dermatology*, vol. 138, no. 3, pp. 490–499, 2018.
- [61] X. Meng, Y. Li, S. Li et al., “Dietary sources and bioactivities of melatonin,” *Nutrients*, vol. 9, no. 4, p. 367, 2017.
- [62] N. Ferlazzo, G. Andolina, A. Cannata et al., “Is melatonin the cornucopia of the 21st century?,” *Antioxidants*, vol. 9, no. 11, p. 1088, 2020.
- [63] S.-Y. Wang, X.-C. Shi, and P. Laborda, “Indole-based melatonin analogues: synthetic approaches and biological activity,” *European Journal of Medicinal Chemistry*, vol. 185, p. 111847, 2020.
- [64] R. Hardeland, “Melatonin and the electron transport chain,” *Cellular and Molecular Life Sciences*, vol. 74, no. 21, pp. 3883–3896, 2017.
- [65] G. Xu, J. Zhao, H. Liu, J. Wang, and W. Lu, “Melatonin inhibits apoptosis and oxidative stress of mouse Leydig cells via a SIRT1-dependent mechanism,” *Molecules*, vol. 24, no. 17, p. 3084, 2019.
- [66] R. He, M. Cui, H. Lin et al., “Melatonin resists oxidative stress-induced apoptosis in nucleus pulposus cells,” *Life Sciences*, vol. 199, pp. 122–130, 2018.
- [67] J. Jaworek, A. Leja-Szpak, K. Nawrot-Porąbka et al., “Effects of melatonin and its analogues on pancreatic inflammation, enzyme secretion, and tumorigenesis,” *International Journal of Molecular Sciences*, vol. 18, no. 5, p. 1014, 2017.
- [68] W. Talib, “Melatonin and cancer hallmarks,” *Molecules*, vol. 23, no. 3, p. 518, 2018.
- [69] R. Reiter, S. Rosales-Corral, D.-X. Tan et al., “Melatonin, a full service anti-cancer agent: inhibition of initiation, progression and metastasis,” *International Journal of Molecular Sciences*, vol. 18, no. 4, p. 843, 2017.
- [70] Y. Li, S. Li, Y. Zhou et al., “Melatonin for the prevention and treatment of cancer,” *Oncotarget*, vol. 8, no. 24, pp. 39896–39921, 2017.
- [71] S. Proietti, A. Cucina, M. Minini, and M. Bizzarri, “Melatonin, mitochondria, and the cancer cell,” *Cellular and Molecular Life Sciences*, vol. 74, no. 21, pp. 4015–4025, 2017.
- [72] F. Moradkhani, M. Moloudizargari, M. Fallah, N. Asghari, H. Heidari Khoei, and M. H. Asghari, “Immunoregulatory role of melatonin in cancer,” *Journal of Cellular Physiology*, vol. 235, no. 2, pp. 745–757, 2020.
- [73] S. Bhattacharya, K. K. Patel, D. Dehari, A. K. Agrawal, and S. Singh, “Melatonin and its ubiquitous anticancer effects,” *Molecular and Cellular Biochemistry*, vol. 462, no. 1–2, pp. 133–155, 2019.
- [74] W. Huo, Z.-M. Li, X.-Y. Pan, Y.-M. Bao, and L.-J. An, “Antioxidant enzyme levels in pathogenesis of oral squamous cell carcinoma (OSCC),” *Drug Research*, vol. 64, no. 11, pp. 629–632, 2014.
- [75] S. Manoharan, K. Kolanjiappan, K. Suresh, and K. Panjamurthy, “Lipid peroxidation & antioxidants status in patients with oral squamous cell carcinoma,” *The Indian Journal of Medical Research*, vol. 122, pp. 529–534, 2005.
- [76] K. E. Sabitha and C. S. Shyamaladevi, “Oxidant and antioxidant activity changes in patients with oral cancer and treated with radiotherapy,” *Oral Oncology*, vol. 35, no. 3, pp. 273–277, 1999.
- [77] P. Patel, J. Patel, F. Shah, S. Shukla, and P. Shah, “Role of nitric oxide and antioxidant enzymes in the pathogenesis of oral cancer,” *Journal of Cancer Research and Therapeutics*, vol. 5, no. 4, p. 247, 2009.
- [78] *World Health Organization International classification of diseases for mortality and morbidity statistics (11th Revision)*<https://icd.who.int/browse11/l-m/en>.
- [79] United Nations, *Population Division World Population Ageing*, Department of Economic and Social Affairs, 2015.
- [80] H. P. Misra and I. Fridovich, “The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase,” *The Journal of Biological Chemistry*, vol. 247, no. 10, pp. 3170–3175, 1972.
- [81] R. F. Beers and I. W. Sizer, “A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase,” *The Journal of Biological Chemistry*, vol. 195, no. 1, pp. 133–140, 1952.
- [82] D. E. Paglia and W. N. Valentine, “Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase,” *The Journal of Laboratory and Clinical Medicine*, vol. 70, pp. 158–169, 1967.
- [83] J. A. Buege and S. D. Aust, “Microsomal lipid peroxidation,” in *In Methods in Enzymology Vol. 52*, pp. 302–310, Academic press, United States, 1978.
- [84] H. Esterbauer and K. H. Cheeseman, “[42] Determination of aldehydic lipid peroxidation products: Malonaldehyde and 4-hydroxynonenal,” in *Oxygen Radicals in Biological Systems Part B: Oxygen Radicals and Antioxidants vol. 186*, pp. 407–421, Elsevier, 1990.
- [85] D. L. Drabkin, “Spectrophotometric studies: XIV. The crystallographic and optical properties of the hemoglobin of man in comparison with those of other species,” *The Journal of Biological Chemistry*, vol. 164, no. 2, pp. 703–723, 1946.

- [86] S. Gurudath, K. S. Ganapathy, A. Pai, S. Ballal, and M. L. Asha, "Estimation of superoxide dismutase and glutathione peroxidase in oral submucous fibrosis, oral leukoplakia and oral cancer - a comparative study," *Asian Pacific Journal of Cancer Prevention*, vol. 13, no. 9, pp. 4409–4412, 2012.
- [87] R. Subapriya, R. Kumaraguruparan, S. Nagini, and A. Thangavelu, "Oxidant-antioxidant status in oral precancer and oral cancer patients," *Toxicology Mechanisms and Methods*, vol. 13, no. 1, pp. 77–81, 2003.
- [88] V. Marakala, M. Malathi, and A. R. Shivashankara, "Lipid peroxidation and antioxidant vitamin status in oral cavity and oropharyngeal cancer patients," *Asian Pacific Journal of Cancer Prevention*, vol. 13, no. 11, pp. 5763–5765, 2012.
- [89] U. U. Malik, I. A. Siddiqui, Z. Hashim, and S. Zarina, "Measurement of serum paraoxonase activity and MDA concentrations in patients suffering with oral squamous cell carcinoma," *Clinica Chimica Acta*, vol. 430, pp. 38–42, 2014.
- [90] R. Chole, R. Patil, A. Basak, K. Palandurkar, and R. Bhowate, "Estimation of serum malondialdehyde in oral cancer AND precancer and its association with healthy individuals, gender, alcohol, and tobacco abuse," *Journal of Cancer Research and Therapeutics*, vol. 6, no. 4, p. 487, 2010.
- [91] R. Liu, H. Wang, M. Deng et al., "Melatonin Inhibits Reactive Oxygen Species-Driven Proliferation, Epithelial-Mesenchymal Transition, and Vasculogenic Mimicry in Oral Cancer," *Oxidative Medicine and Cellular Longevity*, vol. 2018, Article ID 3510970, 2018.
- [92] C.-Y. Yang, C.-K. Lin, C.-H. Tsao et al., "Melatonin exerts anti-oral cancer effect via suppressing LSD1 in patient-derived tumor xenograft models," *Oncotarget*, vol. 8, no. 20, pp. 33756–33769, 2017.
- [93] H.-Y. Ho, C.-W. Lin, M.-H. Chien et al., "Melatonin suppresses TPA-induced metastasis by downregulating matrix metalloproteinase-9 expression through JNK/SP-1 signaling in nasopharyngeal carcinoma," *Journal of Pineal Research*, vol. 61, no. 4, pp. 479–492, 2016.
- [94] A. Verma, V. K. Vincent-Chong, H. DeJong, P. A. Hershberger, and M. Seshadri, "Impact of dietary vitamin D on initiation and progression of oral cancer," *The Journal of Steroid Biochemistry and Molecular Biology*, vol. 199, p. 105603, 2020.
- [95] I. S. Bakr, A. M. Zaki, R. M. El-Moslemany, and R. O. Elsaka, "Vitamin D oral gel for prevention of radiation-induced oral mucositis: a randomized clinical trial," *Oral Diseases*, vol. 27, no. 5, pp. 1197–1204, 2021.
- [96] A. Anand, S. Singh, A. A. Sonkar et al., "Expression of vitamin D receptor and vitamin D status in patients with oral neoplasms and effect of vitamin D supplementation on quality of life in advanced cancer treatment," *Contemporary Oncology*, vol. 2, pp. 145–151, 2017.
- [97] D. Jeyasivanesan, S. Mohamed, D. Pandiar, and S. Basheer, "Immunohistochemical analysis of osteopontin expression in oral squamous cell carcinoma," *Indian Journal of Dental Research*, vol. 30, no. 4, p. 539, 2019.
- [98] M. R. Darling, M. Gauthier, L. Jackson-Boeters, T. D. Daley, A. F. Chambers, and A. B. Tuck, "Osteopontin expression in salivary gland tumours," *Oral Oncology*, vol. 42, no. 4, pp. 363–369, 2006.
- [99] T. Muramatsu, K. Shima, K. Ohta et al., "Inhibition of osteopontin expression and function in oral cancer cell lines by antisense oligonucleotides," *Cancer Letters*, vol. 217, no. 1, pp. 87–95, 2005.
- [100] M. Taylor, T. Carr, O. Oke et al., "E-cigarette aerosols induce lower oxidative stress in vitro when compared to tobacco smoke," *Toxicology Mechanisms and Methods*, vol. 26, no. 6, pp. 465–476, 2016.
- [101] R. Carnevale, S. Sciarretta, F. Violi et al., "Acute impact of tobacco vs electronic cigarette smoking on oxidative stress and vascular function," *Chest*, vol. 150, no. 3, pp. 606–612, 2016.
- [102] S. Abrahams, W. L. Haylett, G. Johnson, J. A. Carr, and S. Bardien, "Antioxidant effects of curcumin in models of neurodegeneration, aging, oxidative and nitrosative stress: a review," *Neuroscience*, vol. 406, pp. 1–21, 2019.
- [103] J. Luo, K. Mills, S. le Cessie, R. Noordam, and D. van Heemst, "Ageing, age-related diseases and oxidative stress: what to do next?," *Ageing Research Reviews*, vol. 57, p. 100982, 2020.
- [104] B. Speckmann, H. Steinbrenner, T. Grune, and L.-O. Klotz, "Peroxynitrite: from interception to signaling," *Archives of Biochemistry and Biophysics*, vol. 595, pp. 153–160, 2016.
- [105] A. Sofo, A. Scopa, M. Nuzzaci, and A. Vitti, "Ascorbate peroxidase and catalase activities and their genetic regulation in plants subjected to drought and salinity stresses," *International Journal of Molecular Sciences*, vol. 16, no. 12, pp. 13561–13578, 2015.
- [106] G. Gómez-Moreno, J. Guardia, M. Ferrera, A. Cutando, and R. Reiter, "Melatonin in diseases of the oral cavity," *Oral Diseases*, vol. 16, no. 3, pp. 242–247, 2010.
- [107] R. J. Reiter, S. A. Rosales-Corral, X. Y. Liu, D. Acuna-Castroviejo, G. Escames, and D.-X. Tan, "Melatonin in the oral cavity: physiological and pathological implications," *Journal of Periodontal Research*, vol. 50, no. 1, pp. 9–17, 2015.
- [108] A. Cutando, J. Aneiros-Fernández, J. Aneiros-Cachaza, and S. Arias-Santiago, "Melatonin and cancer: current knowledge and its application to oral cavity tumours," *Journal of Oral Pathology & Medicine*, vol. 40, no. 8, pp. 593–597, 2011.
- [109] S. Najeeb, Z. Khurshid, S. Zohaib, and M. S. Zafar, "Therapeutic potential of melatonin in oral medicine and periodontology," *The Kaohsiung Journal of Medical Sciences*, vol. 32, no. 8, pp. 391–396, 2016.
- [110] R. Hardeland, "Aging, melatonin, and the pro- and anti-inflammatory networks," *International Journal of Molecular Sciences*, vol. 20, no. 5, p. 1223, 2019.
- [111] P. Pludowski, M. F. Holick, W. B. Grant et al., "Vitamin D supplementation guidelines," *The Journal of Steroid Biochemistry and Molecular Biology*, vol. 175, pp. 125–135, 2018.
- [112] J. Botelho, V. Machado, L. Proença, A. S. Delgado, and J. J. Mendes, "Vitamin D deficiency and oral health: a comprehensive review," *Nutrients*, vol. 12, no. 5, p. 1471, 2020.
- [113] E. S. Santos, J. C. Ramos, A. L. O. C. Roza, B. A. L. A. Mariz, and A. F. Paes Leme, "The role of osteopontin in oral cancer: a brief review with emphasis on clinical applications," *Oral Diseases*, pp. 1–10, 2020.