

Research Article

Antioxidant Mechanisms of the Oligopeptides (FWKVV and FMPLH) from Muscle Hydrolysate of Miiuy Croaker against Oxidative Damage of HUVECs

Yue-Zhen Wang,¹ Yu-Mei Wang,¹ Xin Pan,² Chang-Feng Chi^{ID},² and Bin Wang^{ID}¹

¹Zhejiang Provincial Engineering Technology Research Center of Marine Biomedical Products, School of Food and Pharmacy, Zhejiang Ocean University, Zhoushan 316022, China

²National and Provincial Joint Laboratory of Exploration and Utilization of Marine Aquatic Genetic Resources, National Engineering Research Center of Marine Facilities Aquaculture, School of Marine Science and Technology, Zhejiang Ocean University, Zhoushan 316022, China

Correspondence should be addressed to Chang-Feng Chi; chichangfeng@hotmail.com and Bin Wang; wangbin4159@hotmail.com

Received 28 March 2021; Revised 13 July 2021; Accepted 3 August 2021; Published 23 August 2021

Academic Editor: Marina Sokovi

Copyright © 2021 Yue-Zhen Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this work, the antioxidant mechanisms of bioactive oligopeptides (FWKVV and FMPLH) from protein hydrolysate of miiuy croaker muscle against H₂O₂-damaged human umbilical vein endothelial cells (HUVECs) were researched systematically. The finding demonstrated that the HUVEC viability treated with ten antioxidant peptides (M1 to M10) at 100.0 μM for 24 h was not significantly affected compared with that of the normal group ($P < 0.05$). Furthermore, FWKVV and FMPLH at 100.0 μM could very significantly enhance the viabilities ($75.89 \pm 1.79\%$ and $70.03 \pm 4.37\%$) of oxidative-damaged HUVECs by H₂O₂ compared with those of the model group ($51.66 \pm 2.48\%$) ($P < 0.001$). The results indicated that FWKVV and FMPLH played their protective functions through increasing the levels of antioxidant enzymes including superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) and decreasing the levels of reactive oxygen species (ROS), malondialdehyde (MDA), and nitric oxide (NO) in oxidative-damaged HUVECs in a dose-dependent manner. In addition, the comet assay revealed that FWKVV and FMPLH could dose-dependently protect deoxyribonucleic acid (DNA) from oxidative damage in the HUVEC model. These results suggested that antioxidant pentapeptides (FWKVV and FMPLH) could serve as potential antioxidant additives applied in the food products, pharmaceuticals, and health supplements.

1. Introduction

Bioactive peptides (BPs) produced from food resources are short amino acid sequences with 2 to 20 residues, which are inactive in the sequence of the original proteins [1–3]. Generally, BPs are liberated by protease hydrolysis during either food processing or gastrointestinal digestion in the body [4, 5]. Based on the types of amino acids, sequences, and spatial structures, BPs can effectively perform multiple functions, such as anticancer, antioxidant, angiotensin-converting enzyme (ACE) inhibition, immunomodulatory, antithrombotic hypocholesterolemic, antibacterial, hypotensive, and hypolipidemic activities [2, 6, 7], and their beneficial func-

tions, especially on high nutritional value, health promotion, and chronic disease adjuvant treatment and prevention, are being increasingly recognized [4, 8].

Antioxidant peptides (APs) are one of the most popular types of BPs and have been isolated from animals, plants, and microorganism [9–11]. Liu et al. prepared the antioxidant peptide fraction from the processing of by-product of hazelnut, being composed of DWDPK, ETTL, ADGF, AGGF, AWDPE, and SGAF, which showed high antioxidant and protective ability against angiotensin (Ang) II-caused oxidant damage by upregulating the levels of superoxide dismutase (SOD) and heme oxygenase-1 (HO-1) and downregulating the level of xanthine oxidase-1 (XO-1) to control the

generation of reactive oxygen species (ROS) in human umbilical vein endothelial cells (HUVECs) [12]. An antioxidant peptide YD1 from *Bacillus amyloliquefaciens* with average molecular weight (MW) of ~1.0 kDa could strongly reduce the levels of ROS and nitric oxide (NO) and improve the levels of antioxidant enzymes by heightening the transcriptional and translational ability of NF-E2-related factor-2 (Nrf-2) in murine macrophages (RAW 264.7) [13]. EVSGPGLSPN from walnut showed high protection on PC12 cells against H₂O₂-caused neurotoxicity by controlling the nuclear factor kappa-B (NF- κ B)/caspase pathways and improving the activity of antioxidant enzymes, including glutathione peroxidase (GSH-Px), SOD, and catalase (CAT) [14].

At present, protein hydrolysates and BPs have been prepared from the processing of by-products of multifarious edible marine organisms, such as mollusks, fishes, and seaweeds, and many of them showed significant activities [2, 15–17]. Pan et al. found that four oligopeptides (VPR, IEPH, LEEEE, and IEEEQ) from protein hydrolysate of red stingray cartilages exhibited strong lipid peroxidation inhibition activity, Fe²⁺-chelating ability, reducing power, and radical scavenging activity [7]. Research by Oh et al. indicated that blue mussel- α -chymotrypsin hydrolysate (BMCH) showed high radical scavenging activity and inhibitory activity on Cu²⁺-mediated low-density lipoprotein (LDL) oxidation. Moreover, BMCH could improve the viability of HUVECs and lower ROS generation through downregulating gene expression of p53 and caspase-3, as well as decreasing the ratio of B-cell lymphoma 2- (bcl-2-) associated X (bax)/bcl-2 [18]. In addition, YPPAK, PIIIVYWK, TTANIEDRR, and FSVVPSPK from blue mussel hydrolysate exhibited strong radical scavenging activity, and PIIIVYWK and FSVVPSPK could activate of HO-1 gene expression to play their hepatoprotective effects [15, 19]. Chen and Hou reported that skin gelatin hydrolysate of Pacific cod could protect skin injury caused by ultraviolet (UV) radiation because the gelatin hydrolysate could restrain the decrease in endogenous antioxidant enzymes and hold back the expression of proinflammatory cytokines and NF- κ B [20]. Those studies confirm that APs could serve as antioxidant agents to inhibit the influences associated with oxidative stress in food and living organisms through enhancing the levels of endogenous antioxidant enzymes to prevent or repair the process of oxidation resulting from ROS [9, 15]. Furthermore, these APs showed excellent properties including being easy to absorb, having low MW, and being less toxic or having less side effect and become high-quality ingredients for healthy products and food processing/preservation industries [21, 22].

In our previous work, ten antioxidant peptides (M1 to M10) have been prepared from papain hydrolysate of miiuy croaker muscle and their amino acid sequences were determined to be YASVV (M1), NFWWP(M2), FWKV (M3), TWKV (M4), FMPLH (M5), YFLWP (M6), VIAPW (M7), WVWW (M8), MWKV (M9), and IRWW (M10) [3]. In which, FWKV (M3) and FMPLH (M5) could dramatically control lipid peroxidation and show powerful reducing power and radical scavenging ability. Therefore, the aim of this work was to evaluate the protective functions

of FWKV (M3) and FMPLH (M5) on HUVECs against hydrogen peroxide- (H₂O₂-) induced oxidative damage and illuminate their protective mechanism.

2. Material and Methods

2.1. Reagents. HUVECs were purchased from the Cell Bank of Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). Bicinchoninic acid (BCA), Dulbecco's modified Eagle's medium (DMEM), Hoechst 33342, ethidium bromide (EB), phosphate-buffered saline (PBS, pH 7.2), glycerol, bromophenol blue, acetylcysteine (NAC), L-glutamine, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), and dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich (Shanghai) Trading Co., Ltd. (China). The ten antioxidant peptides (M1-M10) were synthesized in China Peptides Co. (Suzhou, China), and the purity was more than 98% (*w/w*).

2.2. Culture and the Viability Assay of HUVECs. The culture method of HUVECs and the cell viability assay were carried out in accordance with the previous method of Cai et al. [23] and Lim et al. [24]. The MTT assay was replicated for three times and applied to determine the viability of HUVECs. In brief, the basal HUVECs were incubated in a 96-well plate for 24 h. After that, the basal HUVECs were cultured in the sample solutions with the concentration of 100 μ M. After 12 h, the wells were rinsed with PBS two times and the MTT was added to the wells with the final concentration of 0.5 mg/mL. After being incubated for 4 h, the active cells formed formazan crystals, which were added into 150 μ L of DMSO and well combined. Finally, the cell viability was determined according to the absorbance at 570 nm of the blank control group (A_{control}) and the sample group (A_{sample}) by the following formula:

$$\text{Cell viability (\%)} = \left(\frac{A_{\text{sample}}}{A_{\text{control}}} \right) \times 100\%. \quad (1)$$

2.3. Protection of FWKV (M3) and FMPLH (M5) on the Oxidative Damage Model of HUVECs. The oxidative damage model of HUVECs was established in accordance with the method described by Cai et al. [23]. After culturing for 24 h, the supernatant in HUVEC wells was aspirated. Subsequently, H₂O₂ was added and its final concentrations, respectively, reached 100, 200, 300, 400, and 500 mM. After 24 h, the HUVEC viability of each well was measured three times. In addition, the H₂O₂ concentration induced the cell viability; about 50% was selected to establish the oxidative damage model of HUVECs.

After establishing the oxidative damage model of HUVECs under the optimized H₂O₂ conditions, the peptide samples with the final concentrations of 10.0, 50.0, and 100.0 μ g/mL were added to the DMEM and well combined. After culturing for 24 h, the supernatant in HUVEC wells was wiped off. Afterwards, 100 μ L of the peptide samples at the given concentration was put in the protection groups. After 8 h, the peptide sample was cleared up and H₂O₂ at the given concentration was put in the model and peptide

sample groups of HUVECs and treated for 24 h. 100 μL of NAC (1.5 mM) was used in place of 100 μL of peptide sample in the positive group.

2.4. Influences of FWKV (M3) and FMPLH (M5) on the Morphology of HUVECs. The Hoechst 33342 staining assay was performed in accordance with the method described by Cai et al. [23]. After being treated by trypsinization, HUVECs were grown and incubated for 24 h in a 6-well plate. The supernatant in the 6-well plate was wiped off, and 300 μL of NAC, FWKV (M3), and FMPLH (M5) solutions was separately added into different 6-well plates. After incubating for 2 h, NAC, FWKV (M3), and FMPLH (M5) were removed and 300 μL of H_2O_2 was put in the plate separately. After incubating for 24 h, Hoechst 33342 was added into the plate at a concentration of 8 mg/mL. After incubating for 30 min, Hoechst 33342 was cleared away from the plate and HUVECs were washed using serum-free DMEM three times. The fluorescence microscope of LSM710 was employed to observe the morphology of HUVECs, and 550 nm and 460 nm were designed as the excitation and emission wavelengths, respectively.

2.5. Influences of FWKV (M3) and FMPLH (M5) on ROS, SOD, GSH-Px, NO, and MDA. The levels of ROS in blank control, model, and sample groups were measured in accordance with the method described by Cai et al. [23]. In addition, the levels of antioxidases (SOD and GSH-Px) and oxidation-related indicators (NO and MDA) were measured using assay kits in accordance with the manufacturer's protocols (Nanjing Jiancheng Bioengineering Institute Co., Ltd., China), and the BCA method was employed to determine the protein concentrations for normalizing the levels of antioxidases (SOD and GSH-Px). The level of ROS was indicated as % of blank control values. The levels of antioxidases (SOD and GSH-Px) were indicated as U/mg prot (units of enzymatic activity/mg protein).

2.6. Influence of FWKV (M3) and FMPLH (M5) on Oxidative Damage to Deoxyribonucleic Acid (DNA) by H_2O_2 . The protective ability of FWKV (M3) and FMPLH (M5) on supercoiled pBR322 plasmid DNA against H_2O_2 damage was measured in accordance with the method described by Cai et al. [23] and Zhao et al. [25]. Glutathione (GSH) served as the positive control in the experiment.

The DNA comet assay was carried out in accordance with the method described by Cai et al. [23] and applied to analyze the protective ability of FWKV (M3) and FMPLH (M5) on DNA damage in the oxidative damage model of HUVECs.

2.7. Statistical Analysis. All the assays were performed more than three times, and the resulting data were indicated as means \pm standard deviation (SD). The mean value of each treatment was analyzed using the ANOVA test (SPSS Statistics 22.0 software). Duncan's multiple range tests were applied to analyze the significant differences of different groups ($P < 0.05$, $P < 0.01$, and $P < 0.001$).

3. Results and Discussion

3.1. Influence of APs (M1-M10) on the Viability of HUVEC. HUVECs are separated from the umbilical cord vein and play a major role as a model system for the analysis of the regulation of endothelial cells and the effect of antioxidants on endothelial cells and protection from oxidative damage by peroxide, measured by cell viability and its effect on apoptosis [5, 12].

Figure 1 demonstrates the influence of APs (M1-M10) on cell viability. At the concentration of 100 μM , the viability of HUVECs incubated with IRWWW (M10) for 24 h was $93.59 \pm 6.41\%$, which was inferior to those of the blank control and other peptide groups. On the contrary, the viability of cells incubated with YFLWP (M6) was $109.39 \pm 9.39\%$, which was superior to those of the blank control and other peptide groups. However, there was no significant difference between the control group and peptide-treated groups under the same conditions ($P < 0.05$). Cell viability is a determination of living or dead cells, based on a total cell sample. Viability analysis has a tremendous number of applications, especially calculating the effect of antitumor drug candidates on cancer cells and the cytotoxic activity of chemical compound on normal cells [26–28]. Therefore, ten isolated APs (M1-M10) from miiuy croaker revealed the possibility for exploiting nontumor healthy products because they had no significant influence on the conventional proliferation of HUVECs.

In the experiment, H_2O_2 was used to produce hydroxyl radicals ($\cdot\text{OH}$) and oxygen radicals, which further induce oxidative stress in cells [27]. Herein, the oxidative damage model built with H_2O_2 was usually applied to evaluate the antioxidant activity and investigate their molecular mechanisms concerned on the pathogenesis of ROS-caused oxidative injury. Figure 2 shows that the viability of HUVECs decreased with the increase in H_2O_2 concentrations when the H_2O_2 concentrations ranged from 100 μM to 350 μM . In addition, it was found that the viability of HUVECs reduced to $51.66 \pm 2.48\%$ at the H_2O_2 concentration of 200 μM , and there was significant difference with the other tested groups ($P < 0.05$). Then, 200 μM was recognized as the optimal concentration of H_2O_2 for establishing the oxidative damage model of HUVECs.

3.2. Protective Effects of the APs (M1-M10) on the Oxidative Injury of HUVEC by H_2O_2 . The protective ability of the 10 APs (M1-M10) on the oxidative injury model of HUVEC caused by H_2O_2 is presented in Figure 3. The results indicated that the cell viabilities of M1-M10-treated groups ranged from $55.90 \pm 2.95\%$ to $75.89 \pm 1.79\%$, which were superior to that of the oxidative damage model group ($51.66 \pm 2.48\%$). However, the cell viabilities of M1-M10-treated groups were lower than that ($82.90 \pm 3.05\%$) of the positive control group (NAC). In addition, the HUVEC viability of the FWKV- (M3) and FMPLH- (M5) treated group was $75.89 \pm 1.79\%$ and $70.03 \pm 4.37\%$ at the concentration of 100.0 μM , which was very significantly ($P < 0.001$) superior to that of the oxidative damage model group ($51.66 \pm 2.48\%$); the NFWWP (M2), YFLWP (M6),




FIGURE 1: Influences of M1-M10 on the viability of HUVECs at the concentration of 100.0 μM . The experiments were in triplicate ($n = 3$).




FIGURE 2: Influences of H_2O_2 concentrations (100.0 μM to 350.0 μM) on the viability of HUVECs. The experiments were in triplicate ($n = 3$).
^{a-g}Values with the same letters indicate no significant difference ($P > 0.05$).




FIGURE 3: Influences of M1-M10 on the viability of oxidative-damaged HUVECs at the concentration of 100.0 μM . ^{###} $P < 0.001$ vs. the blank control group; ^{*} $P < 0.05$, ^{**} $P < 0.01$, and ^{***} $P < 0.001$ vs. the oxidative damage model group.

WVWWW (M8), MWKVW (M9), and IRWWW (M10) increased the HUVEC viability to $62.36 \pm 2.14\%$, $63.51 \pm 1.34\%$, $67.34 \pm 3.54\%$, $64.70 \pm 1.94\%$, and $65.01 \pm 3.52\%$, respectively, and the viabilities of those peptide-treated groups were significantly ($P < 0.05$) higher than that of the oxidative damage model group. Therefore, these results illustrated that FWKVW (M3) and FMPLH (M5) have the highest protective capability on HUVEC against oxidative damage induced by H_2O_2 among the 10 APs (M1-M10) when the concentration of APs (M1-M10) was $100.0 \mu M$.

The results in Figure 4 illustrated that the concentrations of FWKVW (M3) and FMPLH (M5) and their protective capability on the H_2O_2 -induced HUVEC injury model showed a positive relationship. The HUVEC viability in the FWKVW- (M3) incubated group increased from $55.02 \pm 2.35\%$ to $75.89 \pm 1.79\%$ when its concentration was changed from $10.0 \mu M$ and $100.0 \mu M$. Moreover, the HUVEC viability was very significantly ($P < 0.01$) and significantly ($P < 0.05$) higher than that of the oxidative damage model group at the concentrations of $50.0 \mu M$ and $100.0 \mu M$, respectively. The HUVEC viability of the FMPLH- (M5) treated group showed similar tendency at the tested concentrations. However, there was no significant difference between FWKVW- (M3) and FMPLH- (M5) incubated groups and the oxidative damage model group at the concentration of $10.0 \mu M$ ($P > 0.05$).

Images of the blank control (H_2O), oxidative damage model (H_2O_2), FWKVW (M3), FMPLH (M5), and positive control (NAC) stained with Hoechst 33342 are presented in Figure 5. After incubation for 24 h, HUVECs were plump in shape, uniform in size, and presenting blue fluorescence in the blank control group (Figure 5(a)). However, HUVECs in the model group (Figure 5(b)) showed an apoptosis state because the number of HUVECs significantly reduced, the cells became smaller, and the fluorescence of most of the remaining adherent cells became bright. Compared with the model group, Figures 5(d) and 5(e) present that most of HUVECs in FWKVW (M3) and FMPLH (M5) groups adhered to the wall. In addition, only a handful of HUVECs were washed away and other small quantity of HUVECs that presented blue fluorescence brightened. However, the protective capacity of FWKVW (M3) and FMPLH (M5) on oxidative-damaged HUVECs induced by H_2O_2 was weaker than that of the positive control. The present results suggested that FWKVW (M3) and FMPLH (M5) displayed a significant protective capacity on the H_2O_2 -damaged HUVECs, which agreed with the results found in Figure 2.

3.3. Effects of FWKVW (M3) and FMPLH (M5) on the ROS Levels in the Oxidative Damage Model of HUVECs. The ROS levels in the blank control (H_2O), oxidative damage model (H_2O_2), FWKVW (M3), FMPLH (M5), and positive control (NAC) groups are presented in Figure 6. The data demonstrated that there was an extremely ($P < 0.001$) significant difference between the blank control group and the oxidative damage model group ($231.7 \pm 12.8\%$ of blank control). The finding proved that the induced oxidative damage caused by H_2O_2 resulted in a significant increase in ROS levels in the model group. In peptide groups, the intracellular

ROS levels were significantly decreased by FWKVW (M3) and FMPLH (M5) pretreatment in a concentration-dependent manner. Moreover, FWKVW (M3) showed stronger activity of ROS scavenging than FMPLH (M5) did. At the concentrations of $10 \mu M$, $50 \mu M$, and $100 \mu M$, the intracellular ROS levels of the FWKVW- (M3) incubated group were $178.2 \pm 11.7\%$, $152.3 \pm 8.9\%$, and $133.4 \pm 9.6\%$ of the blank control group. The data indicated that FWKVW (M3) and FMPLH (M5) could play an important role in helping HUVECs from oxidative stress damage through decreasing the levels of ROS.

3.4. Effects of FWKVW (M3) and FMPLH (M5) on the Levels of Antioxidases (SOD and GSH-Px) and Oxidation-Related Indicators (NO and MDA) in the Oxidative Damage Model of HUVECs. The endogenous antioxidant enzymes, such as SOD, GSH-Px, CAT, and GSH-Rx, and nonenzymatic antioxidants form the body's endogenous defense mechanisms to protect vital biomolecules and ultimately body tissues against oxidative damage in cells through catalyzing reactions to neutralize ROS [29]. Under oxidative stress, uncontrolled generation of ROS will be beyond the normal processing capacity of this endogenous defense mechanism, and additional antioxidants are needed to help control oxidative stress damage [2, 30]. Hence, the influences of FWKVW (M3) and FMPLH (M5) on the levels of antioxidant enzymes (SOD and GSH-Px) and oxidation-related indicators (NO and MDA) in HUVECs were determined and are shown in Figure 7 for explaining their protective action on HUVECs against H_2O_2 -induced oxidative damage.

Figures 7(a) and 7(b) show that the levels of antioxidant enzymes (SOD and GSH-Px) in the H_2O_2 -induced HUVEC injury model were 111.35 ± 3.47 U/mg prot and 20.38 ± 0.82 U/mg prot, respectively, and the data were extremely significantly less than that of the normal HUVEC control ($P < 0.001$). Furthermore, Figures 7(a) and 7(b) indicate that there were positive correlations between the levels of antioxidant enzymes (SOD and GSH-Px) and the peptide (FWKVW (M3) and FMPLH (M5)) concentrations. The levels of SOD (148.94 ± 5.64 U/mg prot and 180.62 ± 4.38 U/mg prot) and GSH-Px (28.89 ± 0.96 and 38.67 ± 0.98) of HUVECs incubated with FWKVW (M3) at the concentrations of $50.0 \mu M$ and $100.0 \mu M$ were very significantly ($P < 0.01$) and significantly ($P < 0.05$) higher than those of the oxidative damage model group. At the same concentration, the influence of FMPLH (M5) on the levels of antioxidant enzymes (SOD and GSH-Px) was slightly weaker than that of FWKVW (M3). The levels of SOD (162.64 ± 2.06 U/mg prot and 167.28 ± 4.57 U/mg prot) of HUVECs incubated with FMPLH (M5) at the concentrations of $50.0 \mu M$ and $100.0 \mu M$ were significantly ($P < 0.05$) and very significantly ($P < 0.01$) higher than those of the H_2O_2 -damaged group. In addition, the levels of GSH-Px of HUVECs treated with FMPLH (M5) were 34.68 ± 1.55 U/mg prot at the concentration of $100.0 \mu M$, and the level was very significantly ($P < 0.01$) greater than that (20.38 ± 0.82 U/mg prot) of the oxidative damage model group. Therefore, the present finding indicated that FWKVW (M3) and FMPLH (M5), especially at the high concentration, had a significant influence




FIGURE 4: Influences of FWKV (M3) and FMPLH (M5) on the viability of oxidative-damaged HUVECs at the concentrations of 10.0 μ M, 50.0 μ M, and 100.0 μ M. The experiments were in triplicate ($n = 3$). ### $P < 0.001$ vs. the blank control group; * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ vs. the oxidative damage model group.

on increasing the levels of antioxidant enzymes (SOD and GSH-Px) of HUVECs to deal with the damage of oxidative stress.

ROS generated under oxidative stress can damage the cell membrane and vital macromolecules, which causes the increase in MDA and NO. As shown in Figures 7(c) and 7(d), the contents of oxidation-related indicators including NO and MDA in the oxidative damage model group were significantly higher than those (MDA, 9.67 ± 0.35 nM/mg prot; NO, 3.75 ± 0.32 μ M/L) of the blank control group ($P < 0.001$). The results suggested that H₂O₂ causes oxidative stress in HUVECs at the concentration of 200 μ M and seriously damages the cell membranes. To our excitement, there were negative correlations between the levels of MDA and NO and the peptide (FWKV (M3) and FMPLH (M5)) concentrations in Figures 7(c) and 7(d). The contents of MDA at the concentration of 100.0 μ M were 15.12 ± 0.62 nM/mg prot and 14.13 ± 0.58 nM/mg prot, respectively, and those data were significantly less than those of the oxidative damage model group ($P < 0.01$). The influences of FWKV (M3) and FMPLH (M5) on the NO content in oxidative-damaged HUVECs were more significant than those on the MDA content. The contents of NO of HUVECs incubated with FWKV (M3) and FMPLH (M5) at the concentration of 50.0 μ M were very significantly lower than those of the oxidative damage model group ($P < 0.01$). In FWKV (M3) and FMPLH (M5) treated groups, the content decrease in MDA and NO in oxidative-damaged HUVECs further proved that FWKV (M3) and FMPLH (M5) could increase the levels of antioxidant enzymes to scavenge ROS and alleviate its oxidized damage to cells.

Previous literatures indicated that some protein hydrolysates and BPs displayed intracellular antioxidant functions through controlling the levels and gene expression of antioxidant enzyme in cells and organisms [2]. Homayouni-Tabrizi

et al. reported that YLEELHRLNAGY from camel milk protein hydrolysate could significantly increase the gene expression of SOD and CAT in treated HepG2 cells [31]. Antioxidant peptides derived from *Pinctada fucata* protein played a protective role in the ultraviolet-induced photoaging of mouse skins by significantly controlling the speed of lipid peroxidation and the reduction of the activity of SOD, GSH-Px, hydroxyproline, and CAT [32]. IYVVDLR and IYVFVR from the hydrolysate of soybean protein were able to protect Caco-2 cells against H₂O₂-induced oxidative damage *via* significantly downregulating intracellular ROS generation and lipid peroxidation, statistically upregulating total reduced glutathione (GSH) synthesis, enhancing activities of CAT and GSH-Px, and suppressing ROS-mediated inflammatory responses *via* inhibiting interleukin-8 secretion ($P < 0.05$) [33]. In an *in vivo* experiment, bovine hair hydrolysates (BHP) (with a major sequence of CERPTCCEHS) [34] and oyster meat hydrolysates by Alcalase (OMA) [35] showed a strong capability to decrease the MDA level and enhance the levels of endogenous cellular antioxidant enzymes, including SOD, CAT, and GSH-Px. Those findings in the experiment suggested that FWKV (M3) and FMPLH (M5) had similar cytoprotective functions with those reported antioxidant peptides because they all could reinforce endogenous antioxidant defense systems to help cells against the H₂O₂ damage.

3.5. Protective Capacities of FWKV (M3) and FMPLH (M5) on DNA

3.5.1. Protective Capacities on Plasmid DNA. The superfluous ROS produced in oxidative stress can increase the level of the oxidative DNA damage, which is further implicated in the development of a variety of cancers including colon, breast, and prostate cancer [25, 36, 37]. Therefore, the protective




FIGURE 5: Influences of FWKV (M3) and FMPLH (M5) on the apoptosis of oxidative-damaged HUVECs at the concentration of 100.0 μM .

effects of FWKV (M3) and FMPLH (M5) on plasmid DNA (pBR322 DNA) against oxidative injury by H_2O_2 are displayed in Figure 8. It was found that the plasmid DNA (pBR322 DNA) is principally of the supercoiled (SC) form in normal circumstances (Figure 8(a)). A relaxed open circular (OC) form is produced when one of the phosphodiester chains of the plasmid DNA is cleaved when undergoing oxidative stress and further cut open surrounding the first breaking, which leads to generating the linear (LIN) double-stranded DNA molecules. The LIN and OC formations of DNA are characteristic of the double-strand and single-strand breaks, respectively [25].

In the experiment, the pBR322 DNA was cleaved by the hydroxyl radicals generated from the decomposition of H_2O_2 mediated by Fe^{2+} , which changed the SC form into the OC form (Figure 8(h)). In addition, the trace LIN form of pBR322 DNA was discovered as shown in Figure 8(h),

which suggested that a minute amount of the double strand of DNA was cut open by the superfluous hydroxyl radicals. As shown in Figures 8(b)–8(g), FWKV (M3) and FMPLH (M5) positively affected the amount of the SC form of pBR322 DNA in a dose-dependent manner when their concentrations ranged from 0.5 mg/mL to 2.0 mg/mL. Correspondingly, the OC form of pBR322 DNA was gradually reduced with the increase in the concentrations of FWKV (M3) and FMPLH (M5). However, the amount of the SC form of the plasmid DNA (pBR322 DNA) in the FWKV (M3) and FMPLH (M5) group at the concentration of 2.0 mg/mL was less than that of the positive control (GSH) group (Figure 8(i)), which suggested that the protective effects on oxidative DNA damage of FWKV (M3) and FMPLH (M5) were weaker than that of the positive control (GSH). Therefore, FWKV (M3) and FMPLH (M5) might inhibit the chemical reaction of Fe^{2+} with H_2O_2 and/or





FIGURE 6: Influences of FWKV (M3) and FMPLH (M5) on the levels of ROS in oxidative-damaged HUVECs at the concentrations of 10.0 μM, 50.0 μM, and 100.0 μM. The experiments were in triplicate ($n = 3$). ### $P < 0.001$ vs. the blank control group; ** $P < 0.01$ and *** $P < 0.001$ vs. the oxidative damage model group.

directly clear away generated hydroxyl radicals through giving out the hydrogen atom or electron to play their defensive functions on the supercoiled plasmid DNA. This result agreed with our previous report that FWKV (M3) and FMPLH (M5) had strong ferric reducing power and could effectively scavenge hydroxyl radical *in vitro* [3].


3.5.2. Protective Capacities on DNA against H₂O₂-Induced Injury in the HUVEC Model. Due to the imbalance between the oxidants and antioxidants, redundant ROS is generated and then attacks DNA in cells, which further leads to adverse physiological and biochemical reactions. At present, several methods are applied to estimate the DNA damage in cells, and the comet assay is believed to be a fairly simple, sensitive, and versatile way in single cells to quantitatively and qualitatively evaluate the effects of DNA damage and DNA repair [37]. In the assay, an electrical current is applied to a slide featuring the cell in a gel. The broken strands of DNA will get away from their initial position towards the positive charge and leave a comet trail, which will be easily observable through fluorescence [38]. The comet trail can give out many quantitative data on the degree of DNA damage, which can serve to evaluate the protective effects of antioxidant compounds on DNA.

In the blank control group, the comet with a bright head and almost without comet tail (Figure 9(a)) indicated that the DNA in HUVEC was in a normal state, but the comet with a long and large tail in the oxidative damage model group (Figure 9(b)) suggested that the DNA was seriously injured by H₂O₂ and the H₂O₂-induced oxidative damage model of HUVECs was successfully established. In comparison with the oxidative damage model group, the length and area of comet tails on oxidative-damaged HUVECs were step by step diminished with the increase in the concentrations of FWKV (M3) and FMPLH (M5) from 50.0 μM to 200.0 μM (Figures 9(d)–9(i)). Even more interesting was that the length and area of comet tails in FWKV- (M3) and FMPLH- (M5) incubated groups at the concentration of 200 μM were approximately equal to the length and area of the NAC-treated group (Figure 9(c)).


The images in Figure 9 on the comet tail length or comet area could display visually the experimental results. Moreover, the comet assay had employed some indexes including torque class indicator (Olive tail moment, OTM), head DNA (HDNA), and tail DNA (TDNA) for being more conducive to accurate analysis of the results (Tables 1 and 2). These indexes demonstrated that the HDNA was extremely prominently lower than that of the blank control group ($P < 0.001$); the tail moment (TM), tail length (TL), OTM, and TDNA in the oxidative damage model group were extremely prominently higher than those in the blank control group ($P < 0.001$), and the comet length (CL) in the oxidative damage model group was significantly higher than that of the blank control group ($P < 0.05$). With the increase in the concentration of FWKV (M3) and FMPLH (M5) from 50.0 μM to 200.0 μM, the TDNA, CL, TL, TM, and OTM decreased gradually, but the HDNA of the comet increased step by step. Furthermore, all the measured indicators of FWKV (M3) and FMPLH (M5) groups at the concentrations of 100 and 200 μM, except the CL of the FMPLH (M5) group at 100 μM, showed extremely significant differences with those of the model groups ($P < 0.001$). At the concentration of 50 μM, the HDNA, TL, and TM of FWKV (M3) and FMPLH (M5) groups showed very significant ($P < 0.01$) differences with those of the model groups, respectively; the TDNA and OTM of FWKV (M3) and FMPLH (M5) groups showed very significant differences with those of the model groups, respectively ($P < 0.01$); the OTM of FWKV (M3) and FMPLH (M5) groups showed significant ($P < 0.05$) differences with that of the model groups, respectively, but the CL of FWKV (M3) and FMPLH (M5) groups showed no significant differences with that of the oxidative damage model groups ($P > 0.05$). In consequence, the results of the comet assay demonstrated that FWKV (M3) and FMPLH (M5) showed significantly protective capacity on DNA in the oxidative damage model groups induced by H₂O₂. In addition, FWKV (M3) showed stronger protective effect on DNA in the H₂O₂-induced HUVEC injury model than FMPLH (M5) did at the same concentrations.



(a)



(b)



(c)

FIGURE 7: Continued.




FIGURE 7: Influences of FWKVV (M3) and FMPLH (M5) on the levels of SOD (a), GSH-Px (b), MDA (c), and NO (d) in oxidative-damaged HUVECs at the concentrations of 10.0 μ M, 50.0 μ M, and 100.0 μ M. The experiments were in triplicate ($n = 3$). ### $P < 0.001$ vs. the blank control group; *** $P < 0.001$, ** $P < 0.01$, and * $P < 0.05$ vs. the oxidative damage model group.




FIGURE 8: Protective capacities of FWKVV (M3) and FMPLH (M5) on plasmid pBR322 DNA against H₂O₂-induced oxidative damage. A: pBR322 DNA; B: pBR322 DNA+FeSO₄+FWKVV (M3) (2.0 mg/mL)+H₂O₂; C: pBR322 DNA+FeSO₄+FWKVV (M3) (1.0 mg/mL)+H₂O₂; D: pBR322 DNA+FeSO₄+FWKVV (M3) (0.5 mg/mL)+H₂O₂; E: pBR322 DNA+FeSO₄+FMPLH (M5) (2.0 mg/mL)+H₂O₂; F: pBR322 DNA+FeSO₄+FMPLH (M5) (1.0 mg/mL)+H₂O₂; G: pBR322 DNA+FeSO₄+FMPLH (M5) (0.5 mg/mL)+H₂O₂; H: pBR322 DNA+FeSO₄+H₂O₂; and I: pBR322 DNA+FeSO₄+GSH (2.0 mg/mL)+H₂O₂.

Attack of DNA by ROS produced under conditions of oxidative stress can induce strand breaks, DNA-DNA and DNA-protein crosslinking, and formation of at least 20 modified bases adducts, which play a key role in aging, cancer, arteriosclerosis, neurodegenerative diseases, and diabetes [39, 40]. The antioxidant hydrolysate and BPs are stage by stage accepted as food components applied in functional food and nutraceuticals to effectively adjust and control the oxidative injury to protect DNA, lipid, and protein in the human body [41–43]. Hoki frame protein hydrolysate (APHPH) could decrease *t*-butyl hydroperoxide-caused cytotoxicity on human embryonic lung fibroblasts and observably protect DNA against the free radical-induced injury [44]. Similarly, protein hydrolysates prepared from Nile tilapia [45] and shrimp shell [46] could effectively inhibit the DNA scission caused by H₂O₂ and peroxy radical in a dose-dependent manner. YGDEY could protect HepG2 cells against alcohol-caused oxidative damage by controlling oxidative stress including reducing the degree of DNA damage. The antioxidant mechanism might be bound up with the Akt/NF- κ B/MAPK signal transduction pathways [47]. Sheih

reported that VECYG (VG5) presented the protective ability on DNA against oxidative injury, so it further reinforced the capacity of the APs to protect hydroxyl radical-caused injury [48]. WAFAPA and MYPGLA prepared from the protein hydrolysate of the blue-spotted stingray were superior to carnosine in their capacity to control lipid oxidation caused by H₂O₂. In addition, WAFAPA and MYPGLA could help plasmid DNA against oxidative injury caused by Fenton's reagent [21]. In our previous report, FWKVV (M3) and FMPLH (M5) have been found to have strong lipid peroxidation inhibition, reducing power, and radical scavenging activities [3], and the current study finding demonstrated that FWKVV (M3) and FMPLH (M5) could weaken the oxidative injury of HUVECs caused by H₂O₂ through enhancing the contents of endogenous antioxidants (SOD and GSH-Px), lowering the contents of oxidation-related indicators (NO and MDA), and helping DNA from oxidative injury.

4. Discussion

Oxidative stress contributes to cell pathogenesis [49], ROS-mediated damage to DNA [39, 40], and an altered mitochondrial function [50]. The oxidative damage to DNA is considered a key factor in vascular disorder [51, 52]. Our results showed that the cell viability was significantly decreased, accompanied by DNA damage detected by the DNA comet assay and plasmid DNA assay after being cultured in 200 μ M H₂O₂ medium for 24 h, while H₂O₂-induced cytotoxicity and DNA damage were significantly inhibited when HUVECs were cocubated with FWKVV (M3) or FMPLH (M5). The study also found that the exposure of HUVECs to 200 μ M H₂O₂ for 24 h led to a dramatic increase in ROS production and induced apoptosis. However, treatment with M3 and M5 together with 200 μ M H₂O₂ markedly inhibited ROS formation and diminished DNA damage and cell apoptosis in HUVECs. These data suggest that M3 and M5 have ameliorative effects on H₂O₂-induced endothelial damage in HUVECs.




FIGURE 9: Protective capacities of FWKVV (M3) and FMPLH (M5) on DNA against H_2O_2 -induced oxidative damage in HUVECs: (a) the blank control group; (b) the oxidative damage model group (H_2O_2 , 200.0 μM); (c) positive control (NAC) group; (d) H_2O_2 +FWKVV (M3) (200.0 μM); (e) H_2O_2 +FWKVV (M3) (100.0 μM); (f) H_2O_2 +FWKVV (M3) (50.0 μM); (g) H_2O_2 +FMPLH (M5) (200.0 μM); (h) H_2O_2 +FMPLH (M5) (100.0 μM); (i) H_2O_2 +FMPLH (M5) (50.0 μM).

Excessive oxides deplete tissues of GSH and impair antioxidant defense systems in humans [53, 54]. How to increase the level of GSH becomes important. In this experiment, we observed that the levels of GSH in HUVECs had a marked increase in the groups with the treatment with M3 or M5. The results for the potential mechanism of the promoting production of GSH by M3 or M5 might contribute to a broader biological effect of a protective nature with regard to the general detoxification of environmental agents. No evidence was found that GSH was able to pass through cell membranes freely; thus, M3 or M5 must modulate GSH levels by stimulating cells to synthesize GSH intracellularly. The results indicate that M3 or M5 does protect cells from oxidative stress damage, and the protection includes its capacity to stimulate GSH synthesis.

Cells are in a stable state known as redox homeostasis under normal physiological conditions. Redox homeostasis is maintained by the balance between continuous ROS generation and several mechanisms involved in antioxidant activity [30, 55]. An overwhelming production of ROS leads to a prooxidant state also known as oxidative stress, which is a leading factor in the pathogenesis of vascular disorder complications [2, 56, 57]. NOX is a major contributor to ROS

generation in endothelial cells [58]. Discrete compartmental redox signaling is demonstrated by Nrf2-dependent activation of ARE. Nrf2 is a redox-sensitive transcription factor that is activated by an oxidative signal in the cytoplasm. The Nrf2 pathway is regarded as the most important in the cell to protect against oxidative stress [40, 59]. Upon extracellular stimulation, the nucleus translocation of Nrf2 induces the transcription of several antioxidant genes, such as NADPH quinone oxidoreductase-1 (NQO-1), heme oxygenase 1 (HO-1), and SOD [60–62]. SOD and GSH-Px are two important antioxidant enzymes that remove toxic free radicals [29, 47]. Our study revealed that intracellular ROS were significantly increased when HUVECs were cultured in 200 μM H_2O_2 media. Treatment with M3 or M5 reduced H_2O_2 -induced ROS production. The present study also showed that SOD and GSH-Px activity markedly decreased in HUVECs and increased the levels of the key cytotoxic lipid peroxide MDA and NO production when they were exposed to 200 μM H_2O_2 for 24 h, which was blocked by M3 or M5 cotreatments. Major intracellular antioxidant defenses are the GSH pool and ROS-scavenging enzymes such as GSH-Px and SOD. Increased levels or activation of these endogenous antioxidants or enzymes has been shown to protect cells

TABLE 1: Protective capacity of FWKV (M3) on DNA against H₂O₂-induced oxidative damage in HUVECs.

Group	Blank control	Model (H ₂ O ₂)	Positive control (NAC)	FWKV (M3, μ M)		
				200	100	50
Cell number (<i>n</i>)	104	117	121	113	109	124
HDNA (%)	91.9 ± 4.2	18.2 ± 3.2 ^{###}	80.7 ± 3.8 ^{***}	69.6 ± 6.9 ^{***}	46.8 ± 3.3 ^{***}	32.2 ± 2.1 ^{***}
TDNA (%)	8.1 ± 4.2	81.8 ± 3.2 ^{###}	19.3 ± 4.8 [*]	30.4 ± 6.9 ^{***}	53.2 ± 2.5 ^{***}	67.8 ± 3.0 ^{**}
CL (pix)	68.3 ± 3.3	77.3 ± 13.5 [#]	66.2 ± 3.3 ^{***}	63.0 ± 3.2 ^{***}	66.6 ± 7.3 ^{***}	72.0 ± 5.5
TL (pix)	7.3 ± 0.33	59.3 ± 6.6 ^{###}	27.2 ± 1.6 ^{***}	29.6 ± 3.3 ^{***}	33.7 ± 2.5 ^{***}	47.7 ± 3.1 ^{***}
TM	0.6 ± 0.03	46.4 ± 8.1 ^{###}	5.2 ± 0.24 ^{***}	10.9 ± 1.2 ^{***}	18.0 ± 2.4 ^{***}	39.4 ± 2.8 ^{***}
OTM	2.3 ± 0.1	24.5 ± 4.3 ^{###}	4.7 ± 0.2 ^{***}	8.7 ± 1.0 ^{***}	12.6 ± 1.7 ^{***}	21.1 ± 1.5 ^{**}

All data are presented as the mean ± SD of triplicate results (*n* = 3). [#]*P* < 0.05 and ^{###}*P* < 0.001 vs. the blank control group; ^{*}*P* < 0.05, ^{**}*P* < 0.01, and ^{***}*P* < 0.001 vs. the oxidative damage model group.

TABLE 2: Protective capacity of FMPLH (M5) on DNA against H₂O₂-induced oxidative damage in HUVECs.

Group	Blank control	Model (H ₂ O ₂)	Positive control (NAC)	FMPLH (M5, μ M)		
				200	100	50
Cell number (<i>n</i>)	104	117	121	103	114	106
HDNA (%)	91.9 ± 4.2	18.2 ± 3.2 ^{###}	80.7 ± 3.8 ^{***}	62.3 ± 6.9 ^{***}	44.7 ± 5.4 ^{***}	31.6 ± 3.8
TDNA (%)	8.1 ± 4.2	81.8 ± 3.2 ^{###}	19.3 ± 4.8 [*]	37.7 ± 6.9 ^{***}	55.3 ± 5.4 ^{***}	68.4 ± 8.3 ^{**}
CL (pix)	68.3 ± 3.3	77.3 ± 13.5 [#]	66.2 ± 3.3 ^{***}	62.0 ± 7.3 ^{***}	70.4 ± 6.5 [*]	76 ± 6.2
TL (pix)	7.3 ± 0.33	59.3 ± 6.6 ^{###}	27.2 ± 1.6 ^{***}	35.4 ± 3.3 ^{***}	39.0 ± 2.9 ^{***}	51.7 ± 4.6 ^{**}
TM	0.6 ± 0.03	46.4 ± 8.1 ^{###}	5.2 ± 0.24 ^{***}	16.7 ± 1.6 ^{***}	21.6 ± 2.6 ^{***}	40.9 ± 2.3 ^{**}
OTM	2.3 ± 0.1	24.5 ± 4.3 ^{###}	4.7 ± 0.2 ^{***}	11.5 ± 1.1 ^{***}	13.1 ± 1.0 ^{***}	22.6 ± 1.6 [*]

All data are presented as the mean ± SD of triplicate results (*n* = 3). [#]*P* < 0.05 and ^{###}*P* < 0.001 vs. the blank control group; ^{*}*P* < 0.05, ^{**}*P* < 0.01, and ^{***}*P* < 0.001 vs. the oxidative damage model group.

against oxidative damage. In this study, a marked decrease indeed occurred in the level of GSH after the treatment with H₂O₂; as a result, the decreased levels of GSH might be attributed to the less decreased activity of GSH-Px. The imbalance of the H₂O₂-induced antioxidant status in HUVECs, such as the changes in the activity of antioxidant enzymes and the depletion of GSH, might be major causes of cell injury. Nevertheless, the cytoprotective effects of M3 or M5 may be mediated, in part, by activation of Nrf2, a redox-regulated transcription factor that binds to the antioxidant response element (ARE). These data suggest that H₂O₂-induced ROS production exceeds the natural antioxidant capacity and leads to oxidative stress and that M3 or M5 may exert its protective effect on HUVECs by reducing ROS accumulation and increasing antioxidant enzyme activity.

Finally, in this study, we demonstrated that FWKV (M3) and FMPLH (M5) could significantly protect cells against H₂O₂-induced oxidative stress, DNA damage, and cell apoptosis in cultured HUVECs. Treatment with M3 or M5 reduced the loss of cell viability, improved the antioxidant capacity, and reduced the oxidative damage in HUVECs. Further, we hypothesized that the protective effect of M3 or M5 appears to be mediated by the regulation of the Nrf2 signaling pathway and the maintenance of cellular redox homeostasis. This finding further supports the concept that M3 or M5 may have potential as a vascular protective drug.

5. Conclusion

In this work, the protective activity and antioxidation mechanisms of FWKV (M3) and FMPLH (M5) on HUVECs against H₂O₂-caused oxidative injury were researched carefully. When incubated for 24 h, no significant difference was found on the viability of HUVECs between the blank control group and the AP (M1-M10) groups at 100 μ M (*P* < 0.05). Moreover, FWKV (M3) AND FMPLH (M5) could significantly protect HUVECs from H₂O₂-induced oxidative damage by enhancing the contents of endogenous antioxidant enzymes, lowering the levels of oxidative products, and helping DNA from oxidative injury. These results suggested that FWKV (M3) and FMPLH (M5) could serve as great potential antioxidants in the health care products. In addition, the molecular mechanism and *in vivo* antioxidant experiments of FWKV (M3) and FMPLH (M5) will also be carried out in our laboratory.

Data Availability

The datasets are available from the corresponding author on reasonable request.

Conflicts of Interest

The authors declare that there is no conflict of interest.

Authors' Contributions

BW and CFC designed the experiment. YZW, YMW, and XP performed the experiment and analyzed the experimental data. YZW, CFC, and BW wrote and revised the manuscript. All authors critically reviewed the manuscript for intellectual content and gave the final approval for the version to be published.

Acknowledgments

This work was funded by the National Natural Science Foundation of China (No. 82073764) and Ten-thousand Talents Plan of Zhejiang Province (No. 2019R52026).

References

- [1] A. Pihlanto and H. Korhonen, "Bioactive peptides and proteins," *Advances in Food and Nutrition Research*, vol. 47, pp. 175–276, 2003.
- [2] A. Sila and A. Bougatef, "Antioxidant peptides from marine by-products: Isolation, identification and application in food systems. A review," *Journal of Functional Foods*, vol. 21, pp. 10–26, 2016.
- [3] Y. He, X. Pan, C. F. Chi, K. L. Sun, and B. Wang, "Ten new pentapeptides from protein hydrolysate of miiuy croaker (*Miichthys miiuy*) muscle: Preparation, identification, and antioxidant activity evaluation," *LWT- Food Science and Technology*, vol. 105, pp. 1–8, 2019.
- [4] P. A. Harnedy, M. B. O'Keeffe, and R. J. FitzGerald, "Fractionation and identification of antioxidant peptides from an enzymatically hydrolysed *Palmaria palmata* protein isolate," *Food Research International*, vol. 100, Part 1, pp. 416–422, 2017.
- [5] Y. Q. Zhao, L. Zhang, J. Tao, C. F. Chi, and B. Wang, "Eight antihypertensive peptides from the protein hydrolysate of Antarctic krill (*Euphausia superba*): Isolation, identification, and activity evaluation on human umbilical vein endothelial cells (HUVECs)," *Food Research International*, vol. 121, pp. 197–204, 2019.
- [6] X. Pan, Y. Q. Zhao, F. Y. Hu, and B. Wang, "Preparation and identification of antioxidant peptides from protein hydrolysate of skate (*Raja porosa*) cartilage," *Journal of Functional Foods*, vol. 25, pp. 220–230, 2016.
- [7] X. Y. Pan, Y. M. Wang, L. Li, C. F. Chi, and B. Wang, "Four antioxidant peptides from protein hydrolysate of red stingray (*Dasyatis akajei*) cartilages: isolation, identification, and *in vitro* activity evaluation," *Marine Drugs*, vol. 17, no. 5, article 263, 2019.
- [8] C. F. Chi, B. Wang, Y. M. Wang, B. Zhang, and S. G. Deng, "Isolation and characterization of three antioxidant peptides from protein hydrolysate of bluefin leatherjacket (*Navodon septentrionalis*) heads," *Journal of Functional Foods*, vol. 12, pp. 1–10, 2015.
- [9] J. M. Lorenzo, P. E. S. Munekata, B. Gómez et al., "Bioactive peptides as natural antioxidants in food products - a review," *Trends in Food Science & Technology*, vol. 79, pp. 136–147, 2018.
- [10] J. Sheng, X. Yang, J. Chen et al., "Antioxidative effects and mechanism study of bioactive peptides from defatted walnut (*Juglans regia* L.) meal hydrolysate," *Journal of Agricultural and Food Chemistry*, vol. 67, no. 12, pp. 3305–3312, 2019.
- [11] M. Tu, S. Cheng, W. Lu, and M. Du, "Advancement and prospects of bioinformatics analysis for studying bioactive peptides from food-derived protein: sequence, structure, and functions," *TrAC-Trends in Analytical Chemistry*, vol. 105, pp. 7–17, 2018.
- [12] C. Liu, D. Ren, J. Li et al., "Cytoprotective effect and purification of novel antioxidant peptides from hazelnut (*C. heterophylla* Fisch) protein hydrolysates," *Journal of Functional Foods*, vol. 42, pp. 203–215, 2018.
- [13] M. S. Rahman, Y. Hee Choi, Y. Seok Choi, M. B. Alam, S. Han Lee, and J. Cheol Yoo, "A novel antioxidant peptide, purified from *Bacillus amyloliquefaciens*, showed strong antioxidant potential via Nrf-2 mediated heme oxygenase-1 expression," *Food Chemistry*, vol. 239, pp. 502–510, 2018.
- [14] C. Liu, Y. Guo, F. Zhao et al., "Potential mechanisms mediating the protective effects of a peptide from walnut (*Juglans mandshurica* Maxim.) against hydrogen peroxide induced neurotoxicity in PC12 cells," *Food & Function*, vol. 10, no. 6, pp. 3491–3501, 2019.
- [15] B. Wang, L. Li, C. F. Chi, J. H. Ma, H. Y. Luo, and Y. F. Xu, "Purification and characterisation of a novel antioxidant peptide derived from blue mussel (*Mytilus edulis*) protein hydrolysate," *Food Chemistry*, vol. 138, no. 2–3, pp. 1713–1719, 2013.
- [16] C. B. Ahn, J. G. Kim, and J. Y. Je, "Purification and antioxidant properties of octapeptide from salmon byproduct protein hydrolysate by gastrointestinal digestion," *Food Chemistry*, vol. 147, pp. 78–83, 2014.
- [17] J. B. Zhang, Y. Q. Zhao, Y. M. Wang, C. F. Chi, and B. Wang, "Eight Collagen peptides from hydrolysate fraction of Spanish mackerel Skins: isolation, identification, and In Vitro antioxidant activity Evaluation," *Marine Drugs*, vol. 17, no. 4, article 224, 2019.
- [18] Y. Oh, C. B. Ahn, K. H. Nam, Y. K. Kim, N. Y. Yoon, and J. Y. Je, "Amino acid composition, antioxidant, and cytoprotective effect of blue mussel (*Mytilus edulis*) hydrolysate through the inhibition of caspase-3 activation in oxidative stress-mediated endothelial cell injury," *Marine Drugs*, vol. 17, no. 2, article 135, 2019.
- [19] S. Y. Park, Y. S. Kim, C. B. Ahn, and J. Y. Je, "Partial purification and identification of three antioxidant peptides with hepatoprotective effects from blue mussel (*Mytilus edulis*) hydrolysate by peptic hydrolysis," *Journal of Functional Foods*, vol. 20, pp. 88–95, 2016.
- [20] T. Chen and H. Hou, "Protective effect of gelatin polypeptides from Pacific cod (*Gadus macrocephalus*) against UV irradiation-induced damages by inhibiting inflammation and improving transforming growth factor- β /Smad signaling pathway," *Journal of Photochemistry and Photobiology B Biology*, vol. 162, pp. 633–640, 2016.
- [21] F. C. Wong, J. Xiao, M. G. L. Ong et al., "Identification and characterization of antioxidant peptides from hydrolysate of blue-spotted stingray and their stability against thermal, pH and simulated gastrointestinal digestion treatments," *Food Chemistry*, vol. 271, pp. 614–622, 2019.
- [22] Y. Li, J. Li, S. J. Lin, Z. S. Yang, and H. X. Jin, "Preparation of antioxidant peptide by Microwave-Assisted hydrolysis of collagen and its protective effect against H₂O₂-induced damage of RAW264.7 cells," *Marine Drugs*, vol. 17, no. 11, article 642, 2019.
- [23] S. Y. Cai, Y. M. Wang, Y. Q. Zhao, C. F. Chi, and B. Wang, "Cytoprotective effect of antioxidant pentapeptides from the protein hydrolysate of swim bladders of miiuy croaker

- (*Miichthys miiuy*) against H₂O₂-mediated human umbilical vein endothelial cell (HUVEC) injury,” *International Journal of Molecular Sciences*, vol. 20, no. 21, p. 5425, 2019.
- [24] S. Lim, A. H. Choi, M. Kwon et al., “Evaluation of antioxidant activities of various solvent extract from *Sargassum serratifolium* and its major antioxidant components,” *Food Chemistry*, vol. 278, pp. 178–184, 2019.
- [25] Y. Q. Zhao, L. Zeng, Z. S. Yang, F. F. Huang, G. F. Ding, and B. Wang, “Anti-fatigue effect by peptide fraction from protein hydrolysate of croceine croaker (*Pseudosciaena crocea*) swim bladder through inhibiting the oxidative reactions including DNA damage,” *Marine Drugs*, vol. 14, no. 12, p. 221, 2016.
- [26] L. J. Gómez, N. A. Gómez, J. E. Zapata, G. López-García, A. Cilla, and A. Alegría, “In-vitro antioxidant capacity and cytoprotective/cytotoxic effects upon Caco-2 cells of red tilapia (*Oreochromis spp.*) viscera hydrolysates,” *Food Research International*, vol. 120, pp. 52–61, 2019.
- [27] J. Tao, Y. Q. Zhao, C. F. Chi, and B. Wang, “Bioactive peptides from cartilage protein hydrolysate of spotless smoothhound and their antioxidant activity *in vitro*,” *Marine Drugs*, vol. 16, no. 4, article 100, 2018.
- [28] S. Y. Lee and S. J. Hur, “Neuroprotective effects of different molecular weight peptide fractions obtained from beef by hydrolysis with commercial enzymes in SH-SY5Y cells,” *Food Research International*, vol. 121, pp. 176–184, 2019.
- [29] O. M. Ighodaro and O. A. Akinloye, “First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): their fundamental role in the entire antioxidant defence grid,” *Alexandria Journal of Medicine*, vol. 54, no. 4, pp. 287–293, 2018.
- [30] W. H. Zhao, Q. B. Luo, X. Pan, C. F. Chi, K. L. Sun, and B. Wang, “Preparation, identification, and activity evaluation of ten antioxidant peptides from protein hydrolysate of swim bladders of miiuy croaker (*Miichthys miiuy*),” *Journal of Functional Foods*, vol. 47, pp. 503–511, 2018.
- [31] M. Homayouni-Tabrizi, A. Asoodeh, and M. Soltani, “Cytotoxic and antioxidant capacity of camel milk peptides: effects of isolated peptide on superoxide dismutase and catalase gene expression,” *Journal of Food and Drug Analysis*, vol. 25, no. 3, pp. 567–575, 2017.
- [32] Y. Wu, Q. Tian, L. Li et al., “Inhibitory effect of antioxidant peptides derived from *Pinctada fucata* protein on ultraviolet-induced photoaging in mice,” *Journal of Functional Foods*, vol. 5, no. 2, pp. 527–538, 2013.
- [33] Q. Zhang, X. Tong, Y. Li et al., “Purification and characterization of antioxidant peptides from Alcalase-hydrolyzed soybean (*Glycine max* L.) hydrolysate and their cytoprotective effects in human intestinal Caco-2 cells,” *Journal of Agricultural and Food Chemistry*, vol. 67, no. 20, pp. 5772–5781, 2019.
- [34] W. C. Zeng, Q. Sun, W. H. Zhang, X. P. Liao, and B. Shi, “Antioxidant activity *in vivo* and biological safety evaluation of a novel antioxidant peptide from bovine hair hydrolysates,” *Process Biochemistry*, vol. 56, pp. 193–198, 2017.
- [35] Z. Zhang, G. Su, F. Zhou, L. Lin, X. Liu, and M. Zhao, “Alcalase-hydrolyzed oyster (*Crassostrea rivularis*) meat enhances antioxidant and aphrodisiac activities in normal male mice,” *Food Research International*, vol. 120, pp. 178–187, 2019.
- [36] M. S. Cooke, M. D. Evans, M. Dizdaroglu, and J. Lunec, “Oxidative DNA damage: mechanisms, mutation, and disease,” *FASEB Journal*, vol. 17, no. 10, pp. 1195–1214, 2003.
- [37] V. Gunasekarana, G. V. Raj, and P. Chand, “A comprehensive review on clinical applications of comet assay,” *Journal of Clinical and Diagnostic Research*, vol. 9, no. 3, pp. GE01–GE05, 2015.
- [38] N. P. Singh, “The comet assay: reflections on its development, evolution and applications,” *Mutation Research-Reviews in Mutation Research*, vol. 767, pp. 23–30, 2016.
- [39] M. A. Lovell and W. R. Markesbery, “Oxidative DNA damage in mild cognitive impairment and late-stage Alzheimer’s disease,” *Nucleic Acids Research*, vol. 35, no. 22, pp. 7497–7504, 2007.
- [40] Y. Liang, B. Dong, N. Pang, and J. Hu, “ROS generation and DNA damage contribute to abamectin-induced cytotoxicity in mouse macrophage cells,” *Chemosphere*, vol. 234, pp. 328–337, 2019.
- [41] R. Liu, L. Xing, Q. Fu, G. H. Zhou, and W. G. Zhang, “A review of antioxidant peptides derived from meat muscle and by-products,” *Antioxidants (Basel)*, vol. 5, no. 3, p. 32, 2016.
- [42] L. Zhang, G. X. Zhao, Y. Q. Zhao, Y. T. Qiu, C. F. Chi, and B. Wang, “Identification and active evaluation of antioxidant peptides from protein hydrolysates of skipjack tuna (*Katsuwonus pelamis*) head,” *Antioxidants (Basel)*, vol. 8, no. 8, article 318, 2019.
- [43] A. Sannasimuthu and J. Arockiaraj, “Intracellular free radical scavenging activity and protective role of mammalian cells by antioxidant peptide from thioredoxin disulfide reductase of *Arthrospira platensis*,” *Journal of Functional Foods*, vol. 61, article 103513, 2019.
- [44] S. Y. Kim, J. Y. Je, and S. K. Kim, “Purification and characterization of antioxidant peptide from hoki (*Johnius belengerii*) frame protein by gastrointestinal digestion,” *Journal of Nutritional Biochemistry*, vol. 18, no. 1, pp. 31–38, 2007.
- [45] S. Yarnpakdee, S. Benjakul, H. G. Kristinsson, and H. E. Bakken, “Preventive effect of Nile tilapia hydrolysate against oxidative damage of HepG2 cells and DNA mediated by H₂O₂ and AAPH,” *Journal of Food Science and Technology*, vol. 52, no. 10, pp. 6194–6205, 2015.
- [46] P. Ambigaipalan and F. Shahidi, “Bioactive peptides from shrimp shell processing discards: antioxidant and biological activities,” *Journal of Functional Foods*, vol. 34, pp. 7–17, 2017.
- [47] M. F. Chen, F. Gong, Y. Y. Zhang et al., “Preventive effect of YGDEY from tilapia fish skin gelatin hydrolysates against alcohol-induced damage in HepG2 cells through ROS-mediated signaling pathways,” *Nutrients*, vol. 11, no. 2, p. 392, 2019.
- [48] I.-C. Sheih, “The antioxidant properties of a new antioxidative peptide (VG5) in different oxidation system,” *Journal of Nutrition & Food Sciences*, vol. 8, pp. 546–553, 2016.
- [49] H. Z. Pan, H. Zhang, D. Chang, H. Li, and H. Sui, “The change of oxidative stress products in diabetes mellitus and diabetic retinopathy,” *British Journal of Ophthalmology*, vol. 92, no. 4, pp. 548–551, 2008.
- [50] R. A. Kowluru and P. S. Chan, “Oxidative stress and diabetic retinopathy,” *Experimental Diabetes Research*, vol. 2007, Article ID 43603, 12 pages, 2007.
- [51] R. A. Kowluru, “Diabetic retinopathy: mitochondrial dysfunction and retinal capillary cell death,” *Antioxidants & Redox Signaling*, vol. 7, no. 11–12, pp. 1581–1587, 2005.
- [52] R. A. Kowluru, L. Atasi, and Y. S. Ho, “Role of mitochondrial superoxide dismutase in the development of diabetic

- retinopathy,” *Investigative Ophthalmology & Visual Science*, vol. 47, no. 4, pp. 1594–1599, 2006.
- [53] G. Bounous, “Whey protein concentrate (WPC) and glutathione modulation in cancer treatment,” *Anticancer Research*, vol. 20, no. 6C, pp. 4785–4792, 2000.
- [54] L. C. Lands, V. L. Grey, and A. A. Smountas, “Effect of supplementation with a cysteine donor on muscular performance,” *Journal of Applied Physiology*, vol. 87, no. 4, pp. 1381–1385, 1999.
- [55] W. Luczaj, A. Gegotek, and E. Skrzydlewska, “Antioxidants and HNE in redox homeostasis,” *Free Radical Biology and Medicine*, vol. 111, pp. 87–101, 2017.
- [56] T. Szasz, K. Thakali, G. D. Fink, and S. W. Watts, “A comparison of arteries and veins in oxidative stress: producers, destroyers, function, and disease,” *Experimental Biology and Medicine*, vol. 232, no. 1, pp. 27–37, 2007.
- [57] I. G. Obrosova, “Increased sorbitol pathway activity generates oxidative stress in tissue sites for diabetic complications,” *Antioxidants & Redox Signaling*, vol. 7, no. 11-12, pp. 1543–1552, 2005.
- [58] A. Petry, T. Djordjevic, M. Weitnauer, T. Kietzmann, J. A. Hess, and A. Görlach, “NOX2 and NOX4 mediate proliferative response in endothelial cells,” *Antioxidants & Redox Signaling*, vol. 8, no. 9-10, pp. 1473–1484, 2006.
- [59] A. K. Jaiswal, “Nrf2 signaling in coordinated activation of antioxidant gene expression,” *Free Radical Biology & Medicine*, vol. 36, no. 10, pp. 1199–1207, 2004.
- [60] M. T. Do, H. G. Kim, J. H. Choi, and H. G. Jeong, “Metformin induces microRNA-34a to downregulate the Sirt1/Pgc-1 α /Nrf2 pathway, leading to increased susceptibility of wild-type p53 cancer cells to oxidative stress and therapeutic agents,” *Free Radical Biology and Medicine*, vol. 74, pp. 21–34, 2014.
- [61] F. Xue, J. W. Huang, P. Y. Ding et al., “Nrf2/antioxidant defense pathway is involved in the neuroprotective effects of Sirt1 against focal cerebral ischemia in rats after hyperbaric oxygen preconditioning,” *Behavioural Brain Research*, vol. 309, pp. 1–8, 2016.
- [62] Y. W. Ding, G. J. Zhao, X. L. Li et al., “SIRT1 exerts protective effects against paraquat-induced injury in mouse type II alveolar epithelial cells by deacetylating NRF2 *in vitro*,” *International Journal of Molecular Medicine*, vol. 37, no. 4, pp. 1049–1058, 2016.