1 2	Supplementary materials for Choline protects heart from doxorubicin-induced cardiotoxicity through vagal activation and Nrf2/HO-1 pathway
3	
4	Authors: Fuding Guo, MD ^a [†] , Yueyi Wang, MD, PhD ^{a,} [†] , Jun Wang, MD ^a [†] , Zhihao
5	Liu, MD ^a , Yanqiu Lai, MD ^a , Zhen Zhou, MD ^a , Zihan Liu, MD ^a , Yuyang Zhou, MD ^a ,
6	Xiao Xu, MD ^a , Zeyan Li, MD ^a , Meng Wang, MD ^a , Fu Yu, MD ^a , Ruijie Hu, MD ^a ,
7	Liping Zhou, MD ^a , Hong Jiang, MD, PhD ^{a, *}
8	
9	Affiliation:
10	^a Department of Cardiology, Renmin Hospital of Wuhan University; Cardiac
11	Autonomic Nervous System Research Centre of Wuhan University; Cardiovascular
12	Research Institute, Wuhan University; Hubei Key Laboratory of Cardiology, Wuhan,
13	China
14	† These authors contributed equally to this work.
15	
16	*Address for correspondence:
17	Hong Jiang, MD, PhD
18	Department of Cardiology, Renmin Hospital of Wuhan University,
19	No. 238 Jiefang Road, Wuhan City, Hubei Provicne 430060, P.R. China,
20	Tel: +86 27 88041911; Fax: +86 27 88040334,
21	E-mail: <u>hong-jiang@whu.edu.cn</u>
22	
23	
24	
25	
26	
27	
28	
29	
30	

- 31 Supplemental Material includes:
- 32 Graphical Abstract
- 33



Heart Injury

Heart Protection

Central illustration of related mechanisms of choline in preventing DOX-induced cardiotoxicity. The protective effect of choline may be associated with the enhancement of vagal activity and the activation of the Nrf2/HO-1 signaling pathway in DOX-treated rats. A simple and non-invasive pharmacological treatment with oral choline may represent a valuable adjunct therapeutic approach for the alleviation of DOX-induced cardiotoxicity in clinic.