

## Research Article

# Differences in Nonspecific Low Back Pain between Young Adult Females with and without Lumbar Scoliosis

Wangshu Yuan <sup>1</sup>, Jianxiong Shen <sup>2</sup>, Lixia Chen <sup>1</sup>, Hai Wang <sup>2</sup>, Keyi Yu,<sup>2</sup> Hui Cong <sup>1</sup>, Jingya Zhou,<sup>3</sup> and Youxi Lin<sup>2</sup>

<sup>1</sup>Department of Rehabilitation and Physical Therapy, Peking Union Medical College Hospital, Beijing, China

<sup>2</sup>Department of Orthopedic Surgery, Peking Union Medical College Hospital, Beijing, China

<sup>3</sup>Department of Medical Records & WHO-FIC Collaborating Center, Peking Union Medical College Hospital, Beijing, China

Correspondence should be addressed to Jianxiong Shen; [sjxpumch@163.com](mailto:sjxpumch@163.com) and Lixia Chen; [clx727@126.com](mailto:clx727@126.com)

Received 14 June 2018; Revised 16 November 2018; Accepted 9 December 2018; Published 3 March 2019

Guest Editor: Baogan Peng

Copyright © 2019 Wangshu Yuan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Study design.** Retrospective characterization of nonspecific low back pain (NSLBP) in young adult female patients with and without lumbar scoliosis. **Background.** There is no consensus as to whether NSLBP in scoliosis patients is related to scoliosis per se or is just a normal symptom that could happen in anyone. **Objectives.** The aim of this study was to compare the differences in NSLBP between young adult female patients with and without lumbar scoliosis and to provide a theoretical basis for differential treatment of NSLBP in patients with and without lumbar scoliosis. **Methods.** Ninety female young adults with NSLBP were divided into scoliosis and nonscoliosis groups. Characteristics of pain, lumbar mobility, muscle strength, Cobb angle, axial trunk rotation (ATR) angle, and surface electromyography (SEMG) signal were compared between the two groups. **Results.** The pain location in scoliotic patients was more concentrated on the left side of the lumbar spine ( $P \leq 0.001$ ). The area affected by pain ( $P = 0.028$ ) and the numerical pain rating scale (NPRS) scores ( $P = 0.014$ ) of scoliotic patients were less than those of nonscoliotic patients. The difference between side-bending in scoliotic patients was greater than that in nonscoliotic patients ( $P = 0.001$ ). Scoliotic patients exhibited a significantly better ability for flexion ( $P = 0.001$ ) and extension ( $P = 0.017$ ) than nonscoliotic patients. The posterior muscles in scoliotic patients were stronger than those in nonscoliotic patients ( $P = 0.014$ ). The ratio of root-mean-square (RMS) on paraspinal muscles in scoliotic patients was greater than that in nonscoliotic patients ( $P \leq 0.001$ ). Scoliotic patients exhibited greater relaxation time during the flexion-relaxation phenomenon (FRP) than nonscoliotic patients ( $P = 0.024$ ). **Conclusions.** The characteristics of NSLBP experienced by patients with lumbar scoliosis were distinct from those of NSLBP experienced by nonscoliotic patients. The treatment of NSLBP in scoliotic patients should be different from that in nonscoliotic patients.

## 1. Introduction

Idiopathic scoliosis is a common spine deformity. A majority of cases of idiopathic scoliosis occur in adolescents, and the disease progression is much faster and greater in adolescents; therefore, adolescent idiopathic scoliosis (AIS) has received more attention than other kinds of idiopathic scoliosis [1]. Owing to the matured skeleton and typically slow rate of disease progression, young adult idiopathic scoliosis (YAIS) has not garnered much attention, either from a clinical or a societal perspective. However, all AIS patients eventually become YAIS patients, and there will be some new cases diagnosed in young adults; therefore, the

number of YAIS patients is much greater than that of AIS patients. Generally, YAIS patients with mild disease experience no functional limitation and their aesthetic appearance is also acceptable; therefore, these patients typically consult a doctor because of low back pain (LBP). Research on the relationship between scoliosis and LBP has largely been focused on adolescent people and most cases of AIS experience nonspecific low back pain (NSLBP). Studies have shown that patients with AIS have a higher prevalence of LBP and experience more severe and longer duration of pain as compared to normal controls [2–4]. However, there is no definitive evidence to suggest that scoliosis is indeed the underlying cause of NSLBP in patients with AIS or whether

the NSLBP is similar to that occurring in nonscoliotic patients [5, 6]. In this retrospective study, we sought to assess the differences between NSLBP experienced by young adult females with and without scoliosis and investigated the associated factors. Our aim was to provide a theoretical basis for a clearer understanding of NSLBP in young adult patients with scoliosis. To the best of our knowledge, this is the first study that compares the NSLBP experienced by scoliotic and nonscoliotic young adults.

## 2. Materials and Methods

**2.1. Subjects.** All patients were enrolled at the Department of Physical Therapy and Rehabilitation between January 2016 and May 2017, and were diagnosed as NSLBP by orthopedic surgeons. NSLBP has been widely described as pain or discomfort that is localised below the costal margin and above the inferior gluteal folds, with or without leg pain, but not attributable to a known or specific pathology [7]. Inclusion criteria were as follows: (1) female patients diagnosed as NSLBP by both orthopedic surgeons and rehabilitation physicians; (2) age range, 18–30 years; (3) duration of pain: 1–6 months; no history of treatment (physical or pharmacological); (4) patients with idiopathic scoliosis only had a lumbar curve (apex located between L2 and L4), and the Cobb angle was  $>10^\circ$ . The exclusion criteria were as follows: (1) LBP caused by any specific reason; (2) history of spine surgery or injury; and (3) patients with intellectual disability and those unable to understand the instructions. A total of 41 patients with scoliosis suffering from NSLBP (mean age:  $24.95 \pm 2.90$  years; mean height:  $162.41 \pm 3.82$  cm; mean body weight:  $55.12 \pm 7.41$  kg) and 49 nonscoliotic patients with NSLBP (mean age:  $24.73 \pm 2.83$  years; mean height:  $162.85 \pm 3.72$  cm; mean body weight:  $55.64 \pm 6.42$  kg) qualified the study selection criteria and were recruited in the study.

There were no significant differences between the two groups with respect to baseline characteristics (Table 1). All patients with scoliosis had undergone radiographic imaging and standard evaluation. The Institution Review Board of the Peking Union Medical Hospital has approved the study and the consent form.

### 2.2. Evaluation and Procedure

**2.2.1. Description of Low Back Pain.** All subjects were asked to answer the following four questions. First question, where is the location of the pain exactly? The response options were as follows: (1) left side of the lumbar spine; (2) right side of the lumbar spine; (3) middle of the lumbar spine; and (4) both sides of the lumbar spine. Second question, how many square centimeters ( $\text{cm}^2$ ) of the painful area? The response options were as follows: (1)  $<5 \text{ cm}^2$ ; (2)  $5\text{--}10 \text{ cm}^2$ ; and (3)  $>10 \text{ cm}^2$ . Third question, how severe is the pain? The subjects were asked to rate the intensity of symptoms on 10-point numerical pain rating scale (NPRS), where 0 indicated no pain and 10 indicated the maximum imaginable pain [8]. Fourth question, does the pain happen suddenly or slowly at the very beginning? The response options were as follows: (1) sudden onset and (2) slow onset.

TABLE 1: Descriptive analysis of the characteristic of subjects in the two groups in this study (SD, standard deviation).

|                                | Scoliosis group   | Nonscoliosis group | <i>P</i> value | <i>t</i> value |
|--------------------------------|-------------------|--------------------|----------------|----------------|
| Age (years)                    | $24.95 \pm 2.90$  | $24.73 \pm 2.83$   | 0.721          | -0.358         |
| Height (standing) (m)          | $162.41 \pm 3.82$ | $162.85 \pm 3.72$  | 0.578          | -0.558         |
| Weight (kg)                    | $55.12 \pm 7.41$  | $55.64 \pm 6.42$   | 0.722          | 0.357          |
| BMI ( $\text{kg}/\text{m}^2$ ) | $20.78 \pm 2.66$  | $21.13 \pm 2.67$   | 0.538          | 0.618          |

**2.2.2. Mobility of the Lumbar Spine.** A physical therapist (PT) evaluated the mobility of the lumbar spine in three planes. (1) In the frontal plane, the subject was made to stand straight against the wall and asked to bend towards the left and right sides to the maximum possible extent. The distance from the middle fingertip to the floor in both standing and the bended position were recorded. The difference between the distance in the standing and side bended position denoted the ability of side-bending. (2) In the sagittal plane, the subject assumed the prone position and pushed her upper body up slowly with her arms, while the PT held her sacrum; the subject was asked to stop moving at the point when the PT felt lifting of her sacrum. The distance of the jugular notch from the bed was recorded to denote the extent of extension. The distance from the middle fingertip to the feet in sit-and-reach was recorded to denote the extent of flexion. (3) In the horizontal plane, the subject held the gymnastics rod on her shoulders and rotated her trunk to both the left and the right side without moving her buttocks in the sitting position. The angle of the rotation of the gymnastics rod was recorded to denote the angle of left and right rotation.

**2.2.3. Strength of the Core Muscles around the Lumbar Spine.** The strength of the anterior and posterior muscles of the lumbar spine was evaluated by a PT. The subject lifted her legs in the supine position, held the thigh at an angle of  $45^\circ$  to the ground, and the calf kept parallel to the ground. The duration of time was recorded to denote the strength of the anterior muscles of the lumbar spine. Subject was asked to undergo a body weight-dependent isometric back extension test (Sorensen Test, ST) whilst lying on a bed [9]. The time duration was recorded to denote the strength of the posterior muscles of the lumbar spine.

**2.2.4. Cobb Angle.** The Cobb angle was measured from the anterior-posterior X-rays by recording the angle between the upper and lower most-tilted end vertebra by a spine specialist orthopedist ( $>20$  years experience in scoliosis operative treatment) and a PT ( $>5$  years experience in scoliosis conservative therapy) [10, 11]. The average Cobb angle is  $26^\circ$  degree and the maximum angle is  $40^\circ$ .

**2.2.5. Axial Trunk Rotation Angle.** The subject was in standing position with 30 cm distance between the feet, and the PT sat behind them. The subjects were asked to bend forward slowly until the point when the deformity of the spine was most prominent and the top of the hump was parallel to the PT's line of sight. The scoliometer was laid

across the apical spinous process and the axial trunk rotation (ATR) angle recorded.

**2.2.6. Surface Electromyography.** The back of the subject was exposed and cleaned with 70% alcohol to reduce skin impedance. For scoliotic patients, the PT palpated the spinous processes, identified the apical spinous process and the intervertebral space between L4 and L5, and marked the skin for placement of electrodes. Six electrodes were placed bilaterally on the marked skin over the erector spinae muscle at the corresponding apical vertebral level and with a 3 cm gap. The other six electrodes were placed bilaterally on the marked skin over the intervertebral space between L4 and L5. For nonscoliotic patients, the electrodes were placed bilaterally on the L2 and L4-L5. Surface electromyography (SEMG) signals were recorded during two movements.

**(1) Extension Force.** The subject assumed the sitting position with his back to the wall (distance between the back and the wall was 10–20 cm) and extended the trunk until the occiput touched the wall. At that point, the subject was asked to push the occiput against the wall with as much force as possible for 30 seconds. Values of root-mean-square (RMS; microvolt) as the SEMG signal were collected. In the upper segment (apical spinous process or L2), the greater value was the numerator, and the smaller value was the denominator. The ratio of RMS value on bilateral lumbar was used to assess the difference:

$$\text{ratio of RMS} = \frac{\text{RMS value (greater value)}}{\text{RMS value (smaller value)}} \quad (1)$$

**(2) FRP [12].** In the standing position, the subject was asked to bend over to the maximum possible extent on hearing the tone and then to maintain relaxation for 10 s. Subsequently, the subject was asked to straighten up on hearing the tone again; the entire process was performed three times. Values of RMS as the SEMG signal were collected during all the processes. In the lower segment (L4-L5), we recorded the number of times the average RMS value was smaller than 5 mV in the stooped position.

**2.3. Statistical Analysis.** All statistical analyses were conducted using SPSS (version 19.0) software (SPSS, Chicago, IL). The data were analyzed using the chi-squared test, analysis of variance, Student's *t*-test, Spearman correlation coefficient, and linear regression.  $P < 0.05$  was considered indicative of statistical significance.

### 3. Results

The distribution of pain in scoliotic patients was largely unilateral. In the scoliosis group, 32 patients had left-sided lumbar pain and 9 patients had right-sided lumbar pain; and 78% of scoliotic subjects had pain on the convex side. In contrast, 83.7% subjects in the nonscoliosis group had midline or symmetrical pain. The painful area in scoliotic patients was smaller than that in the nonscoliotic patients. Nonscoliotic patients had higher NPRS scores than scoliotic patients (Table 2).

TABLE 2: Descriptive statistics of description of the low back pain in two groups.

|               | Scoliosis group | Nonscoliosis group | <i>P</i> value | <i>Z</i> value |
|---------------|-----------------|--------------------|----------------|----------------|
| Pain location | 1 (1, 2.5)      | 3 (3, 4)           | $\leq 0.001^*$ | -4.316         |
| Painful area  | 2 (2, 2)        | 2 (2, 3)           | 0.028*         | -2.400         |
| NPRS          | 3.5 (2.5, 6.0)  | 5.5 (3.5, 7.5)     | 0.014*         | -2.443         |
| Pain onset    | 2 (1, 2)        | 1 (1, 2)           | 0.181          | -1.604         |

\*Difference between two groups is significant,  $P < 0.05$ .

The extent of side-bending in the scoliosis group was significantly greater than that in the nonscoliosis group. The extent of flexion and extension in the scoliosis group were significantly greater than that in the nonscoliosis group. Other movements of the lumbar spine were comparable between the two groups (Table 3).

The posterior muscles in scoliotic patients were significantly stronger than those in nonscoliotic patients. No significant between-group difference was observed with respect to the strength of anterior muscles (Table 4).

Scoliotic patients showed significantly greater Cobb angle and ATR angle as compared to nonscoliotic patients (Table 5).

The ratio of RMS value on bilateral lumbar in scoliotic patients was significantly greater than that in nonscoliotic patients. Scoliotic patients showed significantly greater relaxation times in TFRP than in nonscoliotic patients (Table 6).

### 4. Discussion

NSLBP and idiopathic scoliosis are two diseases with unclear etiology; currently, the two conditions can only be distinguished based on symptoms and clinical evaluation. Contrary to the findings reported by Gremeaux et al. and Jackson et al. [13, 14], patients with scoliosis showed lesser severity of pain in the present study. The study by Gremeaux et al. found no significant difference between scoliotic and nonscoliotic patients with regard to pain severity. In the study by Jackson et al., pain severity was greater in scoliotic patients. In the present study, most patients in the scoliosis group had unilateral pain especially on the convex side of the lumbar curve, which is different from the results reported by Joncas et al. [15]. These differences could be attributable to different age profile of the study population. The average age of subjects in these studies was  $>50$  years, while the average age of subjects in the present study was 24 years. Younger subjects are less affected by age-related degenerative changes. It is likely that the pain was more closely associated with the scoliosis at the very beginning; with increase in age, the patients gradually experienced pain from NSLBP. With further progression of age, scoliotic patients experience pain due to both scoliosis and NSLBP; therefore, the pain severity was even greater than that in the control group. The location of pain on the convex side is not just because of the uneven biomechanics caused by vertebral deviation and rotation but may also be related to psychological factors [16, 17]. Most

TABLE 3: Mobility of the lumbar spine in 3 planes in two groups.

|                                      | Scoliosis group | Nonscoliosis group | <i>P</i> value | <i>t/Z</i> value |
|--------------------------------------|-----------------|--------------------|----------------|------------------|
| Left side-bending (cm)               | 17.73 ± 3.91    | 17.47 ± 3.67       | 0.744          | -0.328           |
| Right side-bending (cm)              | 19.41 ± 3.85    | 18.55 ± 3.56       | 0.273          | -1.013           |
| Difference between side-bending (cm) | 3 (1, 5)        | 1 (1, 2)           | 0.001*         | -3.329           |
| Left rotation (°)                    | 57.20 ± 16.43   | 55.00 ± 18.54      | 0.558          | -0.589           |
| Right rotation (°)                   | 61.95 ± 14.31   | 57.22 ± 17.82      | 0.175          | -1.368           |
| Difference between rotation (°)      | 5 (0, 10)       | 5 (0, 5)           | 0.103          | -1.629           |
| Flexion (cm)                         | 5 (0, 15)       | 0 (-12.5, 6)       | 0.001*         | -3.462           |
| Extension (cm)                       | 18.22 ± 3.34    | 16.39 ± 3.76       | 0.017*         | -2.423           |

\*Difference between two groups is significant,  $P < 0.05$ .

TABLE 4: Descriptive statistics of strength of the core muscles around the lumbar spine in both groups.

|                     | Scoliosis group | Nonscoliosis group | <i>P</i> value | <i>Z</i> value |
|---------------------|-----------------|--------------------|----------------|----------------|
| Anterior muscle(s)  | 60 (41, 69)     | 41 (36, 70)        | 0.340          | -0.955         |
| Posterior muscle(s) | 74 (48, 154)    | 64 (35, 75)        | 0.014*         | -2.469         |

\*Difference between two groups is significant,  $P < 0.05$ .

TABLE 5: Descriptive statistics of Cobb angle and ATR angle in two groups.

|                | Scoliosis group | Nonscoliosis group | <i>P</i> value | <i>Z</i> value |
|----------------|-----------------|--------------------|----------------|----------------|
| Cobb angle (°) | 26 (20, 30)     | 1 (0, 5)           | ≤0.001*        | -5.508         |
| ATR angle (°)  | 8 (7, 9)        | 2 (0, 2)           | ≤0.001*        | -5.555         |

\*Difference between two groups is significant,  $P < 0.05$ .

TABLE 6: SEMG activities in two groups.

|               | Scoliosis group | Nonscoliosis group | <i>P</i> value | <i>t/Z</i> value |
|---------------|-----------------|--------------------|----------------|------------------|
| Ratio of RMS  | 1.59 ± 0.44     | 1.24 ± 0.40        | ≤0.001*        | 3.956            |
| Times of FERP | 3 (1.5, 3)      | 2 (0, 3)           | 0.024*         | -2.225           |

\*Difference between two groups is significant,  $P < 0.05$ .

female young adults with scoliosis attach importance to the aesthetic aspects; therefore, the presence of a hump on the lower back is likely to induce regional pain due to psychological reasons.

In the present study, scoliotic patients exhibited uneven side-bending which is different from the results reported by Veldhuizen and Scholten [18]. From a biomechanical perspective, patients with a left lumbar curve experience restriction of bending towards the left side. However, the left lumbar curve is likely to induce trunk deviation to the left side, which facilitates bending towards the left side. This represents a natural compensatory mechanism. Therefore, in the study by Veldhuizen et al., there was no difference between the extent of bending on the two sides. However, the presence of pain, especially unilateral pain, serves to further exaggerate the uneven side-bending. Studies by Hultman et al. [19] and Taechasubamorn et al. [20] showed that patients with low back pain have significantly lesser sagittal flexibility and endurance of back extensors than healthy people. In this study, patients in the scoliosis group showed significantly greater sagittal flexibility and endurance of back extensor muscles as compared to patients in the

nonscoliosis group. This suggests that the NSLBP in nonscoliotic patients caused significantly greater disability than in scoliotic patients. In other words, scoliotic patients showed a less stronger relationship with the restricted flexibility and weak muscle strength.

RMS is the time threshold indicator of SEMG; it was shown to reflect the functional capacity of the muscles and was linked to neuromuscular efficiency [21]. In the present study, the RMS value on the convex side of the apical vertebral region was greater than that on the concave side, which is consistent with the results of previous studies [22–25]. During isometric contraction of homologous muscles under the same load, atrophic muscles were shown to produce greater RMS values than fatigued muscles, and fatigued muscles produced greater RMS values than trained muscles [26]. In patients with scoliosis, the convex side muscles generated greater RMS value than the concave side; therefore, we predict that the convex side muscles were weaker than those on the concave side and that the uneven functional capacity of paraspinal muscles was one of the reasons why a significant proportion of patients in the scoliosis group had unilateral pain. FRP was commonly observed in healthy subjects. In contrast, few patients with low back pain demonstrated FRP [27]. In this study, FRP in nonscoliotic patients with NSLBP was less commonly observed than that in scoliotic patients with NSLBP. These findings imply that the FRP in scoliotic patients with NSLBP was more akin to that in healthy subjects.

In conclusion, irrespective of the pain description or pain-related parameters, NSLBP in young adult females with scoliosis is not the same as that in nonscoliotic patients. These findings suggest that the treatment of NSLBP in scoliotic patients should be different from that in nonscoliotic patients.

## Data Availability

The data used to support the findings of this study are restricted by the Institution Review Board of Peking Union Medical College Hospital in order to protect patient privacy.

Data are available from Yuan Wangshu (ywsartemis@126.com) for researchers who meet the criteria for access to confidential data.

## Conflicts of Interest

The authors declare that there are no conflicts of interest.

## References

- [1] H. K. Wong and K. J. Tan, "The natural history of adolescent idiopathic scoliosis," *Indian Journal of Orthopaedics*, vol. 44, no. 1, pp. 9–13, 2010.
- [2] T. Sato, T. Hirano, T. Ito et al., "Back pain in adolescents with idiopathic scoliosis: epidemiological study for 43,630 pupils in Niigata City, Japan," *European Spine Journal*, vol. 20, no. 2, pp. 274–279, 2011.
- [3] Z. Landman, T. Oswald, J. Sanders et al., "Prevalence and predictors of pain in surgical treatment of adolescent idiopathic scoliosis," *Spine*, vol. 36, no. 10, pp. 825–829, 2011.
- [4] Y. Smorgick, Y. Mirovsky, K. C. Baker et al., "Predictors of back pain in adolescent idiopathic scoliosis surgical candidates," *Journal of Pediatric Orthopaedics*, vol. 33, no. 3, pp. 289–292, 2013.
- [5] F. M. Kovacs, M. Gestoso, M. T. Gil del Real, J. López, N. Mufraggi, and J. I. Méndez, "Risk factors for non-specific low back pain in school children and their parents: a population based study," *Pain*, vol. 103, no. 3, pp. 259–268, 2003.
- [6] F. Balagué and F. Pellisé, "Adolescent idiopathic scoliosis and back pain," *Scoliosis Spinal Disord*, vol. 11, no. 1, p. 27, 2016.
- [7] A. K. Burton, F. Belague, G. Cardon et al., "Chapter 2 European guidelines for prevention in low back pain November 2004," *European Spine Journal*, vol. 15, no. 2, pp. 136–168, 2006.
- [8] J. D. Childs, S. R. Piva, and J. M. Fritz, "Responsiveness of the numeric pain rating scale in patients with low back pain," *Spine*, vol. 30, no. 11, pp. 1331–1334, 2005.
- [9] A. Champagne, M. Descarreaux, and D. Lafond, "Back and hip extensor muscles fatigue in healthy subjects: task-dependency effect of two variants of the sorensen test," *European Spine Journal*, vol. 17, no. 12, pp. 1721–1726, 2008.
- [10] J. Cheung, D. J. Wever, A. G. Veldhuizen et al., "The reliability of quantitative analysis on digital images of the scoliotic spine," *European Spine Journal*, vol. 11, no. 6, pp. 535–542, 2002.
- [11] J. R. Cobb, "Outline for the study of scoliosis. Instructional course lectures," *American Academy of Orthopaedic Surgeons*, vol. 5, pp. 261–275, 1948.
- [12] R. Neblett, "Surface electromyographic (SEMG) biofeedback for chronic low back pain," *Healthcare*, vol. 4, no. 2, p. 27, 2016.
- [13] V. Gremaux, J. Casillas, P. Fabbro-Peray et al., "Analysis of low back pain in adults with scoliosis," *Spine*, vol. 33, no. 4, pp. 402–405, 2008.
- [14] R. P. Jackson, E. H. Simmons, and D. Stripinis, "Incidence and severity of back pain in adult idiopathic scoliosis," *Spine*, vol. 8, no. 7, pp. 749–756, 1983.
- [15] J. Joncas, H. Labelle, and B. Poitras, "Dorso-lumbal pain and idiopathic scoliosis in adolescence," *Annales de Chirurgie*, vol. 50, no. 8, pp. 637–640, 1996.
- [16] L. Saccomani, F. Vercellino, and P. Rizzo, "Adolescents with scoliosis: psychological and psychopathological aspects," *Minerva Pediatrica*, vol. 50, no. 1-2, pp. 9–14, 1998.
- [17] T. N. Theologis, R. J. Jefferson, and A. H. Simpson, "Quantifying the cosmetic defect of adolescent idiopathic scoliosis," *Spine*, vol. 18, no. 7, pp. 909–912, 1993.
- [18] A. G. Veldhuizen and P. J. Scholten, "Flexibility in structurally normal young females and in young females with idiopathic scoliosis," *Clinical Biomechanics*, vol. 5, no. 2, pp. 117–119, 1990.
- [19] G. Hultman, H. Saraste, and H. Ohlsen, "Anthropometry, spinal canal width, and flexibility of the spine and hamstring muscles in 45-55-year-old men with and without low back pain," *Journal of Spinal Disorders*, vol. 5, no. 3, pp. 245–253, 1992.
- [20] P. Taechasubamorn, T. Nopkesorn, and S. Pannarunothai, "Comparison of physical fitness between rice farmers with and without chronic low back pain: a cross-sectional study," *Journal of the Medical Association of Thailand*, vol. 93, no. 12, pp. 1415–1421, 2010.
- [21] H. A. DeVries, "Efficiency of electrical activity as a physiological measure of the functional state of muscle tissue," *American Journal of Physical Medicine & Rehabilitation*, vol. 47, no. 1, pp. 10–22, 1968.
- [22] G. Kwok, J. Yip, M. C. Cheung et al., "Evaluation of myoelectric activity of paraspinal muscles in adolescents with idiopathic scoliosis during habitual standing and sitting," *BioMed Research International*, vol. 2015, Article ID 958450, 9 pages, 2015.
- [23] C. Zetterberg, R. Björk, R. Ortengren, and G. B. Andersson, "Electromyography of the paravertebral muscles in idiopathic scoliosis. Measurements of amplitude and spectral changes under load," *Acta Orthopaedica Scandinavica*, vol. 55, no. 3, pp. 304–309, 1984.
- [24] M. A. Alexander and E. H. Season, "Idiopathic scoliosis: an electromyographic study," *Archives of Physical Medicine and Rehabilitation*, vol. 59, no. 7, pp. 314–315, 1978.
- [25] M. Reuber, A. Schultz, T. McNeill et al., "Trunk muscle myoelectric activities in idiopathic scoliosis," *Spine*, vol. 8, no. 5, pp. 447–456, 1983.
- [26] H. Christenstensen, M. Lo Monaco, K. Dahl, and A. Fuglsang-Frederiksen, "Progressing of electrical activity in human muscle during a gradual increase in force," *Electroencephalography and Clinical Neurophysiology*, vol. 58, no. 3, pp. 230–239, 1984.
- [27] O. Shirado, T. Ito, K. Kaneda et al., "Flexion-relaxation phenomenon in the back muscles. A comparative study between healthy subjects and patients with chronic low back pain," *American Journal of Physical Medicine & Rehabilitation*, vol. 74, no. 2, pp. 139–144, 1995.



**Hindawi**

Submit your manuscripts at  
[www.hindawi.com](http://www.hindawi.com)

