

## Review Article

# Accurate Diagnosis and Treatment of Painful Temporomandibular Disorders: A Literature Review Supplemented by Own Clinical Experience

Adam Andrzej Garstka <sup>1</sup>, Lidia Kozowska <sup>2</sup>, Konrad Kijak <sup>3</sup>, Monika Brzózka <sup>1</sup>,  
Helena Gronwald <sup>1</sup>, Piotr Skomro <sup>1</sup> and Danuta Lietz-Kijak <sup>1</sup>

<sup>1</sup>Department of Propaedeutic, Physical Diagnostics and Dental Physiotherapy, Faculty of Medicine and Dentistry, Pomeranian Medical University, Szczecin, Poland

<sup>2</sup>Student Scientific Society, Pomeranian Medical University, Szczecin, Poland

<sup>3</sup>Student Scientific Society, Medical University of Silesia, Zabrze, Poland

Correspondence should be addressed to Danuta Lietz-Kijak; danuta.lietzkijak@gmail.com

Received 21 March 2022; Revised 4 September 2022; Accepted 21 January 2023; Published 31 January 2023

Academic Editor: Tomasz Kaczmarzyk

Copyright © 2023 Adam Andrzej Garstka et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Introduction.** Temporomandibular disorders (TMD) is a multifactorial group of musculoskeletal disorders often with combined etiologies that demand different treatment plans. While pain is the most common reason why patients decide to seek help, TMD is not always painful. Pain is often described by patients as a headache, prompting patients to seek the help of neurologists, surgeons, and ultimately dentists. Due to the unique characteristics of this anatomical area, appropriate diagnostic tools are needed, as well as therapeutic regimens to alleviate and/or eliminate the pain experienced by patients. *Aim of the Study.* The aim of this study is to collect and organize information on the diagnosis and treatment of pain in TMD, through a review of the literature supplemented by our own clinical experience. *Material and Methods.* The study was conducted by searching scientific databases PubMed, Scopus, and Google Scholar for documents published from 2002–2022. The following keywords were used to build the full list of references: TMD, pain, temporomandibular joint (TMJ), TMJ disorders, occlusal splint, relaxing splints, physiotherapy TMD, pharmacology TMD, natural therapy TMD, diagnostic criteria for TMD, and DC/TMD. The literature review included 168 selected manuscripts, the content of which was important for pain diagnosis and clinical treatment of TMD. *Results.* An accurate diagnosis of TMD is the foundation of appropriate treatment. The most commonly described treatments include physiotherapy, occlusal splints therapy, and pharmacological treatment tailored to the type of TMD. *Conclusions.* Based on the literature review and their own experience, the authors concluded that there is no single ideal form of pain therapy for TMD. Treatment of TMD should be based on a thorough diagnostic process, including the DC/TMD examination protocol, psychological evaluation, and cone beam computer tomography (CBCT) imaging. Following the diagnostic process, once a diagnosis is established, a treatment plan can be constructed to address the patient's complaints.

## 1. Introduction

Temporomandibular disorders (TMD) can present with pain, prompting patients to seek help from various specialists [1–4]. TMD is most frequently seen in people aged between 20 and 40 years [5–8] and is more common in women due to hormonal changes and greater influence of psychosocial factors [9–12]. Thus, it can be concluded that

TMD is a civilization problem, which may escalate due to the increasing pace of life, omnipresent stress, and improper use of the masticatory system [13–24]. One unquestionable causative factor is stress, which has a destructive effect on all masticatory structures, and if chronic, it may expose or aggravate temporomandibular disorders [25–32].

Pain in temporomandibular disorders may have a diverse etiology, i.e., central or peripheral, as demonstrated by

the 2020 study by Yin et al., finding that TMD is accompanied by functional and structural changes in the primary somatosensory cortex, prefrontal cortex, and basal ganglia of the brain, which should inform treatment decisions [33]. Temporomandibular disorders (TMD) are characterized by abnormalities in the temporomandibular joint, masticatory muscles, and other adjacent structures, often described by patients as a headache [34–40]. According to research findings, typical TMD symptoms are more common in patients with migraine or tension headaches. It has also been shown that patients with diagnosed TMD are more likely to experience migraines, and the coexistence of both problems exacerbates the symptoms of each [41–47]. This unique anatomical region does not lend itself easily to diagnosis and treatment. It is not uncommon for patients to be referred to neurologists, otolaryngologists, surgeons, and dentists. Undoubtedly, the involvement of many specialists in the problems affecting this area may be beneficial in the classification and differentiation of disorders [48–51].

Masticatory dysfunction can be diagnosed when at least three of the following symptoms are reported: pain and acoustic symptoms during mandibular movements, limited mandibular mobility, difficulty with jaw opening, and occlusal or nonocclusal parafunction. The modern diagnosis of TMD should be based on the DC/TMD examination protocol because only with the correct diagnosis is the correct treatment possible [52–54].

## 2. Aim of the Study

The aim of this study is to collect and organize information on the accurate diagnosis and treatment of pain in TMD through a review of the literature supplemented by our own clinical experience.

## 3. Materials and Methods

The study was conducted by searching scientific databases PubMed, Scopus, and Google Scholar for documents published from 2002–2022. The literature review included 168 selected manuscripts, the content of which was important for pain diagnosis and clinical treatment of TMD. These aspects mentioned previously were the criteria for the inclusion of the manuscripts in the review. The following keywords were used to build the full list of references: TMD, pain, TMJ disorders, occlusal splints, relaxing splints, physiotherapy TMD, pharmacology TMD, and natural therapy TMD.

## 4. The Essence of the Matter

### 4.1. TMD Pain Diagnosis

**4.1.1. Myalgia.** Myalgia (muscle pain) can be caused by mandibular movements, parafunctions, and excessive muscle tension due to the increased activity of masticatory muscles. Pain occurs upon provocation testing. The patient's history may include pain in the jaw, temple, ear, or in front of the ear. Pain may be modified with jaw movement, function, or parafunction.

Upon physical examination of the patient, the physician is able to confirm the location of pain in the temporalis or masseter muscle, additionally using muscle palpation and maximum unassisted or assisted jaw opening [55–75].

**4.1.2. Myofascial Pain.** Myofascial pain can be local or referred to and is experienced by the patient as deep and dull. Unlike myalgia, this pain spreads beyond the palpated area, remaining inside the boundary of the examined muscle or in the case of referred myofascial pain—beyond the area of the examined muscle. Myofascial trigger points may also be felt during palpation [76–80].

**4.1.3. Arthralgia.** The term arthralgia refers to pain in the temporomandibular joint without signs of joint inflammation. The onset of pain is associated with mandibular movement, function, and parafunction. Pain is also triggered during provocation testing. The patient's history includes pain in the jaw, temple, ear, or in front of the ear. On physical examination, the physician confirms pain in the TMJ area, especially the lateral region, and examines the maximum range of jaw opening with and without assistance [81–83].

**4.1.4. TMD-Attributed Headache.** Headache attributed to temporomandibular dysfunction is characterized by a history of temporal pain of any nature. The pain can be modified by mandibular movement, function, and parafunction. Upon physical examination, pain in the temporalis region can also be observed in provocative tests. Pain may occur during palpation and when testing jaw opening [84].

**4.1.5. Disc Displacement with Reduction or with Intermittent Locking.** An intracapsular disorder involving the condyle-disc complex. To make the diagnosis, it is necessary to determine the closed mouth position according to the protocol. At the Department of Propaedeutics, Physical Diagnostics, and Dental Physiotherapy, we ask the patient to assume their habitual occlusion and then relax the mandible. In this way, we are able to assess the actual intraarticular status, which is confirmed by palpation, joint sound inspection (with a stethoscope), and diagnostic imaging (CBCT). When performing diagnostic imaging, it is essential to perform the examination under the same conditions, without the bite stick.

In this disorder, the disc is positioned anteriorly relative to the condylar head and reduces with mouth opening movements. In some cases, medial and lateral displacement of the articular disc can be observed, as well as noises such as clicking, crackling, or popping [85–93]. Please note that if the patient has a history of joint locking and chewing problems, this diagnosis is ruled out.

To make the diagnosis, the patient is asked to report all TMJ noises that have occurred in the last 30 days during mandibular movements, and additionally, the patient should report any noises during the examination:

- (i) Clicking, popping, and/or snapping noise during both opening and closing movements, detected with palpation during at least one of three repetitions of jaw opening and closing movements; or
- (ii) Clicking, popping, and/or snapping noise detected with palpation during at least one of three repetitions of opening or closing movement(s);
- (iii) Clicking, popping, and/or snapping noise detected with palpation during at least one of three repetitions of right or left lateral, or protrusive movement(s) [94–100].

When discussing this disorder, it should be stated that imaging should be the reference standard for this diagnosis [101–103].

**4.1.6. Disc Displacement without Reduction with Limited Opening.** In this intracapsular disorder, in the closed mouth position, the disc is positioned anteriorly relative to the condylar head and does not reduce in size with the opening of the mouth. Characteristically, the disorder is associated with persistent limited mandibular opening, sometimes referred to as a closed lock, which is not resolved by a manipulative manoeuvre performed by the physician.

Patient history includes a locked jaw, limited movement, and eating difficulties. In physical examination, during assisted jaw opening, the distance between the upper and lower incisors is less than 40 mm. Passive movements may be accompanied by noise [104–107].

**4.1.7. Osteoarthritis of the Temporomandibular Joint.** This disorder involves joint tissue deterioration with concomitant osseous changes in the condylar head and/or articular eminence.

In history, the patient reports noise when chewing or opening the mouth in the last 30 days, and these phenomena may also appear during the examination. On physical examination, the physician detects snapping, popping sounds in the joint during the abduction, adduction, and lateral or protrusive movements. Imaging is required, as CBCT may help visualize subchondral cysts, erosions, generalized sclerosis/calcification, or osteophytes [108–111].

**4.1.8. Subluxation.** A hypermobility disorder involving the disc-condyle complex and the articular eminence. In the open mouth position, the disc is anterior to the articular eminence and the normally closed mouth position cannot be restored without a manipulative manoeuvre. The difference between subluxation and luxation is that in the former the patient is able to reduce the dislocation on their own, whereas the latter requires professional intervention. Patient history includes jaw locking upon abduction movement in the last 30 days. These locks may have been incidental and temporary, resulting in an inability to close the mouth [112, 113].

The RDC/TMD and DC/TMD protocols make it possible to establish a diagnosis but do not shed any light on the etiology of the disorder, and elimination of the cause or an attempt to create the optimal conditions will be crucial in the treatment process.

At the Department of Propaedeutics, Physical diagnostics, and Dental Physiotherapy, the treatment team consists of an orthodontist, a physician dealing with dental prosthetics and restorative dentistry, a physiotherapist, and a dentist who coordinates the work of the whole team [114]. One of the most common signs of a disease process within the TMJ are sounds emitted by the articular structures, such as popping, clicking, humming, grinding, or crunching [114].

Egermark et al., after examining 320 children aged 7, 11, and 15 years, reported that acoustic symptoms were more common in those with malocclusion (24%), with a predominance of transverse malocclusion. In their conclusions, they noted that there were no significant differences in the prevalence of masticatory dysfunction in the studied population between patients with malocclusion and those with a normal bite [115].

Research findings provide no clear-cut conclusion as to how temporomandibular joint disorders are affected by a malocclusion. The consequences of malocclusion in terms of TMD development may be manifold and are undoubtedly related to age, gender, as well as the severity of the disorder.

A fairly significant problem reported and observed in patients is nocturnal bruxism, which affects 8% of the population, and awake bruxism, the prevalence of which is estimated at 20%. At present, bruxism is defined not as a disorder but as a physiological stress-coping mechanism [116–121].

Based on our own experience, we would like to note the relatively frequent coexistence of TMD with orthodontic disorders and temporomandibular disorders in post-orthodontic patients, where the teeth were often aligned in arches while the condylar heads were displaced posteriorly with reduced joint space [122]. In addition, it is important to consider that dental arches are somatic sites where excessive emotional tension can be diffused and reduced [123].

Research into the associations between malocclusion and TMD, as well as the influence of malocclusion treatments on TMD should be conducted in large study samples.

## 4.2. TMD Pain Therapy

**4.2.1. Natural Methods.** Acupuncture is the best-known method of traditional Chinese medicine that is often used, also in Poland, in the treatment of chronic pain. Acupuncture points often coincide with so-called trigger points and correspond to sites of increased density of A- $\delta$  and C fibre nerve endings that conduct pain sensations. Warm compress therapy is used for chronic inflammation and muscle strains. Ideally, a warm compress at 35–40 degrees C should be applied for 20–30 minutes. Cold compresses, on the other hand, are good for acute inflammation with pain and swelling [124, 125].

*4.2.2. Psychological and Behavioural Methods.* Psychological and behavioural programmes are effective in alleviating the psychological crisis, allowing the patient to change their perception of pain and improving functioning in patients with chronic pain. The therapeutic effect is not affected by the duration of the programme or by whether the treatment is delivered in an individual or group setting.

Behavioural approaches aim to reduce the frequency of pain-promoting behaviours and increase the frequency of health-promoting behaviours. They include:

- (i) improving physical fitness
- (ii) social and employment activation
- (iii) reducing the amount of medication
- (iv) reducing overuse of health services

Psychological methods include the following:

- (i) modifying ways of thinking about pain (misconceptions about pain) that cause prolonged suffering and disability
- (ii) replacing a sense of helplessness with a sense of control over pain and one's own life
- (iii) developing strategies for adequate and effective pain management
- (iv) returning to work and promoting an active lifestyle [126, 127]

It must be remembered that effective pain control requires a multidimensional approach, aiming to reduce the pain but also to improve the patient's quality of life.

*4.2.3. Interventional Methods-Splint Therapy.* Occlusal splint therapy can be used in all TMD disorders; however, it is vitally important to use the right splint for the patient's unique situation.

An occlusal splint is an appliance that affects the mutual relationship of the upper and lower teeth and, consequently, the relationship of the condylar process to the mandibular fossa and articular eminence within the TMJ. The purpose of splints is to stabilize occlusion or to protect teeth from excessive abrasion [128, 129].

According to numerous studies, the use of splints has a significant effect on alleviating or even eliminating the patient's pain symptoms. In cases of disc displacement, repositioning splints are used to stabilize the mandible in the centric relation, and in cases of masticatory muscle disorders, relaxation splints are used to prevent parafunctional effects [130, 131].

Splints are most commonly made by obtaining dental impressions and making a bite registration with wax or silicone mass. An intraoral scanner and electronic bite registration can also be used.

The technique recommended by our team for making occlusal splints is 3D printing using special resin, which makes it possible to avoid the mistakes common in the conventional hand-made process. On the basis of our own experience, research findings, and patient feedback, we use two types of splints in the Department of Propaedeutics,

Physical diagnostics, and Dental Physiotherapy: the Michigan-type relaxation splint and the maxillomandibular repositioning splint [132, 133].

The Michigan-type relaxation splint with canine guidance is used in cases involving: myalgia, myofascial pain, and TMD-attributed headache.

The relaxation splint is made from hard resin and always applied to a single arch, with the upper usually being the arch of choice—unless there are missing teeth in the back. Importantly, in the case of missing teeth, the design of the splint should allow for retention elements.

The hard repositioning splint joined interocclusal in the correct construction bite relationship is used in the following situations: arthralgia, disc displacement with reduction, disc displacement with reduction with intermittent locking, disc displacement without reduction with a limited opening, disc displacement without reduction and without limited opening, osteoarthritis of the temporomandibular joint, subluxation.

*4.2.4. Physiotherapy.* Physiotherapy is a discipline of health science that aims to eliminate, alleviate, and prevent various ailments, as well as restore functional ability through movement and various physical agents. Physiotherapists are part of the treatment process in the case of dysfunctions involving the neuromuscular, musculoskeletal, and other systems [134].

In their work, physiotherapists use kinesiotherapy and physical therapy techniques.

- (i) Self-therapy and muscle training. The patient is taught how to perform the correct opening, closing, lateral and protrusive movements of the mandible, as well as how to deal with sudden pain. Exercises should be performed daily in front of a mirror, and if the treatment includes a splint, it should also be used during exercises. The purpose of the exercises is to shorten the overstretched muscles and relax them, which may help improve symmetry and regulate muscle tone [135].
- (ii) Manual therapy makes use of trigger points. For disc displacement, a joint mobilization technique is applied, which involves the physiotherapist performing traction and gliding movements with low velocity but increasing amplitude. These movements are performed parallel and perpendicular to the joint surface. If the mandibular range of motion is limited, muscle energy techniques (MET) can be used. Treatments using the MET involve the repetition of three steps: in step one, the muscle is stretched to the point of resistance of the tissues; in step two, the patient slightly contracts muscles for about 10 seconds trying to resist the force generated by the physiotherapist; in the last step, the patient relaxes the muscles [136].
- (iii) Massage is used for myofascial pain in order to achieve pain relief and improve muscle length and flexibility, as well as loosen fascia [137, 138]. The

frequency of massage sessions should be 30 minutes twice a week. With subsequent visits, the treatment should be applied with increasing force.

- (iv) Physical therapy, such as ultrasound and transcutaneous electrical nerve stimulation (TENS) can be used for pain of muscular origin. Therapeutic ultrasound can be applied in three modalities: using continuous waves, short bursts (pulsed ultrasound), and ultrasound combined with electrical stimulation, the latter of which has proven to be the most effective.

TENS relieves pain and relaxes masticatory muscles in symptomatic patients with TMD [139–142]. In the pain of intracapsular origin, positive results have been observed after the application of a magnetic field combined with LED light therapy. The Solux infrared lamp can be used in cases of arthropathy and rheumatic diseases. The beneficial effects of heat therapy include the alleviation of pain.

- (i) The Kinesio Taping method is used for TMJ stabilization. It should be applied bilaterally. The tapes work by reducing the tension in the masticatory muscles, as well as the adjacent structures such as the muscles of the neck, shoulders, and spine [143–146]. The application of tapes also stimulates lymphatic drainage, which has a beneficial effect on inflammation accompanied by tissue swelling.
- (ii) Iontophoresis is the use of direct electrical current to accelerate the transdermal delivery of nonsteroidal anti-inflammatory drugs (NSAIDs), corticosteroids, and analgesics. While it is not associated with pain relief, a significant improvement in the range of motion in the joint has been observed [147].

**4.2.5. Pharmacotherapy.** The decision on the use of medications in temporomandibular disorders should be preceded by a thorough analysis of the risks and benefits of the drug [148–152]. Medications used to treat TMD include analgesics, nonsteroidal anti-inflammatory drugs, anticonvulsants, muscle relaxants, and benzodiazepines [153, 154].

**4.2.6. Nonsteroidal Anti-Inflammatory Drugs (NSAID).** NSAIDs are beneficial for patients with acute temporomandibular arthritis resulting from sudden disc displacement. Treatment should continue for a minimum of two weeks, and it is important to combine NSAIDs with gastroprotective agents. Among NSAIDs, ibuprofen appears to be the safest for the gastrointestinal tract [155].

It should also be noted that taking NSAIDs for more than 5 days may reduce the efficacy of antihypertensive drugs, such as diuretics, beta-blockers, and ACE inhibitors [154, 155]. In addition, NSAIDs used with anticoagulants such as warfarin or acenocoumarol may increase the risk of bleeding.

**4.2.7. Opioids.** Due to the interactions of NSAIDs with anticoagulants, as well as the risk of gastritis, physicians

sometimes choose to administer oral opioids, such as codeine and oxycodone. The intraarticular delivery route has been studied, but the findings are conflicting [156]. It is essential to bear in mind the side effects of opioid use, which include: dizziness, excessive sedation, nausea, vomiting, constipation, physical dependence and addiction, and respiratory depression. Because of the mentioned reasons, the use of opioids for the management of TMD should be discouraged [157–159].

**4.2.8. Corticosteroids.** Corticosteroids are helpful in the treatment of moderate to severe TMD. They can be administered by intraarticular injection or by oral route. They have an anti-inflammatory effect which can help relieve pain.

For intraarticular injections, it is a good idea to combine corticosteroid preparation with a local anaesthetic, such as lidocaine. According to research findings, this approach provides for a significant reduction in pain, lasting 4 to 6 weeks, and a reduced risk of complications.

Corticosteroids should be used with caution or discontinued in patients with hypertension, adrenal disease, or electrolyte problems. On day 4 after injection, it is recommended to introduce NSAIDs [160–163].

**4.2.9. Myorelaxants.** Muscle relaxants are used to reduce skeletal muscle tone and, therefore, may be helpful in the management of TMD of muscular origin and chronic orofacial pain [164]. The most common myorelaxants include cyclobenzaprine, metaxalone, methocarbamol, and carisoprodol. Based on numerous studies, cyclobenzaprine is considered to be the drug of choice due to relieving the pain of muscular origin and improving sleep quality [165].

Caution should be exercised when using this type of medication due to its potential to induce significant sedation. These drugs are contraindicated in patients with hyperthyroidism, heart failure, after myocardial infarction, and heart rhythm disorders. The recommended dose is 10 mg at bedtime for 30 days, followed by a 2-week period to flush the drug out of the system and a medical follow-up. In the course of the therapy, the patient should always remain under medical supervision.

**4.2.10. Anticonvulsants.** When discussing anticonvulsants, it is worth noting gabapentin, a GABA analogue. Gabapentin is thought to inhibit neurotransmitter release and reduce postsynaptic excitability [166].

The use of gabapentin reduces the pain of muscular origin, particularly from the temporal and masseter muscles. The drug is generally well tolerated and is associated with transient and mild side effects, including dizziness, drowsiness, dry mouth, weight gain, and impaired concentration [167].

**4.2.11. Benzodiazepines.** Benzodiazepines facilitate transmission in the GABAergic system. They have been found to produce anxiolytic, sedative, hypnotic, anticonvulsant, and

myorelaxant effects. Due to the risk of tolerance and dependence, as well as side effects including confusion, amnesia, and impaired motor coordination, these drugs are not recommended for the treatment of TMD [168].

## 5. Summary

Based on the literature review, the authors concluded that there is no single, ideal form of pain therapy for TMD. Treatment of TMD should be based on a thorough diagnostic process, including the DC/TMD examination protocol, psychological evaluation, and CBCT imaging. Following the diagnostic process, once a diagnosis is established, a treatment plan can be constructed to address the patient's complaints.

The treatment of temporomandibular dysfunctions requires a thorough diagnostic process, taking into account the etiology of the disorder. Having reviewed the relevant literature, the authors emphasize the need to combine multiple methods. For severe pain, pharmacotherapy may be used, while in other cases, it will be more appropriate to apply a combination of splint therapy and physiotherapy. While waiting for a custom-tailored occlusal splint, the patient can take advantage of behavioural and psychological methods, which should be continued after they have been fitted with the splint, as well as during physiotherapy treatments. Follow-up visits are an essential part of the TMD treatment process. The first follow-up visit should take place after one month of therapy and the next after three months. In the meantime, the patient should keep a diary describing their symptoms, pain levels, sleep quality, and wellbeing upon awakening and at bedtime. These observations, which should be reviewed at the follow-up visit, help build a full picture of the effects of the splint and other treatments, as well as inform the psychological assessment of the patient. An accurate diagnosis of TMD is the foundation of appropriate treatment. The most commonly described treatments include physiotherapy, occlusal splint therapy, and pharmacological treatment tailored to the type of TMD.

## Data Availability

No data were used to support this study.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

This work was supported by the Faculty of Medicine and Dentistry, Pomeranian Medical University, Szczecin, Poland.

## References

- [1] C. Di Paolo, D. Costanzo, F. Panti et al., "Epidemiological analysis on 2375 patients with TMJ disorders: basic statistical aspects," *Annali di Stomatologia*, vol. IV, no. 1, pp. 161–169, 2013.
- [2] A. Gil-Martínez, A. Paris-Aleman, I. López-de-Uralde-Villanueva, and R. La Touche, "Management of pain in patients with temporomandibular disorder (TMD): challenges and solutions," *Journal of Pain Research*, vol. 11, no. 11, pp. 571–587, 2018.
- [3] M. Więckiewicz, K. Boening, P. Wiland, Y. Y. Shiau, and A. Paradowska-Stolarz, "Reported concepts for the treatment modalities and pain management of temporomandibular disorders," *The Journal of Headache and Pain*, vol. 16, no. 106, 2015.
- [4] D. T. S. Li, Y. Y. Leung, and T. Disorders, "Current concepts and controversies in diagnosis and management," *Diagnostics*, vol. 11, no. 3, p. 459, 2021.
- [5] J. E. Loster, M. A. Osiewicz, M. Groch, W. Ryniewicz, and A. Wieczorek, "The prevalence of TMD in Polish young adults," *Journal of Prosthodontics*, vol. 26, no. 4, pp. 284–288, 2017.
- [6] L. B. Calixtre, B. L. D. S. Gruninger, T. C. Chaves, and A. B. D. Oliveira, "Is there an association between anxiety/depression and temporomandibular disorders in college students?" *Journal of Applied Oral Science*, vol. 22, no. 1, pp. 15–21, 2014.
- [7] D. Manfredini, F. Piccotti, G. Ferronato, and L. Guardanardini, "Age peaks of different RDC/TMD diagnoses in a patient population," *Journal of Dentistry*, vol. 38, no. 5, pp. 392–399, 2010.
- [8] A. A. Köhler, A. Nydell-Helkimo, T. Magnusson, and A. Hugoson, "Prevalence of symptoms and signs indicative of temporomandibular disorders in children and adolescents, A cross-sectional epidemiological investigation covering two decades," *European Archives of Paediatric Dentistry*, vol. 10, no. S1, pp. 16–25, 2009.
- [9] T. V. Macfarlane, A. S. Blinkhorn, R. M. Davies, J. Kincey, and H. V. Worthington, "Association between female hormonal factors and oro-facial pain: study in the community," *Pain*, vol. 97, no. 1, pp. 5–10, 2002.
- [10] A. Nekora-Azak, "RETRACTED: temporomandibular disorders in relation to female reproductive hormones: a literature review," *The Journal of Prosthetic Dentistry*, vol. 91, no. 5, pp. 491–493, 2004.
- [11] I. M. Nilsson, M. Drangsholt, and T. List, "Impact of temporomandibular disorder pain in adolescents: differences by age and gender," *Journal of Orofacial Pain*, vol. 23, no. 2, pp. 115–122, 2009.
- [12] J. Wang, Y. Chao, Q. Wan, and Z. Zhu, "The possible role of estrogen in the incidence of temporomandibular disorders," *Medical Hypotheses*, vol. 71, no. 4, pp. 564–567, 2008.
- [13] D. Manfredini, A. Bandettini di Poggio, E. Cantini, L. Dell'Osso, and M. Bosco, "Mood and anxiety psychopathology and temporomandibular disorder: a spectrum approach," *Journal of Oral Rehabilitation*, vol. 31, no. 10, pp. 933–940, 2004.
- [14] E. Salameh, F. Alshaarani, H. A. Hamed, and J. A. Nassar, "Investigation of the relationship between psychosocial stress and temporomandibular disorder in adults by measuring salivary cortisol concentration: a case-control study," *Journal of Indian Prosthodontic Society*, vol. 15, no. 2, pp. 148–152, 2015.
- [15] S. Anna, K. Joanna, S. Teresa, G. Maria, and W. Aneta, "The influence of emotional state on the masticatory muscles function in the group of young healthy adults," *BioMed Research International*, vol. 2015, Article ID 174013, 7 pages, 2015.

- [16] A. Stocka, T. Sierpinska, J. Kuc, and M. Golebiewska, "Relationship between depression and masticatory muscles function in a group of adolescents," *Cranio: The Journal of Craniomandibular & Sleep Practice*, vol. 36, no. 6, pp. 390–395, 2018.
- [17] A. Sojka, B. Stelcer, M. Roy, E. Mojs, and M. Pryliński, "Is there a relationship between psychological factors and TMD?" *Brain Behav*, vol. 9, no. 9, Article ID e01360, 2019.
- [18] A. U. J. Yap, E. K. Chua, S. F. Dworkin, H. H. Tan, and K. B. C. Tan, "Multiple pains and psychosocial functioning/psychologic distress in TMD patients," *The International Journal of Prosthodontics*, vol. 15, no. 5, pp. 461–466, 2002.
- [19] A. U. J. Yap, E. K. Chua, and J. K. E. Hoe, "Clinical TMD, pain-related disability and psychological status of TMD patients," *Journal of Oral Rehabilitation*, vol. 29, no. 4, pp. 374–380, 2002.
- [20] A. U. J. Yap, E. K. Chua, and K. B. C. Tan, "Depressive symptoms in Asian TMD patients and their association with non-specific physical symptoms reporting," *Journal of Oral Pathology & Medicine*, vol. 33, no. 5, pp. 305–310, 2004.
- [21] A. U. Yap, K. B. Tan, C. Prosthodont, E. K. Chua, and H. H. Tan, "Depression and somatization in patients with temporomandibular disorders," *The Journal of Prosthetic Dentistry*, vol. 88, no. 5, pp. 479–484, 2002.
- [22] R. Čelić, J. Pandurić, and N. Dulčić, "Psychologic status in patients with temporomandibular disorders," *International Journal of Prosthodontics*, vol. 19, no. 1, pp. 28–29, 2006.
- [23] R. B. Fillingim, R. Ohrbach, J. D. Greenspan et al., "Psychological factors associated with development of TMD: the OPPERA prospective cohort study," *The Journal of Pain*, vol. 14, no. 12, pp. T75–T90, 2013.
- [24] F. Licini, A. Nojelli, M. Segù, and V. Collesano, "Role of psychosocial factors in the etiology of temporomandibular disorders: relevance of a biaxial diagnosis," *Minerva Stomatologica*, vol. 58, no. 11–12, pp. 557–566, 2009.
- [25] H. Kanehira, A. Agariguchi, H. Kato, S. Yoshimine, and H. Inoue, "Association between stress and temporomandibular disorder," *Nippon Hotetsu Shika Gakkai Zasshi*, vol. 52, no. 3, pp. 375–380, 2008.
- [26] D. R. Reissmann, M. T. John, O. Schierz, H. Seedorf, and S. Doering, "Stress-related adaptive versus maladaptive coping and temporomandibular disorder pain," *Journal of Orofacial Pain*, vol. 26, no. 3, pp. 181–190, 2012.
- [27] M. Schmitter, L. Keller, N. Giannakopoulos, and P. Rammelsberg, "Chronic stress in myofascial pain patients," *Clinical Oral Investigations*, vol. 14, no. 5, pp. 593–597, 2010.
- [28] V. G. Augusto, K. C. B. Perina, D. S. G. Penha, D. C. A. d. Santos, and V. A. S. Oliveira, "Temporomandibular dysfunction, stress and common mental disorder in University students," *Acta Ortopédica Brasileira*, vol. 24, no. 6, pp. 330–333, 2016.
- [29] L. R. Bonjardim, M. B. D. Gavião, L. J. Pereira, and P. M. Castelo, "Anxiety and depression in adolescents and their relationship with signs and symptoms of temporomandibular disorders," *The International Journal of Prosthodontics*, vol. 18, no. 4, pp. 347–352, 2005.
- [30] B. D. Furquim, L. M. S. P. Flamengui, and P. C. R. Conti, "TMD and chronic pain: a current view," *Dental Press Journal Orthodontics*, vol. 20, no. 1, pp. 127–133, 2015.
- [31] A. U. J. Yap, S. F. Dworkin, E. K. Chua, T. List, K. B. C. Tan, and H. H. Tan, "Prevalence of temporomandibular disorder subtypes, psychologic distress and psychosocial dysfunction in Asian patients," *Journal of Orofacial Pain*, vol. 17, no. 1, pp. 21–28, 2003.
- [32] D. Manfredini, L. Borella, L. Favero, G. Ferronato, and L. Guarda-Nardini, "Chronic pain severity and depression/somatization levels in TMD patients," *The International Journal of Prosthodontics*, vol. 23, no. 6, pp. 529–534, 2010.
- [33] Y. Yin, S. He, J. Xu et al., "The neuro-pathophysiology ofttemporomandibular disorders-related pain:a systematic review of structural andfunctional MRI studies," *The Journal of Headache and Pain*, vol. 21, no. 1, p. 78, 2020.
- [34] P. E. Bohm, F. F. Stancampiano, and T. D. Rozen, "Migraine headache: updates and future developments," *Mayo Clinic Proceedings*, vol. 93, no. 11, pp. 1648–1653, 2018.
- [35] C. Wober-Bingol, "Epidemiology of migraine and headache in children and adolescents," *Current Pain and Headache Reports*, vol. 17, no. 6, p. 341, 2013.
- [36] D. Bree and D. Levy, "Development of CGRP-dependent pain and headache related behaviours in a rat model of concussion: implications for mechanisms of post-traumatic headache," *Cephalalgia*, vol. 38, no. 2, pp. 246–258, 2018.
- [37] S. Mittrattanakul and R. L. Merrill, "Headache impact in patients with orofacial pain," *Journal of the American Dental Association*, vol. 137, no. 9, pp. 1267–1274, 2006.
- [38] D. A. G. Goncalves, M. E. Bigal, L. C. F. Jales, C. M. Camparis, and J. G. Speciali, "Headache and symptoms of temporomandibular disorder: an epidemiological study," *Headache: The Journal of Head and Face Pain*, vol. 50, no. 2, pp. 231–241, 2010.
- [39] D. Millstine, C. Y. Chen, and B. Bauer, "Complementary and integrative medicine in the management of headache," *BMJ*, vol. 357, p. j1805, 2017.
- [40] V. Ballegaard, P. Thede-Schmidt-Hansen, P. Svensson, and R. Jensen, "Are headache and temporomandibular disorders related? A blinded study," *Cephalalgia*, vol. 28, no. 8, pp. 832–841, 2008.
- [41] R. Ramachandran, "Neurogenic inflammation and its role in migraine," *Seminars in Immunopathology*, vol. 40, no. 3, pp. 301–314, 2018.
- [42] I. de Boer, A. M. van den Maagdenberg, and G. M. Terwindt, "Advance in genetics of migraine," *Current Opinion in Neurology*, vol. 32, no. 3, pp. 413–421, 2019.
- [43] X. Moisset, G. Bommelaer, M. Boube et al., "Migraine prevalence in inflammatory bowel disease patients: a tertiary-care centre cross-sectional study," *European Journal of Pain*, vol. 21, no. 9, pp. 1550–1560, 2017.
- [44] A. A. Gelfand, "Episodic syndromes of childhood associated with migraine," *Current Opinion in Neurology*, vol. 31, no. 3, pp. 281–285, 2018.
- [45] A. Charles, "The pathophysiology of migraine: implications for clinical management," *The Lancet Neurology*, vol. 17, no. 2, pp. 174–182, 2018.
- [46] M. G. Miglis, "Migraine and autonomic dysfunction: which is the horse and which is the jockey?" *Current Pain and Headache Reports*, vol. 22, no. 3, 2018.
- [47] A. L. Franco, D. A. Goncalves, S. M. Castanharo, J. G. Speciali, M. E. Bigal, and C. M. dCamparis, "Migraine is the most prevalent primary headache in individuals with temporomandibular disorders," *Journal of Orofacial Pain*, vol. 24, no. 3, pp. 287–292, 2010.
- [48] S. F. Dworkin, "Temporomandibular disorder (TMD) pain-related disability found related to depression, non-specific physical symptoms, and pain duration at 3 international sites," *Journal of Evidence-Based Dental Practice*, vol. 11, no. 3, pp. 143–144, 2011.

- [49] R. Poveda Roda, J. V. Bagan, J. M. Diaz Fernández, S. Hernández Bazán, and Y. Jiménez Soriano, "Review of temporomandibular joint pathology. Part I: classification, epidemiology and risk factors," *Medicina Oral, Patología Oral Y Cirugía Bucal*, vol. 12, no. 4, pp. E292–E298, 2007.
- [50] D. Manfredini, L. Lombardo, and G. Siciliani, "Temporomandibular disorders and dental occlusion. A systematic review of association studies: end of an era?" *Journal of Oral Rehabilitation*, vol. 44, no. 11, pp. 908–923, 2017.
- [51] M. T. John, S. F. Dworkin, and L. A. Mancl, "Reliability of clinical temporomandibular disorder diagnoses," *Pain*, vol. 118, no. 1, pp. 61–69, 2005.
- [52] E. Schiffman, R. Ohrbach, E. Truelove et al., "Diagnostic criteria for temporomandibular disorders (DC/TMD) for clinical and research applications: recommendations of the international RDC/TMD consortium network\* and orofacial pain special interest group," *The Journal of Oral & Facial Pain and Headache*, vol. 28, no. 1, pp. 6–27, 2014.
- [53] C. C. Peck, J. P. Goulet, F. Lobbezoo et al., "Expanding the taxonomy of the diagnostic criteria for temporomandibular disorders," *Journal of Oral Rehabilitation*, vol. 41, no. 1, pp. 2–23, 2014.
- [54] R. Rongo, E. C. Ekberg, I. M. Nilsson, and A. Michelotti, "Diagnostic criteria for temporomandibular disorders (DC/TMD) for children and adolescents: an international delphi study—Part 1-development of axis I," *Journal of Oral Rehabilitation*, vol. 48, pp. 836–845, 2021.
- [55] E. Kijak, D. Lietz-Kijak, Z. Śliwiński, and B. Frączak, "Muscle activity in the course of rehabilitation of masticatory motor system functional disorders," *Postępy Higieny I Medycyny Doświadczalnej*, vol. 67, pp. 507–516, 2013.
- [56] S. M. Shaffer, J. M. Brismée, P. S. Sizer, and C. A. Courtney, "Temporomandibular disorders. Part 1: anatomy and examination/diagnosis," *Journal of Manual & Manipulative Therapy*, vol. 22, no. 1, pp. 2–12, 2014 Feb.
- [57] R. Akhter, N. M. M. Hassan, J. Aida, T. Kanehira, K. U. Zaman, and M. Morita, "Association between experience of stressful life events and muscle-related temporomandibular disorders in patients seeking free treatment in a dental hospital," *European Journal of Medical Research*, vol. 12, no. 11, pp. 535–540, 2007.
- [58] T. I. Suvinen and P. Kempainen, "Review of clinical EMG studies related to muscle and occlusal factors in healthy and TMD subjects," *Journal of Oral Rehabilitation*, vol. 34, no. 9, pp. 631–644, 2007.
- [59] J. E. Lindroth, J. E. Schmidt, and C. R. Carlson, "A comparison between masticatory muscle pain patients and intracapsular pain patients on behavioral and psychosocial domains," *Journal of Orofacial Pain*, vol. 16, no. 4, pp. 277–283, 2002.
- [60] R. L. Gauer and M. J. Semidey, "Diagnosis and treatment of temporomandibular disorders," *American Family Physician*, vol. 91, no. 6, pp. 378–386, 2015.
- [61] J. Y. Ro, P. Svensson, and N. F. Capra, "Effects of experimental muscle pain on electromyographic activity of masticatory muscles in the rat," *Muscle & Nerve*, vol. 25, no. 4, pp. 576–584, 2002.
- [62] C. Bodere, S. H. Tea, M. A. Giroux-Metges, and A. Woda, "Activity of masticatory muscles in subjects with different orofacial pain conditions," *Pain*, vol. 116, no. 1, pp. 33–41, 2005.
- [63] B. Hatef, S. Talebian, F. Hashemirad, and M. Ghaffarpour, "Effect of pain on the timing pattern of masseter muscle activity during the open-close-clench cycle in the migraine without aura and tension type headaches," *Iranian Journal of Neurology*, vol. 11, no. 4, pp. 146–150, 2012.
- [64] C. Peck, G. Murray, and T. Gerzina, "How does pain affect jaw muscle activity? The integrated pain adaptation model," *Australian Dental Journal*, vol. 53, no. 3, pp. 201–207, 2008.
- [65] L. Bendtsen and C. Fernández-de-la-Peñas, "The role of muscles in tension-type headache," *Current Pain and Headache Reports*, vol. 15, no. 6, pp. 451–458, 2011.
- [66] Y. Dong, X. M. Wang, M. Q. Wang, and S. E. Widmalm, "Asymmetric muscle function in patients with developmental mandibular asymmetry," *Journal of Oral Rehabilitation*, vol. 35, no. 1, pp. 27–36, 2008.
- [67] J. Li, T. Jiang, H. Feng, K. Wang, Z. Zhang, and T. Ishikawa, "The electromyographic activity of masseter and anterior temporalis during orofacial symptoms induced by experimental occlusal highspot," *Journal of Oral Rehabilitation*, vol. 35, no. 2, pp. 79–87, 2008.
- [68] T. Sierpińska, P. Jacunski, J. Kuc, M. Golebiewska, A. Wiczorek, and S. Majewski, "Effect of the dental arches morphology on the masticatory muscles activities in normal occlusion young adults," *Cranio*, vol. 33, no. 2, pp. 134–141, 2015.
- [69] A. Wiczorek, J. Loster, and B. W. Loster, "Relationship between occlusal force distribution and the activity of masseter and anterior temporalis muscles in asymptomatic young adults," *BioMed Research International*, vol. 2013, Article ID 354017, 7 pages, 2013.
- [70] T. K. Goto, T. Yamada, and K. Yoshiura, "Occlusal pressure, contact area, force and the correlation with the morphology of the jaw-closing muscles in patients with skeletal mandibular asymmetry," *Journal of Oral Rehabilitation*, vol. 35, no. 8, pp. 594–603, 2008.
- [71] P. Urbański, B. Trybulec, and M. Pihut, "The application of manual techniques in masticatory muscles relaxation as adjunctive therapy in the treatment of temporomandibular joint disorders," *International Journal of Environmental Research and Public Health*, vol. 18, no. 24, Article ID 12970, 2021.
- [72] S. E. Forrester, S. J. Allen, R. G. Presswood, A. C. Toy, and M. T. G. Pain, "Neuromuscular function in healthy occlusion," *Journal of Oral Rehabilitation*, vol. 37, no. 9, pp. 663–669, 2010.
- [73] D. Rodrigues-Bigaton, K. C. S. Berni, A. F. N. Almeida, and M. T. Silva, "Activity and asymmetry index of masticatory muscles in women with and without dysfunction temporomandibular," *Electromyography & Clinical Neurophysiology*, vol. 50, no. 7-8, pp. 333–338, 2010.
- [74] D. Manfredini, F. Cocilovo, L. Favero, G. Ferronato, S. Tonello, and L. Guarda-Nardini, "Surface electromyography of jaw muscles and kinesiographic recordings: diagnostic accuracy for myofascial pain," *Journal of Oral Rehabilitation*, vol. 38, no. 11, pp. 791–799, 2011.
- [75] V. Scopel, G. S. Alves Da Costa, and D. Urias, "An electromyographic study of masseter and anterior temporalis muscles in extra-articular myogenous TMJ pain patients compared to an asymptomatic and normal population," *Cranio: The Journal of Craniomandibular & Sleep Practice*, vol. 23, no. 3, pp. 194–203, 2005.
- [76] M. K. A. van Selms, F. Lobbezoo, C. M. Visscher, and M. Naeije, "Myofascial temporomandibular disorder pain, parafunctions and psychological stress," *Journal of Oral Rehabilitation*, vol. 35, no. 1, pp. 45–52, 2008.
- [77] D. Božović, N. Ivkovic, M. Račić, and S. Ristić, "Salivary cortisol responses to acute stress in students with myofascial



- pain," *Srpski arhiv za celokupno lekarstvo*, vol. 146, no. 1–2, pp. 20–25, 2018.
- [78] M. B. Yunus, "The prevalence of fibromyalgia in other chronic pain conditions," *Pain Res Treat*, vol. 2012, Article ID 584573, 8 pages, 2012.
- [79] R. Balasubramaniam, R. de Leeuw, H. Zhu, R. B. Nickerson, J. P. Okeson, and C. R. Carlson, "Prevalence of temporomandibular disorders in fibromyalgia and failed back syndrome patients: a blinded prospective comparison study," *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology & Endodontics*, vol. 104, no. 2, pp. 204–216, 2007.
- [80] L. LeResche, L. A. Mancl, M. T. Drangsholt, G. Huang, and M. Von Korff, "Predictors of onset of facial pain and temporomandibular disorders in early adolescence," *Pain*, vol. 129, no. 3, pp. 269–278, 2007.
- [81] A. G. Glaros, K. Williams, and L. Lausten, "The role of parafunctions, emotions and stress in predicting facial pain," *Journal of The American Dental Association*, vol. 136, no. 4, pp. 451–458, 2005.
- [82] S. G. Cortese and A. M. Biondi, "Relationship between dysfunctions and parafunctional oral habits, and temporomandibular disorders in children and teenagers," *Archivos Argentinos de Pediatría*, vol. 107, no. 2, pp. 134–138, 2009.
- [83] R. M. Feteih, "Signs and symptoms of temporomandibular disorders and oral parafunctions in urban Saudi Arabian adolescents: a research report," *Head & Face Medicine*, vol. 2, no. 1, 2006.
- [84] A. Sojka, M. Żarowski, B. Steinborn, W. Hedzelek, B. Spychala, and B. Dorocka-Bobkowska, "Temporomandibular disorders in adolescents with headache," *Advances in Clinical and Experimental Medicine*, vol. 27, no. 2, pp. 193–199, 2018.
- [85] D. E. Harper, A. Schrepf, and D. J. Clauw, "Pain mechanisms and centralized pain in temporomandibular disorders," *Journal of Dental Research*, vol. 95, no. 10, pp. 1102–1108, 2016.
- [86] V. F. Ferrario, G. M. Tartaglia, A. Galletta, G. P. Grassi, and C. Sforza, "The influence of occlusion on jaw and neck muscle activity: a surface EMG study in healthy young adults," *Journal of Oral Rehabilitation*, vol. 33, no. 5, pp. 341–348, 2006.
- [87] A. J. Hassel, P. Rammelsberg, and M. Schmitter, "Inter-examiner reliability in the clinical examination of temporomandibular disorders: influence of age," *Community Dentistry and Oral Epidemiology*, vol. 34, no. 1, pp. 41–46, 2006.
- [88] A. F. Marangoni, C. H. de Godoy, D. A. Biasotto-Gonzalez et al., "Assessment of type of bite and vertical dimension of occlusion in children and adolescents with temporomandibular disorders," *Journal of Bodywork and Movement Therapies*, vol. 18, 2013.
- [89] B. Y. Li, L. J. Zhou, S. X. Guo, Y. Zhang, L. Lu, and M. Q. Wang, "An investigation on the simultaneously recorded occlusion contact and surface electromyographic activity for patients with unilateral temporomandibular disorders pain," *Journal of Electromyography and Kinesiology*, vol. 28, pp. 199–207, 2016.
- [90] M. Funato, Y. Ono, K. Baba, and Y. Kudo, "Evaluation of the non-functional tooth contact in patients with temporomandibular disorders by using newly developed electronic system," *Journal of Oral Rehabilitation*, vol. 41, no. 3, pp. 170–176, 2014.
- [91] L. S. R. Vilanova, R. C. M. R. Garcia, T. List, P. Alstergren, and P. Alstergren, "Diagnostic criteria for temporomandibular disorders: self-instruction or formal training and calibration?" *The Journal of Headache and Pain*, vol. 16, no. 1, 2015.
- [92] G. D. Klasser and J. P. Okeson, "The clinical usefulness of surface electromyography in the diagnosis and treatment of temporomandibular disorders," *Journal of The American Dental Association*, vol. 137, no. 6, pp. 763–771, 2006.
- [93] X. R. Wang, Y. Zhang, N. Xing, Y. F. Xu, and M. Q. Wang, "Stable tooth contacts in intercuspal occlusion makes for utilities of the jaw elevators during maximal voluntary clenching," *Journal of Oral Rehabilitation*, vol. 40, no. 5, pp. 319–328, 2013.
- [94] D. Ciavarella, V. Parziale, M. Mastrovincenzo et al., "Condylar position indicator and T-scan system II in clinical evaluation of temporomandibular intracapsular disease," *Journal of Cranio-Maxillofacial Surgery*, vol. 40, no. 5, pp. 449–455, 2012.
- [95] U. Santana-Mora, J. Cudeiro, M. J. Mora-Bermúdez et al., "Changes in EMG activity during clenching in chronic pain patients with unilateral temporomandibular disorders," *Journal of Electromyography and Kinesiology*, vol. 19, no. 6, pp. e543–e549, 2009.
- [96] M. Venegas, J. Valdivia, M. Javiera Fresno et al., "Clenching and grinding: effect on masseter and sternocleidomastoid electromyographic activity in healthy subjects," *Cranio: The Journal of Craniomandibular & Sleep Practice*, vol. 27, no. 3, pp. 159–166, 2009.
- [97] H. Minakuchi, M. Fujisawa, Y. Abe et al., "Managements of sleep bruxism in adult: a systematic review," *Japanese Dental Science Review*, vol. 58, pp. 124–136, 2022.
- [98] B. Koos, A. Godt, C. Schille, and G. Göz, "Precision of an instrumentation-based method of analyzing occlusion and its resulting distribution of forces in the dental arch," *Journal of Orofacial Orthopedics/Fortschritte der Kieferorthopaedie*, vol. 71, no. 6, pp. 403–410, 2010.
- [99] A. Saraçoğlu and B. Özpınar, "Vivo and in vitro evaluation of occlusal indicator sensitivity," *The Journal of Prosthetic Dentistry*, vol. 88, no. 5, pp. 522–526, 2002.
- [100] R. B. Kerstein, "Combining technologies: a computerized occlusal analysis system synchronized with a computerized electromyography system," *Cranio: The Journal of Craniomandibular & Sleep Practice*, vol. 22, no. 2, pp. 96–109, 2004.
- [101] D. Karakis, M. Bagkur, and B. Toksoy, "Comparison of simultaneously recorded computerized occlusal analysis and surface electromyographic activity of masticatory muscles between patients with unilateral TMD and healthy controls," *The International Journal of Prosthodontics*, vol. 34, no. 5, pp. 554–559, 2021.
- [102] M. Pihut, A. Gala, R. Obuchowicz, and K. Chmura, "Influence of ultrasound examination on diagnosis and treatment of temporomandibular disorders," *Journal of Clinical Medicine*, vol. 11, no. 5, p. 1202, 2022.
- [103] K. Hashimoto, S. Kawashima, S. Kameoka et al., "Comparison of image validity between cone beam computed tomography for dental use and multidetector row helical computed tomography," *Dentomaxillofacial Radiology*, vol. 36, no. 8, pp. 465–471, 2007.
- [104] R. De-Leeuw and G. D. Klasser, "Orofacial pain: guidelines for assessment, diagnosis, and management," *American Academy of Orofacial Pain*, Quintessence, Newhaven, UK, 5th edition, 2013.
- [105] J. P. Okeson, *The Clinical Management of Temporomandibular Disorders and Occlusion*, Mosby, Maryland, Mo, United States, 7th edition, 2013.

- [106] M. Naeije, A. H. Te Veldhuis, E. C. Te Veldhuis, C. M. Visscher, and F. Lobbezoo, "Disc displacement within the human temporomandibular joint: a systematic review of a 'noisy annoyance,'" *Journal of Oral Rehabilitation*, vol. 40, no. 2, pp. 139–158, 2013.
- [107] E. L. Schiffman, J. O. Look, J. S. Hodges et al., "Randomized effectiveness study of four therapeutic strategies for TMJ closed lock," *Journal of Dental Research*, vol. 86, no. 1, pp. 58–63, 2007.
- [108] B. Craane, P. U. Dijkstra, K. Stappaerts, and A. De Laat, "Randomized controlled trial on physical therapy for TMJ closed lock," *Journal of Dental Research*, vol. 91, no. 4, pp. 364–369, 2012.
- [109] I. Q. Gutiérrez, H. Sábado-Bundó, and C. Gay-Escoda, "Intraarticular injections of platelet rich plasma and plasma rich in growth factors with arthrocentesis or arthroscopy in the treatment of temporomandibular joint disorders: a systematic review," *Journal of Stomatology, Oral and Maxillofacial Surgery*, vol. 123, no. 5, pp. e327–e335, 2022.
- [110] A. U. Yap, X. H. Zhang, Y. Cao, and K. Y. Fu, "Functional, physical and psychosocial impact of degenerative temporomandibular joint disease," *Journal of Oral Rehabilitation*, vol. 49, no. 3, pp. 301–308, 2022.
- [111] M. Wu, F. T. Almeida, and R. Friesen, "A systematic review on the association between clinical symptoms and CBCT findings in symptomatic TMJ degenerative joint disease," *J Oral Facial Pain Headache*, vol. 35, no. 4, pp. 332–345, 2021.
- [112] M. Tuijt, J. H. Koolstra, F. Lobbezoo, and M. Naeije, "Biomechanical modeling of open locks of the human temporomandibular joint," *Clinical Biomechanics*, vol. 27, no. 8, pp. 749–753, 2012.
- [113] D. W. Nitzan, "Temporomandibular joint "open lock" versus condylar dislocation: signs and symptoms, imaging, treatment, and pathogenesis," *Journal of Oral and Maxillofacial Surgery*, vol. 60, no. 5, pp. 506–511, 2002.
- [114] AA. Garstka, M. Brzózka, A. Bitenc-Jasiejko et al., "Cause-effect relationships between painful TMD and postural and functional changes in the musculoskeletal system: a preliminary report," *Pain Research and Management*, vol. 202214 pages, 2022.
- [115] I. Egermark, T. Magnusson, and G. E. Carlsson, "A 20-year follow-up of signs and symptoms of temporomandibular disorders and malocclusions in subjects with and without orthodontic treatment in childhood," *The Angle Orthodontist*, vol. 73, no. 2, pp. 109–115, 2003.
- [116] F. Lobbezoo, J. Ahlberg, A. G. Glaros et al., "Bruxism defined and graded: an international consensus," *Journal of Oral Rehabilitation*, vol. 40, no. 1, pp. 2–4, 2013.
- [117] A. Pergamalian, T. E. Rudy, H. S. Žaki, and C. M. Greco, "The association between wear facets, bruxism, and severity of facial pain in patients with temporomandibular disorders," *The Journal of Prosthetic Dentistry*, vol. 90, no. 2, pp. 194–200, 2003.
- [118] S. Bulanda, D. Ilczuk-Rypuła, A. Nitecka-Buchta, Z. Nowak, S. Baron, and L. Postek-Stefańska, "Sleep bruxism in children: etiology, diagnosis, and treatment-A literature review," *International Journal of Environmental Research and Public Health*, vol. 18, no. 18, p. 9544, 2021.
- [119] D. Manfredini and F. Lobbezoo, "Role of psychosocial factors in the etiology of bruxism," *Journal of Orofacial Pain*, vol. 23, no. 2, pp. 153–166, 2009.
- [120] G. Melo, J. Duarte, P. Pauletto et al., "Bruxism: an umbrella review of systematic reviews," *Journal of Oral Rehabilitation*, vol. 46, no. 7, pp. 666–690, 2019.
- [121] C. Sato, S. Sato, H. Takashina, H. Ishii, M. Onozuka, and K. Sasaguri, "Bruxism affects stress responses in stressed rats," *Clinical Oral Investigations*, vol. 14, no. 2, pp. 153–160, 2010.
- [122] A. Wiczorek and J. E. Loster, "Activity of the masticatory muscles and occlusal contacts in young adults with and without orthodontic treatment," *BMC Oral Health*, vol. 15, no. 1, Article ID 116, 2015.
- [123] S. E. Forrester, R. G. Presswood, A. C. Toy, and M. T. G. Pain, "Occlusal measurement method can affect SEMG activity during occlusion," *Journal of Oral Rehabilitation*, vol. 38, no. 9, pp. 655–660, 2011.
- [124] D. Lietz-Kijak, Ł. Kopacz, R. Ardan, M. Grzegocka, and E. Kijak, "Assessment of the short-term effectiveness of kinesiotaping and trigger points release used in functional disorders of the masticatory muscles," *Pain Research and Management*, vol. 2018, no. 2, Article ID 5464985, 7 pages, 2018.
- [125] Ł. Kopacz, Ż. Ciosek, H. Gronwald, P. Skomro, R. Ardan, and D. Lietz-Kijak, "Comparative analysis of the influence of selected physical factors on the level of pain in the course of temporomandibular joint disorders," *Pain Research and Management*, vol. 2020, no. 2, 8 pages, 2020.
- [126] S. Hu, Z. Shao, and L. Deng, "Clinical manifestations, imaging features, and pathogenic/prognostic risk factors for temporomandibular disorders (TMD): a case-control study based on psychogenic factors of patients," *Computational and Mathematical Methods in Medicine*, vol. 25, Article ID 8279357, 2022.
- [127] S. B. Syrop, "Initial management of temporomandibular disorders," *Dentistry Today*, vol. 21, no. 8, pp. 52–57, 2002.
- [128] M. M. Hamata, P. R. J. Zuim, and A. R. Garcia, "Comparative evaluation of the efficacy of occlusal splints fabricated in centric relation or maximum intercuspation in temporomandibular disorders patients," *Journal of Applied Oral Science*, vol. 17, no. 1, pp. 32–38, 2009.
- [129] A. Sabhlok, S. Gupta, M. Girish, K. V. Rahul Ramesh, H. Shrivastava, and S. Hota, "Practice of occlusal splint therapy for treating temporomandibular disorders by general dentists of Jabalpur - a cross-sectional survey," *Journal of Pharmacy and BioAllied Sciences*, vol. 13, no. 2, pp. S1079–S1083, 2021.
- [130] J. Kuzmanovic Pfcicer, S. Dodic, V. Lazic, G. Trajkovic, N. Milic, and B. Milicic, "Occlusal stabilization splint for patients with temporomandibular disorders: meta-analysis of short and long term effects," *PLoS One*, vol. 12, no. 2, Article ID e0171296, 2017.
- [131] M. Z. Al Ani, S. J. Davies, P. Sloan, and A. M. Glenny, "Stabilization splint therapy for TM pain dysfunction syndrome," *The Cochran Database of Systematic Reviews*, vol. 4, 2016.
- [132] C. Zhang, J. Y. Wu, D. L. Deng et al., "Efficacy of splint therapy for the management of temporomandibular disorders: a meta-analysis," *Oncotarget*, vol. 7, no. 51, pp. 84043–84053, 2016.
- [133] B. Wiechens, S. Paschereit, T. Hampe, T. Wassmann, N. Gersdorff, and R. Bürgers, "Changes in maximum mandibular mobility due to splint therapy in patients with temporomandibular disorders," *Health Care*, vol. 10, no. 6, Article ID 1070, 2022.
- [134] M. S. Medlicott and S. R. Harris, "A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of

- temporomandibular disorder,” *Physical Therapy*, vol. 86, no. 7, pp. 955–973, 2006.
- [135] S. Armijo-Olivo, L. Pitance, V. Singh, F. Neto, N. Thie, and A. Michelotti, “Effectiveness of manual therapy and therapeutic exercise for temporomandibular disorders: systematic review and meta-analysis,” *Physical Therapy*, vol. 96, no. 1, pp. 9–25, 2016.
- [136] T. Yabe, T. Tsuda, S. Hirose, T. Ozawa, and K. Kawai, “Treatment of acute temporomandibular joint dislocation using manipulation technique for disk displacement,” *Journal of Craniofacial Surgery*, vol. 25, no. 2, pp. 596–597, 2014.
- [137] L. B. Calixtre, R. F. C. Moreira, G. H. Franchini, F. Albuquerque-Sendin, and A. B. Oliveira, “Manual therapy for the management of pain and limited range of motion in subjects with signs and symptoms of temporomandibular disorder: a systematic review of randomised controlled trials,” *Journal of Oral Rehabilitation*, vol. 42, no. 11, pp. 847–861, 2015 Nov.
- [138] B. M. F. Alves, C. R. Macedo, E. Januzzi, E. Grossmann, A. N. Atallah, and S. Peccin, “Mandibular manipulation for the treatment of temporomandibular disorder,” *Journal of Craniofacial Surgery*, vol. 24, no. 2, pp. 488–493, 2013.
- [139] V. Kasat, A. Gupta, R. Ladda, M. Kathariya, H. Saluja, and A. A. Farooqui, “Transcutaneous electric nerve stimulation (TENS) in dentistry- A review,” *Journal of Clinical and Experimental Dentistry*, vol. 6, no. 5, pp. e562–e568, 2014.
- [140] D. M. Walsh, T. E. Howe, M. I. Johnson, and K. A. Sluka, “Transcutaneous electrical nerve stimulation for acute pain,” *Cochrane Database of Systematic Reviews*, John Wiley & Sons, Ltd, no. 2, Hoboken, NY, USA, Article ID CD006142, 2008.
- [141] K. H. Awan and S. Patil, “The role of transcutaneous electrical nerve stimulation in the management of temporomandibular joint disorder,” *The Journal of Contemporary Dental Practice*, vol. 16, no. 12, pp. 984–986, 2015.
- [142] U. S. Pal, N. Singh, G. Singh et al., “Trends in management of myofascial pain,” *National Journal of Maxillofacial Surgery*, vol. 5, no. 2, pp. 109–116, 2014.
- [143] M. Mostafavifar, J. Wertz, and J. Borchers, “A systematic review of the effectiveness of Kinesio taping for musculoskeletal injury,” *The Physician and Sportsmedicine*, vol. 40, no. 4, pp. 33–40, 2012.
- [144] K. Kase, J. Wallis, and T. Kase, *Clinical Therapeutic Applications of the Kinesio Taping Method*, Kinesio Taping Association International, Albuquerque, NM, USA, 3rd edition, 2013.
- [145] E. Kaya, M. Zinnuroglu, and I. Tugcu, “Kinesio taping compared to physical therapy modalities for the treatment of shoulder impingement syndrome,” *Clinical Rheumatology*, vol. 30, no. 2, pp. 201–207, 2011.
- [146] P. L. Chen, W. H. Hong, C. H. Lin, and W. C. Chen, “Biomechanics effects of Kinesio taping for persons with patellofemoral pain syndrome during stair climbing,” in *Proceedings of the 4th Kuala Lumpur International Conference on Biomedical Engineering IFMBE*, pp. 395–397, Springer, Berlin, Heidelberg, June 2008.
- [147] R. Butts, J. Dunning, R. Pavkovich, J. Mettelle, and F. Mourad, “Conservative management of temporomandibular dysfunction: a literature review with implications for clinical practice guidelines (Narrative review part 2),” *Journal of Bodywork and Movement Therapies*, vol. 21, no. 3, pp. 541–548, 2017.
- [148] A. Andre, J. Kang, and H. Dym, “Pharmacologic treatment for temporomandibular and temporomandibular joint disorders,” *Oral and Maxillofacial Surgery Clinics of North America*, vol. 34, no. 1, pp. 49–59, 2022.
- [149] E. V. Hersh, R. Balasubramaniam, and A. Pinto, “Pharmacologic management of temporomandibular disorders,” *Oral and Maxillofacial Surgery Clinics of North America*, vol. 20, no. 2, pp. 197–210, 2008.
- [150] E. D. de Andrade, C. M. Rizzatti-Barbosa, and M. L. Pimenta Pinheiro, “Pharmacological guidelines for managing temporomandibular disorders,” *Brazilian Journal of Oral Sciences*, vol. 3, pp. 503–505, 2004.
- [151] A. Ouanounou, M. Goldberg, and D. A. Haas, “Pharmacotherapy in Temporomandibular Disorders: A Review,” *Journal of the Canadian Dental Association*, vol. 83, 2017.
- [152] Y. Sharav and R. Benoliel, “Pharmacotherapy of acute orofacial pain,” in *Orofacial Pain and Headache*, Y. Sharav and R. Benoliel, Eds., pp. 349–376, C. V. Mosby, Maryland, MO, USA, 2008.
- [153] S. Bindu, S. Mazumder, and U. Bandyopadhyay, “Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: a current perspective,” *Biochemical Pharmacology*, vol. 180, Article ID 114147, 2020.
- [154] T. Dreischulte, D. R. Morales, S. Bell, and B. Guthrie, “Combined use of nonsteroidal anti-inflammatory drugs with diuretics and/or renin-angiotensin system inhibitors in the community increases the risk of acute kidney injury,” *Kidney International*, vol. 88, no. 2, pp. 396–403, 2015.
- [155] G. Hijos-Mallada, C. Sostres, and F. Gomollón, “NSAIDs, gastrointestinal toxicity and inflammatory bowel disease,” *Gastroenterología Y Hepatología*, vol. 45, no. 3, pp. 215–222, 2022.
- [156] J. Kunjur, R. Anand, P. A. Brennan, and V. Ilankovan, “An audit of 405 temporomandibular joint arthrocentesis with intra-articular morphine infusion,” *British Journal of Oral and Maxillofacial Surgery*, vol. 41, no. 1, pp. 29–31, 2003.
- [157] K. Hayashi, M. Sugisaiki, S. Ota, and H. Tanabe, “ $\mu$ -Opioid receptor mRNA expression and immunohistochemical localization in the rat temporomandibular joint,” *Peptides*, vol. 23, no. 5, pp. 889–893, 2002.
- [158] Y. Liu, J. S. Wu, Y. L. Tang, Y. J. Tang, W. Fei, and X. H. Liang, “Multiple treatment meta-analysis of intra-articular injection for temporomandibular osteoarthritis,” *Journal of Oral and Maxillofacial Surgery*, vol. 78, no. 3, pp. 373.e1–373.e18, 2020.
- [159] P. M. Mountziaris, P. R. Kramer, and A. G. Mikos, “Emerging intra-articular drug delivery systems for the temporomandibular joint,” *Methods*, vol. 47, no. 2, pp. 134–140, 2009.
- [160] L. Fredriksson, P. Alstergren, and S. Kopp, “Serotonergic mechanisms influence the response to glucocorticoid treatment in TMJ arthritis,” *Mediators of Inflammation*, vol. 2005, no. 4, pp. 194–201, 2005.
- [161] P. Marty, A. Louvrier, E. Weber, P. A. Dubreuil, B. Chatelain, and C. Meyer, “Arthrocentèse de l’articulation temporomandibulaire et injection(s) intra-articulaire(s): un état des lieux [Arthrocentesis of the temporomandibular joint and intra-articular injections: an update],” *Revue de Stomatologie, de Chirurgie Maxillo-faciale et de Chirurgie Orale*, vol. 117, no. 4, pp. 266–272, 2016.
- [162] B. Arabshahi, E. M. Dewitt, A. M. Cahill et al., “Utility of corticosteroid injection for temporomandibular arthritis in children with juvenile idiopathic arthritis,” *Arthritis & Rheumatism*, vol. 52, no. 11, pp. 3563–3569, 2005.

- [163] Z. Shi, C. Guo, and M. Awad, "WITHDRAWN: hyaluronate for temporomandibular joint disorders," *Cochrane Database of Systematic Reviews*, vol. 8, no. 10, Article ID CD002970, 2013.
- [164] D. G. Borenstein and S. Korn, "Efficacy of a low-dose regimen of cyclobenzaprine hydrochloride in acute skeletal muscle spasm: results of two placebo-controlled trials," *Clinical Therapeutics*, vol. 25, no. 4, pp. 1056–1073, 2003.
- [165] H. Moldofsky, H. W. Harris, W. T. Archambault, T. Kwong, and S. Lederman, "Effects of bedtime very low dose cyclobenzaprine on symptoms and sleep physiology in patients with fibromyalgia syndrome: a double-blind randomized placebo-controlled study," *Journal of Rheumatology*, vol. 38, no. 12, pp. 2653–2663, 2011.
- [166] G. J. Sills, "The mechanisms of action of gabapentin and pregabalin," *Current Opinion in Pharmacology*, vol. 6, no. 1, pp. 108–113, 2006.
- [167] Y. Haviv, A. Rettman, D. Aframian, Y. Sharav, and R. Benoliel, "Myofascial pain: an open study on the pharmacotherapeutic response to stepped treatment with tricyclic antidepressants and gabapentin," *The Journal of Oral & Facial Pain and Headache*, vol. 29, no. 2, pp. 144–151, 2015.
- [168] L. M. Arnold, D. L. Goldenberg, S. B. Stanford et al., "Gabapentin in the treatment of fibromyalgia: a randomized, double-blind, placebo-controlled, multi center trial," *Arthritis & Rheumatism*, vol. 56, no. 4, pp. 1336–1344, 2007.