NEW NORTH AMERICAN ODONATA.—II.

BY ALBERT P. MORSE, WELLESLEY, MASS.

Nehalennia gracilis sp. nov.

Very similar to *N. irene* but even slenderer and differing in structure and markings. Post-ocular spots present in the shape of a continuation of the occipital line half-way to each eye. ♂: segments 9 and 10 all blue; 8 blue except a very narrow transverse basal band. Superior appendages one-third (when well extruded) as long as the side of 10, in side view about two-thirds as broad as long, directed backward, upper margin convex, lower margin slightly concave, apex rounded; in dorsal view about twice as broad as long, the inner margins nearly contiguous, the distal margin parallel with the excised margin of 10 but a little concave, the lateral and inner margins equal, slightly convex, directed caudad, the outer apical angle well rounded, the inner less so. Inferior appendages very short, the upper outer angle produced caudad into a stout apical tooth which projects nearly as far as the superior appendages; in dorsal view these teeth are seen at the sides of the latter, directed outward and backward, their inner margins concave, outer margins convex. ♀: similar to *irene* but the hind margin of the prothorax is excavate medially making it bilobed (instead of trilobed as in that species); the markings differ in that a large triangular apical spot on 9 and all of 10 are blue. In three examples the spot on 9 is produced to base and the dorsum of 8 also is nearly or entirely blue medially, widest behind.

Abd.: ♂, 20-22; ♀, 20-22. Hind wing: ♂, 13-14.5; ♀, 14-15.

2 pairs, 13 ♂, 11 ♀, June 24, Sherborn, Mass.; 6 ♂, 2 ♀, June 19; 1 ♀, June 27; 1 ♂, July 19, Wellesley, Mass.

Taken near stagnant pools in peat-bogs in company with *irene*, with which species it has hitherto been confused, as shown by the series in the Hagen collection.

Enallagma laterale sp. nov.

Post-ocular spots of moderate size. Mid-dorsal thoracic and humeral dark stripes of medium width. Abdomen blue with the dorsum black as follows: apical half of 2, fourth of 3, third of 4, half of 5, three-fourths of 6, all of 7 except an interrupted basal ring, a stripe each side on 8, and all of 10. Dorsum of 10 emarginate. Superior appendages short, one-fourth to one-third as long as side of 10, in side view nearly as broad as long, directed caudad or slightly deflexed, sides parallel, apex blunt, emarginate; in dorsal view as broad as long, widened apically, emarginate at end, apex of both limbs showing equally, smoothly rounded, the upper presenting a little tooth on inner margin. Seen from the left dorso-lateral aspect the right (and *vice versa*) appendage is seen to be deeply rounded-emarginate, or slightly bifurcate, the lower limb usually slightly larger and longer. Inferior appendages nearly twice as long as the superior, stout at base but much narrowed dorsally, directed backward and a little upward, sides straight, tapering smoothly to the slightly upturned and incurved tip.

Abd.: ♂, 20.5-22. Hind wing: ♂, 15-16.

14 ♂, May 25; 9 ♂, June 8, Wellesley, Mass.

Enallagma pictum sp. nov.

Allied to *signatum*. Coloration black and yellow. Post-ocular spots reduced to a continuation of the occipital line. A pair of very small pale spots between the anterior ocellus and the posterior two. Dorsum of thorax chiefly black, humeral pale stripes narrow, ♂, or very narrow, ♀. ♂: abdomen black as follows: dorsum of 1-10, on 2 in form of an orbicular apical spot narrowly connected to base, on 3 terminating in a sharp point basally. 1 with wide apical, and 4-7 with narrow interrupted basal ring, yel-

low. Superior appendages two and a half times as long as wide, nearly as long as 10, directed backward, equal or slightly enlarged apically, dorsal margin one-fifth longer, slightly concave, apex obliquely truncate, angles rounded, distal margin concave or shallowly emarginate. In dorsal view stouter at base, only twice as long as wide, the upper limb tapering to a rounded tip which bears a short, stout tooth on the inner side next the rounded apex; the lower limb projects inward making the two sides of the whole appendage nearly parallel, the inner apical margin truncate. The inferior appendages are three-fifths as long as the superior, stout, tapering to a blunt point, directed caudad, a little incurved. ♀: humeral pale stripe one-third as wide as the dark stripe, a line on second lateral suture, and a short line between it and the humeral stripe. Dorsum of 1-10 greenish black, not narrowed, more or less maculate on 10: 3-6 with interrupted basal ring, 7 and 8 with apical margin yellow.

Abd.: ♂, 25; ♀, 26.5. Hind wing: ♂, 16.5; ♀, 18.5.

1 ♂, 1 ♀, Sherborn, Mass., collected by Mr. A. L. Babcock, to whom I am indebted for an opportunity to examine and describe them and who has donated the types to the Hagen collection at Cambridge, Mass., in conjunction with the other types described by me from my own collection in this and the preceding papers.

NOTES ON MOTHS.

Amphion nesus.—Last year I found young larvae of *Amphion nesus* on *Ampelopsis veitchii*, and this year I have found them on *Ampelopsis quinquefolia*, one specimen being of a clear wine-color, with subdorsal and oblique lines pink edged above with deep claret-color. The granulation was pale yellow, as were the face lines and the edge of the anal plate. The larvae drop from the vine at a very slight shake and are easily found in this way.

"Red-fringed" *Actias luna*.—In early June Miss Norton sent me eggs of the *A. luna* having deep red fringes on its wings. These

eggs hatched twelve days after they were laid, and the larvae differed from any I had seen in having a dark brown lateral band from head to anal shield; a dark brown patch on the dorsum of the third and of the pre-anal segments, with the anal plate blackish-brown. The bristles from the tubercles were black with some white ones. The feet were green with a brown patch on each. At the first moult they came out like normal *Luna* larvae.

Attacus promethea.—Is this moth rare this year? I have placed on my window-sill nine females, but in no case has a male been attracted. Last year every female which emerged in my box attracted from five to forty males, and some of the females were not even near the open window.

Caroline G. Soule.

Brookline, Mass., June 30, 1895.

PROCEEDINGS OF THE CLUB.

8 March, 1895. The 186th meeting was held at 156 Brattle St., Mr. S. Henshaw in the chair. Mr. A. G. Mayer was chosen secretary.

Mr. W. L. W. Field was elected a member.

Mr. S. H. Scudder read the delayed address of the retiring President, Mr. T. E. Bean, entitled "A comparison of *Colias hecla* with *Colias meadii* and *Colias elis*."

Mr. S. Henshaw asked what the distribution of *C. hecla* was. Mr. Scudder stated that it was quoted from Greenland, Lapland, Hudson Bay, Bering Straits and Grinnell-land and *C. meadii* from Colorado, Arizona, and New Mexico.

A paper by Mr. W. S. Blatchley on winter collecting (see *Psyche*, vii, 247) was read. Mr. Henshaw stated that he had found between 2-300 species of Coleoptera and a few larvae by sifting leaves during the winter months; the greater number of species thus found were Pselaphidae and Staphylinidae.

Mr. Henshaw exhibited a specimen of *Lycaena xerces* from San Francisco. This species was erroneously supposed to have been exterminated by the extirpation of its food plant.