

its absence in *venustum*. In the Australian *M. tyndalli*, the blade-like portion of the caudal wing was absent. *Neomymar* has normal hind wings except that the petiole is long as in *Mymar*.

---

A METALLIC SPECIES OF *CIRROSPILOPSIS* FROM  
MARYLAND (HYMENOPTERA EULOPHIDÆ).

BY A. A. GIRAULT,

Bureau of Entomology, Washington, D. C.

The following species resembles a *Diaulinopsis* but there are no grooves on the scutellum, the caudal tibial spur is single and the club without a distinct apical nipple.

*Cirrospilopsis metallicus* sp. nov.

*Female*: Length, 1.10 mm. Brassy metallic green, the wings hyaline, the legs (except the coxæ; apex of first coxæ yellow), venation and the antennæ (except the scape and pedicel except both at apex), golden yellow. Body densely scaly, the parapsidal furrows distinct, the propodeum moderately short, noncarinate (a median carina subobsolete), the spiracles small, rounded, cephalad. Bristles on thorax minute, sparse. Postscutellum rather large. Mandibles 4-dentate, the last two teeth much smaller than the others. Pedicel subequal to funicle 1 which is nearly twice longer than wide, 2 somewhat shorter, the club somewhat longer than the funicle; ring-joints distinct. Middle tibial spur long, slender. Stigmal vein not quite half the length of the marginal, the postmarginal two-thirds the length of the marginal.

From two females captured by sweeping in the woods, Glenndale, Md., May 4, 1916.

*Types*: Catalogue No. 20291, U. S. National Museum, the specimens on a tag, a head and two caudal tibiæ on a slide.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

