

Immunohistochemical analysis of histone H3 modification in newt tail tissue cells following amputation

Jiwen Wu^{1,2,3}, Xu Zhang¹, Sekiya Reiko¹, Kiyoshi Aoyagi³, Tao-Sheng Li^{1*}

¹Department of Stem Cell Biology, Atomic Bomb Disease Institute, Nagasaki University, Nagasaki, 852-8523, Japan

²Department of Human Anatomy, Histology and Embryology, School of Basic Medical Sciences, Fujian Medical University, Fuzhou, 350108, China

³Department of Public Health, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki, 852-8523, Japan

***Corresponding author:**

Tao-Sheng Li, MD, Ph.D.

Department of Stem Cell Biology,

Atomic Bomb Disease Institute,


Nagasaki University,

1-12-4 Sakamoto, Nagasaki 852-8523, Japan.


Tel: +81-95-819-7099; Fax: +81-95-819-7100. E-mail: litaoshe@nagasaki-u.ac.jp

Supplemental information


Supplementary Figure 1. Schematic diagram on tissue sampling for experiments.




Supplementary Figure 2. Immunohistochemical analysis on the expression of acetylated H3K9 (AcH3K9) in tissue cells 3 days after amputation. Representative images display in grayscale.



Supplementary Figure 3. Immunohistochemical analysis the expression of acetylated H3K14 (AcH3K14) in tissue cells 3 days after amputation. Representative images display in grayscale.



Supplementary Figure 4. Immunohistochemical analysis on the expression of acetylated H3K27 (AcH3K27) in tissue cells 3 days after amputation. Representative images display in grayscale.



Supplementary Figure 5. Immunohistochemical analysis on the expression of methylated H3K14 (MeH3K27) in tissue cells 3 days after amputation. Representative images display in grayscale.

