

Research Article

Serial Gene Expression Profiling of Neural Stem Cells Shows Transcriptome Switch by Long-Term Physioxia from Metabolic Adaption to Cell Signaling Profile

Lena Braunschweig,^{1,2} Jennifer Lanto,³ Anne K. Meyer,¹ Franz Markert ,³
and Alexander Storch ^{1,2,3,4}

¹Division of Neurodegenerative Diseases, Department of Neurology, Technische Universität Dresden, 01307 Dresden, Germany

²Center for Regenerative Therapies Dresden (CRTD), Technische Universität Dresden, 01307 Dresden, Germany

³Department of Neurology, University of Rostock, 18147 Rostock, Germany

⁴German Centre for Neurodegenerative Diseases (DZNE) Rostock/Greifswald, 18147 Rostock, Germany

Correspondence should be addressed to Alexander Storch; alexander.storch@med.uni-rostock.de

Received 17 May 2022; Revised 30 September 2022; Accepted 26 October 2022; Published 12 November 2022

Academic Editor: Darius Widera

Copyright © 2022 Lena Braunschweig et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oxygen is an essential factor in the cellular microenvironment with pivotal effects on neural development with a particular sensitivity of midbrain neural stem cells (NSCs) to high atmospheric oxygen tension. However, most experiments are still performed at atmospheric O₂ levels (21%, normoxia), whereas mammalian brain tissue is physiologically exposed to substantially lower O₂ tensions around 3% (physioxia). We here performed serial Affymetrix gene array analyses to detect expression changes in mouse fetal NSCs from both midbrain and cortical tissues when kept at physioxia compared to normoxia. We identified more than 400 O₂-regulated genes involved in cellular metabolism, cell proliferation/differentiation, and various signaling pathways. NSCs from both regions showed a low number but high conformity of regulated genes (9 genes in midbrain vs. 34 in cortical NSCs; 8 concordant expression changes) after short-term physioxia (2 days) with *metabolic processes* and *cellular processes* being the most prominent GO categories pointing to cellular adaption to lower oxygen levels. Gene expression profiles changed dramatically after long-term physioxia (13 days) with a higher number of regulated genes and more diverse expression patterns when comparing the two NSC types (338 genes in midbrain vs. 121 in cortical NSCs; 75 concordant changes). Most prominently, we observed a reduction of hits in *metabolic processes* but an increase in *biological regulation* and *signaling* pointing to a switch towards signaling processes and stem cell maintenance. Our data may serve as a basis for identifying potential signaling pathways that maintain stem cell characteristics in cortical versus midbrain physioxia stem cell niches.

1. Introduction

Multipotent tissue-specific neural stem cells (NSCs) with their capacity for self-renewal and multilineage differentiation potentially guarantee increasing cell numbers and diversity during fetal brain development as well as neurogenesis in adult neurogenic brain regions [1, 2]. There is a high interest in NSCs and their derivatives as promising tools for stem cell therapies, tissue engineering, pharmacological testing, and modelling of the development of the brain and

peripheral nervous system tissue [3–8]. Studies of the micro-environmental demands of these NSCs within the neuro-genic niche are thus crucial to maintain their survival, proliferation, and differentiation potential *in vitro* and *in vivo* [9–15]. Oxygen tension is one key factor of the neuro-genic niche regulating cellular functions in stem cells, and lowered oxygen tension of around 3% is beneficial for stem cell proliferation, survival, and maintenance in various NSC types both *in vitro* and *in vivo* [16–21]. Indeed, the physiological oxygen tension within the mammalian brain

ranges from 1% to 5% O₂ (known as physioxia [22, 23]) and thus differs remarkably from the atmospheric oxygen condition routinely used in *in vitro* culture [24–26]. Moreover, sensitivity to oxygen tension seems to vary between NSCs from different fetal brain regions with NSCs isolated from the midbrain showing a selective vulnerability to oxygen as compared to cortical NSCs (cNSCs) [18, 27, 28]: reduced O₂ is vital to *in vitro* cultures of fetal midbrain-derived NSCs (mNSCs) along with proliferation, survival, and potential to dopaminergic differentiation [16, 18, 19, 28, 29].

The mechanisms of oxygen sensitivity of NSCs including the differences between the various NSC types remain however in part enigmatic. Moreover, there are some inconsistencies about the underlying factors which may be caused by differences not only of the tissue of origin of the NSCs but also in the oxygen conditions with levels close to anoxia (<1% O₂) and/or in the cultivation periods in altered oxygen tensions [30–32]. In physioxia, hypoxia-inducible factors (Hifs) are the main regulators of O₂ homeostasis through their target genes such as *vascular endothelial growth factor (Vegf)* or *phosphoglycerate kinase 1 (Pgk1)* inducing the hypoxic response including increased angiogenesis or altered glucose metabolism [29, 33–35]. Consistently, neural-specific Hif-1 α conditional knock-out models showed that expression of Hif-1 α in neural cells is essential for the normal development of the brain [36, 37]. Moreover, in our previous study, we demonstrated a selective vulnerability of mNSCs against Hif-1 α conditional knock-out leading to impaired proliferation, survival, and dopaminergic differentiation solely of mNSCs [28]. Another signaling pathway involved on oxygen-mediated NSC behavior, which is independent of Hif-1 α , is the Wnt/ β -catenin pathway positively stimulating proliferation of physioxia cells by affecting cell cycle regulation [27].

In the present study, we focus on whole transcriptome analyses using gene chip microarrays of fetal mesencephalic and cortical NSCs cultured in normoxia (21% atmospheric oxygen tension) or physioxia (3% oxygen tension) to provide a framework for future studies on the molecular mechanisms of oxygen adaptation of various types of fetal NSC involved in stem cell proliferation, survival, and maintenance.

2. Materials and Methods

2.1. Neural Stem Cell Isolation, Culture, and Treatments. For the present whole microarray analyses, we used the biosamples published previously by Braunschweig and colleagues in 2015 [27]. In brief, wild-type C57Bl/6J mice were purchased from Charles River, Sulzfeld, Germany, and animal procedures were approved by the Animal Rights Committee. Embryos at day 14 were removed by Caesarean section, and mesencephalic and cortical brain regions were dissected. After removal of the meninges, the tissue samples were trypsinized (2.5 ng/ml, Sigma-Aldrich, Seelze, Germany) for 10 min at room temperature, incubated in DNase (40 μ g/ml, Sigma-Aldrich, Seelze, Germany) for 10 min at 37°C, and subsequently triturated to achieve a single-cell suspension. For monolayer cultures, cells were plated onto poly-L-ornithine/fibronectin-precoated culture dishes and maintained in an

expansion medium composed of DMEM (high glucose) supplemented with 32% F12, 2% B27, and 1% penicillin/streptomycin (all from Life Technologies, Darmstadt, Germany) and 20 ng/ml EGF and FGF2 (Sigma-Aldrich, Hamburg, Germany). Medium change was performed three times per week. For the hypoxic culture, cells were maintained in a gas mixture composed of 92% N₂, 5% CO₂, and 3% O₂. To preserve constant O₂ levels, the medium was preequilibrated by exposure to the gas mixture for at least 24 hours. NSCs were used for experiments directly after isolation without cell culture passaging.

2.2. Isolation of RNA. Isolation of RNA was carried out using the RNeasy® Mini Kit (Qiagen, Hilden, Germany) according to manufacturer's instruction. To avoid adaption of the cells to the atmospheric O₂ level, the initiate steps were done quickly (within 6 min). RNA was eluted in RNase-free water by centrifugation, and RNA concentration was determined with the Quant-iT™ RNA Assay Kit (Life Technologies Corporation, Darmstadt, Germany) according to the manufacturer's instruction. For later analysis, RNA was stored at -80°C.

2.3. Quantitative RT-PCR. Quantitative RT-PCR was performed using the SYBR Green PCR Kit (Qiagen, Hilden, Germany) and the Stratagene Mx3000P thermocycler with the following program: 95°C for 15 min followed by 40 cycles of 94°C for 15 sec, 55°C for 30 sec, and 72°C for 30 sec. *Hydroxymethylbilane synthase (Hmbs)* was used as housekeeping genes for normalization. Initial studies on housekeeping gene expression in mNSCs over the cultivation time at the different oxygen levels revealed stable expression of *Hmbs* but not other housekeeping genes such as *Glycerinaldehyd-3-phosphat-dehydrogenase (Gapdh)* or *Tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (Ywhaz)*; Supplementary Figure S1). The following primers were used: *Hes5* F, 5'-GCTCGTAATC GCCTCCA-3'; *Hes5* R, 5'-GGTCCCACGCATCTTCT-3'; *Hif3a* F, 5'-CACTGCTCAGGACATATGAG-3'; *Hif3a* R, 5'-TCCAAAGCGTGGATGTATTC-3'; *Hmbs* F, 5'-TGTA TGCTGTGGGTCAGGGAG-3'; *Hmbs* R, 5'-CTCCTTCCA GGTGCCTCAGA-3'; *Igfbp4* F, 5'-GGTAGATCTTTGCT GTGGGAA-3'; *Igfbp4* R, 5'-CCTCAGACACACACAA TCCAA-3'; *Ldha* F, 5'-GGGTATCTCTGTGTAGCCC TGA-3'; *Ldha* R, 5'-AGAGGCAGGTGGATGTCTGT-3'; *Pdk1* F, 5'-GGTTGGGAACCACTCTTTCA-3'; *Pdk1* R, 5'-GCTTTGGTTACGTGGCATTT-3'; *Sfrp5* F, 5'-GATCTG TGCCAGTGTGAGA-3'; *Sfrp5* R, 5'-TTCAGCTGCC CATAGAAA-3'; *Slc2a1* F, 5'-CGACCCTCTTCTTTCA TCTCCT-3'; *Slc2a1* R, 5'-CAGGTCTCGGGTCACATCG-3'; *Slc2a3* F, 5'-GAACATTGGCTGGATACCTCTA-3'; *Slc2a3* R, 5'-ACAGTCACGGCGAACACC-3'; *Vegf* F, 5'-GCTACTGCCGTCGATTG-3'; and *Vegf* R, 5'-CTCCAG GGCTTCATCGTTAC-3' (all from Eurofins Genomics, Ebersberg, Germany).

2.4. Microarray Gene Chip Analysis. For whole microarray analyses, we used the microarray data from Braunschweig and coworkers [27]. As described in our previous article, mesencephalic and cortical NSCs were maintained in the expansion medium for 2 or 13 days under strict O₂ conditions using preequilibrated medium. After isolation of RNA from three independent cultures per condition and cell type, the concentration was measured using a Nanodrop spectrometer (Agilent Technologies, Santa Clara, CA, USA), and RNA quality was approved by an Agilent 2100 Bioanalyzer (all samples had a RNA integrity number > 9.0). Subsequent hybridization to whole mouse genome microarray gene expression chips (Gene ST 1.0 Arrays, Affymetrix, Santa Clara, CA, USA) was performed following the manufacturer's protocol. Microarray chips were then immediately scanned using the Hewlett-Packard Gene Array Scanner G2500A (Agilent Technologies, Santa Clara, CA, USA). Raw data were processed with the Exon Array Computational Tool (ExACT; Affymetrix) for background correction and normalization, and statistical data analysis was performed in the Agilent GeneSpring GX11 software. Statistical significance was computed using the unpaired Student's *t*-test ($P < 0.05$) with Benjamini-Hochberg as the multiple testing correction.

2.5. Gene Ontology Classification Analysis. The genes, which were up- or downregulated in the microarray analyses with a fold change of ≥ 2 , were analysed in a Gene Ontology analysis via the website "geneontology.org," which is using the PANTHER 17.0 Classification System software (<http://pantherdb.org/>). Of note, setting the fold change threshold to 1.5 led to very similar results. In this process, the regulated genes are classified into the different GO domains: biological process, molecular function, and cellular component, through manual and electronic annotations generated by a computer algorithm based on sequence similarities. The resulting genes are then allocated into different subgroups according to their biological context and occurrence. Based on the number of genes, the percentage of each subgroup in the entire GO domain can be determined. The resulting hits were composed of the diverse biological functions of a gene. For the analysis of the pathways, the PANTHER and Reactome databases were used. Reactome is a manually classified and peer-reviewed pathway database, but it also works in conjunction with PANTHER to identify orthologues of human proteins annotated in Reactome. Fisher's exact tests and Bonferroni correction were used in the PANTHER software to calculate fold enrichment and *P* values [38, 39].

3. Results

3.1. Differential Gene Regulation by Physioxia in Midbrain versus Cortical NSCs. To identify possible stem cell factors and molecular pathways differentially involved in the O₂ adaptation of fetal NSCs derived from midbrain and cortical tissues, Affymetrix microarray gene chip arrays were performed with NSCs cultured in physioxia 3% O₂ or in normoxic 21% O₂ for a short-term (2 days) or a long-term period (13 days). These whole-transcript assays provide a

complete expression profile of 28,853 coding transcripts in the mouse genome. To avoid overinterpretation of the data, the threshold for significance was set at a twofold or greater change in the expression level for all genes analysed (for complete lists of up- and downregulated genes, refer to Supplementary Tables S1–S6). Several of these genes were chosen to confirm fidelity of the microarray data by quantitative RT-PCR. Consistent results in the mode of regulation could be found in all cases proving reliability of the gene chip assays (Figure 1).

For both time points and also for both tissues, we detected more genes that were upregulated than downregulated in the physioxia compared to the normoxic condition (Table 1). None of the 28,853 investigated genes were downregulated in cells which were kept in physioxia for a 2-day period, whereas significant upregulation of expression could be observed for 9 genes in mesencephalic NSC and 34 genes in cortical NSC when kept at low O₂ levels for 2 days. Upon long-term culturing for 13 days, the number of significantly upregulated genes was 226 in midbrain and 96 in cortical NSCs, while 112 genes and 25 genes were downregulated, respectively.

Regarding short-term physioxia, the transcriptomes of midbrain and cortical NSCs showed a high number of identical genes compared with the total number of regulated genes, indicating strict concordance in gene expression. Here, 8 out of 9 more than twofold upregulated genes in the midbrain were also significantly upregulated in the cortex after short-term physioxia (Figure 2). However, long-term physioxia changed the gene expression profiles dramatically. There was an increase in affected genes in both cultures, with the number of matched genes being only 75 out of 338 genes in midbrain cells that were also affected in cortical cells. Interestingly, only 2 out of 9 genes that were induced by short-term physioxia in the midbrain were also significantly upregulated in long-term physioxia midbrain cells. These comparisons revealed that especially long-term physioxia influenced the gene expression pattern.

3.2. PANTHER Classification according to Biological Process of the Genes Regulated by Oxygen Levels in Midbrain versus Cortical NSCs. We then categorized all O₂-regulated genes using the PANTHER classification system (<http://pantherdb.org/>) according to the biological process (PANTHER GO-Slim Biological Process). Since individual genes have multiple functions and are listed in multiple groups, this total number is referred to as hits. The different classes are represented by the percentage of summed hits in the respective biological process category normalized to total hits. The individual hits of the categories represent the individual genes of the categories. The global comparison between short- and long-term cultures showed that their gene expression profiles differed over a wide range from the 28,853 genes analysed (Figures 3 and 4). In general, we observed a rise in the number of total categories as well as the sum of all hits, which indicates an increase in the varieties of biological processes.

In the case of short-term physioxia midbrain cultures, the analysed genes can be functionally classified into the




FIGURE 1: Comparison of gene expression levels measured with microarray gene chip assay and qRT-PCR showing good correlations between both measurements in mNSC. Log₂-transformed fold changes in mRNA levels (3% O₂/21% O₂) and standard error of mean ($n = 3$) were plotted for both array and qRT-PCR.

TABLE 1: Number of genes up- or downregulated more than twofolds in NSCs in physioxia.

	Midbrain NSCs (2 days in culture)	Cortical NSCs (2 days in culture)	Midbrain NSCs (13 days in culture)	Cortical NSCs (13 days in culture)
Upregulated in physioxia (3% O ₂)	9	34	226	96
Downregulated in physioxia (3% O ₂)	0	0	112	25

Displayed are the numbers of regulated genes for the various culture conditions and origins of NSCs. For complete gene lists, refer to Supplementary Tables S1–S6.




FIGURE 2: Comparison of differentially expressed genes in short- versus long-term physioxia of mesencephalic and cortical NSC cultures. The 4-way Venn diagram shows total numbers of O₂-regulated genes when comparing mesencephalic with cortical NSCs and short- (2 days) versus long-term O₂ stimulation (13 days).

two large categories: *metabolic processes* (27% of all hits, 6 hits) and *cellular processes* (23% of all hits, 6 hits) which are represented by genes such as *Pdk1* (pyruvate dehydrogenase kinase 1) or *Tpi1* (triose phosphate isomerase 1;

Figure 3). The number of hits increased from 31 to 801 (up- and downregulated) under long-term physioxia as compared to the normoxia condition leading to much bigger diversity of biological processes (increase in categories from 10 to 15). Despite various additional categories (*biomineralization*, *locomotion*, and *growth*), *cellular processes* (23%, 5 hits to 26%, 135 hits) and *biological regulation* (9%, 2 hits to 16%, 84 hits) even increased in the long-term hypoxic condition. On the other side, total hits in this category are increased, while the proportion of *metabolic processes* decreased from 27% (6 hits) to 15% (80 hits) at long term in midbrain NSC cultures.

Similar results were observed in short-term physioxia cortical NSCs, where 34 out of the 28,853 analysed genes are upregulated (Figure 4). The 63 attributed hits can be classified once more into *cellular processes* (32%, 20 hits) and *metabolic processes* (21%, 13 hits) representing the two largest categories as shown in short-term physioxia mid-brain cultures. In long-term cortical cultures, although not reaching the level of hits from midbrain results, the hits increased to 274 (up- and downregulated). Here, the expression patterns differed from midbrain cultures since both




FIGURE 3: Global comparison of PANTHER gene categories “biological process” in midbrain NSC cultures. Displayed are percentages of hits from oxygen-regulated genes to specific categories according to the PANTHER domain “biological process.” The PANTHER classification includes the total up-/downregulated genes (lower number in the inner circle) with at least a twofold difference for short- and long-term cultures. The grey circles indicate all analysed genes without a change of expression due to O₂ treatment. After 2 days of culture (2 d), no gene was downregulated in hypoxic cells, but 9 genes from different categories were upregulated, which formed a total of 31 hits. 3 genes and 14 hits of them were also upregulated in hypoxic prolonged cultures, while the other 6 genes did not undergo any O₂ change in prolonged cultures. In total, 226 genes were upregulated and formed 529 hits, and 112 genes with 272 hits were downregulated after 13 days of hypoxic culture (13 d). The hits result from the multiple biological functions of a gene. Since a gene can also have multiple functions, one gene also often results in multiple hits. Also shown are the percentages of individual hits in the various biological functions (categories) to the total hits of the regulated genes.

cellular processes (32%, 20 hits to 17%, 34 hits) and metabolic processes (21%, 20 hits to 16%, 34 hits) are reduced when compared to overall hits, and the distribution of categories is even more diverse than that in the midbrain. Next, there was a slight increase of hits from 13% (8 hits) in short-term to 16% (32 hits) in long-term physioxenic cortical cells in the category *biological regulation* and also an increase of the *response to stimulus* category (8%, 5 hits to 14%, 28 hits).

Regarding long-term downregulated genes and their hits, both cultures showed very similar patterns to their hits from upregulated genes: a large percentage of genes belonged to the biological process categories *biological regulation* (midbrain: 17%, 46 hits; cortex: 20%, 16 hits), *cellular processes* (midbrain: 25%, 67 hits; cortex: 23%, 18 hits), and *response to stimulus* (midbrain: 13%, 34 hits; cortex: 17%, 13 hits). Although the distribution of hits still showed a high regulation in the category *signaling* (midbrain: 13%, 34 hits; cortex: 17%, 13 hits), the category was particularly prominent in cortical NSCs. The *signaling* category is represented by genes like *Hey2* (hairly/enhancer-of-split related with YRPW motif protein 2) and *Dll1* (delta-like protein 1) both involved in *Notch signaling* and thus playing a crucial role in the development of the nervous system [40, 41].

In summary, more genes in midbrain and cortex NSCs were regulated by a prolonged physioxenic environment as compared with short-term physioxenia. Most prominently, there was a reduction of hits in *metabolic processes* but an increase in *biological regulation* and *signaling* when com-

pared to overall hits. Despite many differentially regulated genes, the differences in categories and their distribution between midbrain and cortical cultures were small.

3.3. GO Classification of the Genes Regulated by Oxygen Levels in Midbrain versus Cortical NSCs. We then analysed the regulated genes for their classification into the GO domains of biological processes, molecular functions, and cellular components using GO analysis (“geneontology.org”), which also uses the PANTHER database. Overall, detailed analyses of biological processes, molecular functions, and cellular components at short-term physioxenia revealed primarily a link to *glycolysis*, *pyruvate*, and *general carbohydrate metabolism* regardless of the tissue origin (Supplementary Table S7). Genes specifically responsible for the cellular response to low oxygen conditions were also upregulated in both cultures. In this context, the upregulated genes also showed effects on peptidyl proline 4-dioxygenase (or prolyl 4-hydroxylase), which is generally required for the hydroxylation of prolyl residues in proteins such as collagen and, in turn, for the stabilization of hypoxia-inducible factor-1 (HIF-1) [42–44]. Its activity is controlled by several genes, such as *Egln3* (prolyl hydroxylase) and *P4ha1* (prolyl 4-hydroxylase subunit alpha-1), which were also upregulated in both cultures after short-term physioxenia.

The analysis of long-term cultures showed that the fold enrichment of the individual biological processes, cellular




FIGURE 4: Global comparison of PANTHER gene categories “biological process” in cortical NSC cultures. Displayed are percentages of hits from oxygen-regulated genes to specific categories according to the PANTHER domain “biological process.” The PANTHER classification includes the total up-/downregulated genes (lower number in the inner circle) with at least a twofold difference for short- and long-term cultures. The grey circles indicate all analysed genes without a change of expression due to O₂ treatment. After 2 days of culture (2 d), no gene was downregulated in hypoxic cells, but 9 genes from different categories were upregulated, which formed a total of 31 hits. 3 genes and 14 hits of them were also upregulated in hypoxic prolonged cultures, while the other 6 genes did not undergo any O₂ change in prolonged cultures. In total, 226 genes were upregulated and formed 529 hits, and 112 genes with 272 hits were downregulated after 13 days of hypoxic culture (13 d). The hits result from the multiple biological functions of a gene. Since a gene can also have multiple functions, one gene also often results in multiple hits. Also shown are the percentages of individual hits in different biological functions (categories) to the total hits of the regulated genes.

functions, and components was rather low compared to the results in short-term physioxia (Supplementary Tables S8 and S9). On closer inspection, the cellular process *protein hydroxylation* seems to be very present in the upregulated genes under long-term physioxia in a tissue-independent manner (fold enrichment of at least 20.1; Supplementary Table S8). Both NSC types reflected genes associated with the *binding of collagen* as the molecular function, which also played a role in the cellular components of the basement membrane (“*collagen-containing extracellular matrix*”). In contrast, only midbrain NSC cultures had regulated gene functions in angiogenesis and neuron generation after prolonged exposure to physioxia. Interestingly, in the midbrain, genes that positively influence *Notch signaling* and *neurogenesis* appear to be downregulated (Supplementary Table S9).

3.4. GO Cellular Pathway Analyses of the Genes Regulated by Oxygen Levels in Midbrain versus Cortical NSCs. We then categorized the regulated genes in relation to cellular metabolic pathways using GO (Gene Ontology) analysis via geneontology.org using the two types of software and databases PANTHER and Reactome. Analysis of the 9 upregulated genes in short-term midbrain NSC cultures revealed that two of these genes with a fold enrichment greater than 100 were associated with a glycolysis pathway, and 3 genes were associated with a pyruvate pathway, which is required, for example, for the functioning of the citrate acid cycle (Table 2). Cortical cultures also revealed pathway affiliation

with *glycolysis*. In contrast to midbrain cultures, cortical cells showed no affiliation in *pyruvate metabolism* but an additional effect regarding *collagen formation*.

In long-term physioxia NSC cultures, it was generally noticed that the fold enrichment of the cellular pathways was comparatively low compared to the results in short-term physioxia (Tables 3 and 4). The pathway affiliations of midbrain cells showed relations to *Posttranslational protein phosphorylation* and *Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)*. Both pathways could not be found in GO pathway analyses from affected genes in cortical NSCs but showed a link to *amino acid transport across the plasma membrane* with a fold enrichment of 28.63.

Pathway analysis of the downregulated genes in long-term physioxia revealed an association with *Notch signaling* in midbrain (fold enrichment: 18.7) as well as cortical NSCs (fold enrichment: 62.8; Table 4). Once more, there are differences between both cultures since affected genes of midbrain cells are associated with *Wnt signaling: Frzb* (secreted frizzled-related protein 3), *Sfrp5* (secreted frizzled-related protein 5), and *Nkd1* (protein naked cuticle homolog 1). In contrast, affected genes of cortical NSCs are associated with *transmission across chemical synapses* demonstrating once more a tissue-dependent reaction to long-term physioxia.

3.5. Differential Regulation of Differentiation Marker Genes by Physioxia in Midbrain NSC. To further characterize the cNSC and mNSC cultured in 3% O₂ and used for microarray

TABLE 2: Affected pathways from upregulated genes in midbrain and cortex NSCs cultivated for 2 days in physioxia.

	Midbrain NSCs		P value	Number of genes	Number of genes	Cortical NSCs	
	Number of genes	Fold enrichment				Fold enrichment	P value
PANTHER pathways <i>Mus musculus</i> (reference)	2	>100	8.10E-03	—	PANTHER pathways <i>Mus musculus</i> (reference)	4	43.11
Glycolysis					—		
Reactome pathways <i>Mus musculus</i> (reference)	3	>100	3.41E-04		Reactome pathways <i>Mus musculus</i> (reference)	4	32.34
					Glycolysis		1.35E-02
					(i) Glucose metabolism	6	14.59
					(ii) metabolism of carbohydrates	4	5.45E-03
					Collagen formation	4	1.94E-02

TABLE 3: Affected pathways from upregulated genes in midbrain and cortex NSCs cultivated for 13 days in physioxia.

	Midbrain NSCs		Cortical NSCs	
	Number of genes	Fold enrichment	Number of genes	Fold enrichment
		P value		P value
PANTHER pathways <i>Mus musculus</i> (reference)				
—			Amino acid transport across the plasma membrane	28.63
Reactome pathways <i>Mus musculus</i> (reference)			Reactome pathways <i>Mus musculus</i> (reference)	2.86E-02
Regulation of IGF transport and uptake by IGF binding proteins	10	8.25	—	
Post-translational protein phosphorylation	9	7.82	—	




FIGURE 5: Comparison of differentiation-related genes in short- versus long-term physioxia of mesencephalic and cortical NSC cultures. The heat map shows slight changes in fate decisions solely in long-term physioxia mNSC (after 13 days). Fold changes of differentiation-related genes are displayed using colour coding (red: downregulated; green: upregulated). ** $P < 0.01$, t -test with Benjamini-Hochberg procedure for multiple testing correction.

analysis, we displayed a set of specific genes corresponding to neuronal differentiation (Figure 5). The analysis of these markers revealed that lowered oxygen levels did not affect differentiation markers in short-term cultures. On the other hand, there are changes in long-term cultures, which reaches significance solely in mNSCs: two of these markers—*Gfap* and *Map2*—were downregulated, which is comparable with our previous immunohistochemical data [27]. *Ngn2* as a prominent marker for differentiation into various neuronal subtypes was also downregulated [45]. Intriguingly, our data suggested that lowered oxygen levels promote a slight transition of mNSC towards oligodendrocyte precursor cells (OPCs) indicated by significant upregulation of the OPC proliferation regulator *Ascl1* and the OPC-specific marker *Ng2*, while the mature OPC marker *Olig2* is not affected while the radial glia marker *Pax6* is downregulated [46, 47].

4. Discussion

Global comparison of gene expression profiles of short-term (2 days) and long-term (13 days) NSC cultures derived from fetal midbrain or cortical tissue showed that initial adaptation to O_2 deficiency is distinct from maintenance of the response to physioxia culture conditions as suggested in previous reports [48, 49].

In the short-term (2 days) cultures of midbrain and cortical NSCs, very few genes exhibited a twofold or greater change in gene expression upon O_2 alteration, relative to the 28,853 genes examined. There was a high concordance of individual genes in both NSC types. The O_2 -regulated genes mainly involve glycolysis and contribute to the current knowledge that cells reduce glucose production in the presence of decreasing O_2 through a three-step switch from oxidative to glycolytic metabolism to maintain ATP levels [50–52]. The first step involves activated transcription of glucose transporters and glycolytic enzymes to increase glycolytic flux from glucose to pyruvate. Second, the expression of pyruvate dehydrogenase kinase 1 (*Pdk1*) is induced to prevent flux through the tricarboxylic acid cycle and decrease the production of reactive oxygen species. And third, transcription of lactate dehydrogenase A (*Ldha*) is induced to activate the conversion of pyruvate to lactate. Therefore, the switch is known to be regulated via hypoxia-inducible factors (HIFs) which are key regulators for low oxygen conditions [53–55]. Consequently, whole-transcriptome analyses showed that relevant genes such as *Pdk1* and the glucose transporter *Slc2a1* (solute

carrier family 2, member 1) were upregulated, leading to altered biological processes such as pyruvate and glucose metabolism, but also the cellular response to physioxia from gene ontology analysis. These results, consistent with the literature, suggest a largely tissue-independent regulation of cells to short-term hypoxia/physioxia.

The direct response to hypoxia/physioxia of both NSC cultures is also evident in the upregulation of the molecular function peptidyl proline 4-dioxygenase activity involving the *P4ha1* and *Egln3* (*Phd3*) genes. *P4ha1*, only regulated in short-term hypoxia, is known to induce HIF-1 signaling in breast cancer cells through hypoxia [44]. *Phd3* encodes an intracellular prolyl hydroxylase that, distinct from *Phd1* or *Phd2*, primarily hydroxylates the alpha subunit of hypoxia-inducing factor 2 (HIF-2a) acting as a putative feedback control or fast acting regulator in the case of reoxygenation [56–61]. In such normoxic conditions, HIF proteins are continuously hydroxylated by HIF-prolyl hydroxylases (PHD1-3) and degraded by the vHL protein (Von Hippel-Lindau protein). Since these hydroxylation reactions require oxygen, the degradation of HIFs decreases at lower oxygen tension, allowing them to transcribe target genes as transcription factors [62, 63]. Of note, the *Phd3* mRNA level is elevated independent of tissue or physioxia duration, although there are dramatic changes in long-term physioxia gene expression. Thus, our findings likely contribute to the knowledge that it acts as a fast regulator for the case of reoxygenation [64].

While the cellular response to short-term physioxia seems to be related to immediate metabolic changes for cell survival, long-term physioxia (13 days) cultures showed a fundamentally different scenario: from the perspective of chronic physioxia, it may become more important for the cells to prepare for a prolonged absence of O_2 than attempting to further improve supply [26, 32, 65]. Consecutively, the concordance of genes between short-term and long-term physioxia within the same culture (mesencephalon or cortex) was less than 33%, and hypoxia response genes such as *Vegf*, glucose transporter, or *Pdk1* are less or absent in NSCs cultured in long-term physioxia. This results in a switch from affected metabolic pathways in short-term physioxia cultures to signaling pathways such as Notch signaling or Wnt signaling, which is consistent with human studies showing that the response to short-term hypoxia/physioxia differs substantially from adaptation to long-term hypoxia/physioxia [48, 49]. Intriguingly, the affected signaling pathways Notch and Wnt (only in midbrain NSCs) are key

players within embryonic brain development *in vivo* [66, 67]. Associated genes such as *Dll3*, *Hes5*, *Frzb*, or *Cdh8* are all known to be important for adequate brain development and support the relevance of so-called hypoxic niches where neural stem and progenitor cells reside *in vivo* [68, 69]. *Dll1*, *Dll3*, or *Hes5* as part of Notch signaling and downregulated in long-term physioxic midbrain NSC is known to be essential for neurogenesis and is predominantly described for maintenance of stem cell state and blocking of differentiation [70–77]. In long-term physioxic cortical NSCs, similar results were found regarding Notch signaling by downregulation of *Dll1* or *Hey2* [78, 79]. Although most of these studies are only relevant for the cortical situation, they are conflicting with enhanced proliferation under hypoxia from previous reports [27]. This may be explained by the oscillation of Notch genes such as *Dll1* or *Hes1* whose sustained expression would lead to inhibition of proliferation [80, 81].

In long-term physioxic NSC cultures from the midbrain, Wnt signaling is also affected through downregulated genes such as *Celsr1*, *Sfrp5*, *Frzb*, or *Cdh8*. Here, for example, *Celsr1*, *Frzb* (also known as *Sfrp3*) and *Sfrp5* (or its homologs *Sfrp1/2*) functional studies revealed that their inhibition is linked to enhanced stem cell maintenance (Boucherie [82–85]. Vice versa, we could detect upregulated proliferation-enhancing genes of the Wnt pathway like *Ccnd2* known to be expressed during midbrain development [86–88]. We thus conclude that known proliferative effects on midbrain NSC through physioxia might be mediated through Wnt pathways and differ compared to cortical cells. These results are consistent with our previous functional results [16, 27]. Additionally, our analysis of differentiation markers revealed a reduction of *Map2* expression in physioxia, which is consistent with our previous results [27]. This effect is solely detectable in mNSCs, which—in contrast to cNSCs—show nearly no spontaneous differentiation towards *Map2*⁺ cells after cultivation in physioxia [27]. Interestingly, we also observed a slight shift of OPC marker expression towards immature OPCs.

Besides Wnt, we could detect other specific pathways such as *Posttranslational protein phosphorylation* and *Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)* for midbrain NSCs. As an example, *Igfbp4* is normally expressed during rodent brain development but starts to significantly decrease with E14.5 *in vivo* [89]. At this stage, brain vascularization—beginning with its first sprouting at roughly E9—is established, and oxygen can be easily delivered, which may act as a signaling factor during development [36, 90, 91]. Additionally, the upregulated genes *Igf2* and *Igfbp3* are linked to human brain development while *Igfbp3* in particular is able to regulate cell growth at least [92, 93]. Of note, despite GO not revealing this pathway in cortical NSCs, its counterpart *Igfbp4* is also regulated in cortical cells.

Moreno and colleagues found calcineurin-NFATc4 signaling in hypoxic mouse NSCs as an additional signaling pathway acting as a major regulator of self-renewal and proliferation [32]. We have no indication of an importance of this transcription factor pathway in our NSC cultures (fold change of *Nfatc4* in mNSC after 48 h: 1.00; mNSC, 13 d:

1.01; cNSC, 48 h: 1.09; cNSC, 13 d: 1.00). This discrepancy might be due to differences in the experimental paradigm: we here used primarily freshly isolated NSC at 3% for physioxia, while Moreno and coworkers cultured their NSCs at 5% oxygen tension over several cell culture passages. While it is well known that passaging can provoke changes regarding the cellular properties, Horie and coworkers showed—at least for NSC from the ganglionic eminence—that there is a critical shift of NSC behavior at around 4% O₂ which might explain the mentioned differences [94, 95]. Moreover, Moreno and colleagues performed a gene set enrichment analysis (GSEA) using transcription factor target (TFT) and KEGG gene sets. Unfortunately, both gene sets are currently not available for the murine situation in the GSEA software limiting a direct comparison of our data.

Cortical NSCs remarkably show affected pathways regarding *amino acid transport across the plasma membrane* and *transmission across chemical synapses*. Our previous results show that there are only sporadic differentiated cells in long-term physioxic cortical cells [27], so we assume that the regulation of chemical synapses corresponds to signaling regulating stem cell properties rather than its classical role in neuronal functionality. In particular, gamma-aminobutyric acid (GABA) signaling is reported to play a distinct role during brain—particularly cortical—development [96–99]. In agreement with the literature [32, 99], we detected the expression of various GABA receptor subunits with downregulation in long-term physioxia of *Gabbr1* (subunit 1 of the GABA_B receptor) and *Gabrg2* (subunit γ 2 of the GABA_A receptor) in cortical but only *Gabbr2* (subunit 2 of the GABA_B receptor) in midbrain NSCs. In contrast, *Gabra2* (subunit α 2 of the GABA_A receptor) was upregulated in cortical NSC in physioxic conditions. Interestingly, both GABA receptor subtypes have been reported to regulate NSC proliferation [96, 97]. However, there are conflicting results by showing that GABA could also express inhibitory effects on proliferation of cortical progenitors [98]. The effects of GABA on NSC proliferation, maintenance, and differentiation might thus depend on the exact developmental stage of the NSCs and their microenvironment [97, 99].

5. Conclusion

This work focuses on changes in gene expression in midbrain and cortical NSCs that occur under low O₂ culture conditions reflecting physiological oxygen tension and which are beneficial expansion conditions for fetal NSCs as compared to atmospheric O₂. The study shows specific effects of oxygen on NSCs during short-term cultivation mostly affecting metabolic processes which dramatically shifts to signaling processes and stem cell maintenance during long-term cultivation. Strikingly, even though short-term effects are very similar in both NSC types, midbrain and cortical NSCs react differently during long-term cultivation in their physioxic environment underpinning the discrepancies in their oxygen response [16, 18, 19, 28]. These results reveal the extent of molecular mechanisms during O₂ modulation and help to identify pathways that control mammalian stem cell behavior in hypoxic/physioxic stem cell niches.

Data Availability

All data including microarray data can be obtained by contacting the corresponding author without any restrictions.

Conflicts of Interest

The authors do not report any competing financial and nonfinancial interests.

Authors' Contributions

LB, AKM, and AS conceived and designed the experiments. LB did most of the experiments. LB, AKM, FM, JL, GF, and AS analysed, interpreted, and assembled the data. LB, AKM, JL, FM, and AS wrote the paper. All authors read and approved the final manuscript.

Acknowledgments

We are very grateful to Georg Fuellen from the Department of Biostatistics and Informatics in Medicine and Aging Research (IBIMA), Rostock, Germany, for the many fruitful discussions. We would also like to thank Sylvia Kanzler, Andrea Kempe, and Cornelia Mai for their excellent technical assistance. The work was supported by the Deutsche Forschungsgemeinschaft through the Collaborative Research Center 655 "Cells into Tissues: Stem Cell and Progenitor Commitment and Interactions during Tissue Formation" (SFB655, project A23) to AS and the Dresden International Graduate School for Biomedicine and Bioengineering (DIGS-BB) through the DIGS-BB fellow award (to LB).

Supplementary Materials

Supplementary Figure S1: comparison of cycle threshold (CT) values of various housekeeping genes as measured by qRT-PCR in dependency on cultivation times and oxygen levels in mesencephalic NSCs. Supplementary Table S1: upregulated genes in cortical NSCs cultivated for 2 days in physioxia compared to normoxia. Supplementary Table S2: upregulated genes in cortical NSCs cultivated for 13 days in physioxia compared to normoxia. Supplementary Table S3: downregulated genes in cortical NSCs cultivated for 13 days in physioxia compared to normoxia. Supplementary Table S4: upregulated genes in midbrain NSCs cultivated for 2 days in physioxia compared to normoxia. Supplementary Table S5: upregulated genes in midbrain NSCs cultivated for 13 days in physioxia compared to normoxia. Supplementary Table S6: downregulated genes in midbrain NSCs cultivated for 13 days in physioxia compared to normoxia. Supplementary Table S7: affected biological processes, molecular functions, and components from upregulated genes in midbrain and cortical NSCs cultivated for 2 days in physioxia. Supplementary Table S8: affected biological processes, molecular functions, and components from upregulated genes in midbrain and cortical NSCs cultivated for 13 days in physioxia. Supplementary Table S9: affected biological processes, molecular functions, and components from downregulated genes in midbrain and cortical NSCs cultivated for 13 days in physioxia. (*Supplementary Materials*)

References

- [1] A. A. Davis and S. Temple, "A self-renewing multipotential stem cell in embryonic rat cerebral cortex," *Nature*, vol. 372, no. 6503, pp. 263–266, 1994.
- [2] F. H. Gage, "Mammalian neural stem cells," *Science*, vol. 287, no. 5457, pp. 1433–1438, 2000.
- [3] A. Bjorklund and M. Parmar, "Dopamine cell therapy: from cell replacement to circuitry repair," *Journal of Parkinson's Disease*, vol. 11, no. s2, pp. S159–S165, 2021.
- [4] J. Fu, A. Warmflash, and M. P. Lutolf, "Stem-cell-based embryo models for fundamental research and translation," *Nature Materials*, vol. 20, no. 2, pp. 132–144, 2021.
- [5] J. H. Lee and W. Sun, "Neural organoids, a versatile model for neuroscience," *Molecules and Cells*, vol. 45, no. 2, pp. 53–64, 2022.
- [6] B. Martinez and P. V. Peplow, "Biomaterial and tissue-engineering strategies for the treatment of brain neurodegeneration," *Neural Regeneration Research*, vol. 17, no. 10, pp. 2108–2116, 2022.
- [7] R. S. Rathore, R. Ayyannan, and S. K. Mahto, "Emerging three-dimensional neuronal culture assays for neurotherapeutics drug discovery," *Expert Opinion on Drug Discovery*, vol. 17, no. 6, pp. 619–628, 2022.
- [8] L. H. Thompson and A. Bjorklund, "Reconstruction of brain circuitry by neural transplants generated from pluripotent stem cells," *Neurobiology of Disease*, vol. 79, pp. 28–40, 2015.
- [9] J. P. Andreotti, W. N. Silva, A. C. Costa et al., "Neural stem cell niche heterogeneity," *Seminars in Cell & Developmental Biology*, vol. 95, pp. 42–53, 2019.
- [10] N. Genet and K. K. Hirschi, "Understanding neural stem cell regulation in vivo and applying the insights to cell therapy for strokes," *Regenerative Medicine*, vol. 16, no. 9, pp. 861–870, 2021.
- [11] E. A. B. Gilbert, N. Lakshman, K. S. K. Lau, and C. M. Morshead, "Regulating endogenous neural stem cell activation to promote spinal cord injury repair," *Cell*, vol. 111, no. 5, 2022.
- [12] K. Ioannidis, I. Angelopoulos, G. Gakis et al., "3D reconstitution of the neural stem cell niche: connecting the dots," *Frontiers in Bioengineering and Biotechnology*, vol. 9, article 705470, 2021.
- [13] D. Jain, S. Mattiassi, E. L. Goh, and E. K. F. Yim, "Extracellular matrix and biomimetic engineering microenvironment for neuronal differentiation," *Neural Regeneration Research*, vol. 15, no. 4, pp. 573–585, 2020.
- [14] G. Li, J. Liu, Y. Guan, and X. Ji, "The role of hypoxia in stem cell regulation of the central nervous system: from embryonic development to adult proliferation," *CNS Neuroscience & Therapeutics*, vol. 27, no. 12, pp. 1446–1457, 2021.
- [15] C. M. Willis, A. M. Nicaise, G. Krzak et al., "Soluble factors influencing the neural stem cell niche in brain physiology, inflammation, and aging," *Experimental Neurology*, vol. 355, article 114124, 2022.
- [16] J. Milosevic, S. C. Schwarz, K. Krohn, M. Poppe, A. Storch, and J. Schwarz, "Low atmospheric oxygen avoids maturation, senescence and cell death of murine mesencephalic neural precursors," *Journal of Neurochemistry*, vol. 92, no. 4, pp. 718–729, 2005.
- [17] F. Pistollato, H. L. Chen, P. H. Schwartz, G. Basso, and D. M. Panchision, "Oxygen tension controls the expansion of human

- CNS precursors and the generation of astrocytes and oligodendrocytes,” *Molecular and Cellular Neurosciences*, vol. 35, no. 3, pp. 424–435, 2007.
- [18] A. Storch, G. Paul, M. Csete et al., “Long-term proliferation and dopaminergic differentiation of human mesencephalic neural precursor cells,” *Experimental Neurology*, vol. 170, no. 2, pp. 317–325, 2001.
- [19] L. Studer, M. Csete, S. H. Lee et al., “Enhanced proliferation, survival, and dopaminergic differentiation of CNS precursors in lowered oxygen,” *The Journal of Neuroscience*, vol. 20, no. 19, pp. 7377–7383, 2000.
- [20] L. Wagenfuhr, A. K. Meyer, L. Braunschweig, L. Marrone, and A. Storch, “Brain oxygen tension controls the expansion of outer subventricular zone-like basal progenitors in the developing mouse brain,” *Development*, vol. 142, no. 17, pp. 2904–2915, 2015.
- [21] L. Wagenfuhr, A. K. Meyer, L. Marrone, and A. Storch, “Oxygen tension within the neurogenic niche regulates dopaminergic neurogenesis in the developing midbrain,” *Stem Cells and Development*, vol. 25, no. 3, pp. 227–238, 2016.
- [22] A. Carreau, B. El Hafny-Rabbi, A. Matejuk, C. Grillon, and C. Kieda, “Why is the partial oxygen pressure of human tissues a crucial parameter? Small molecules and hypoxia,” *Journal of Cellular and Molecular Medicine*, vol. 15, no. 6, pp. 1239–1253, 2011.
- [23] I. Silver and M. Erecińska, “Oxygen and ion concentrations in normoxic and hypoxic brain cells,” *Advances in Experimental Medicine and Biology*, vol. 454, pp. 7–16, 1998.
- [24] P. Jezek, L. Plecita-Hlavata, K. Smolkova, and R. Rossignol, “Distinctions and similarities of cell bioenergetics and the role of mitochondria in hypoxia, cancer, and embryonic development,” *The International Journal of Biochemistry & Cell Biology*, vol. 42, no. 5, pp. 604–622, 2010.
- [25] D. M. Panchision, “The role of oxygen in regulating neural stem cells in development and disease,” *Journal of Cellular Physiology*, vol. 220, no. 3, pp. 562–568, 2009.
- [26] F. R. Sharp and M. Bernaudin, “HIF1 and oxygen sensing in the brain,” *Nature Reviews. Neuroscience*, vol. 5, no. 6, pp. 437–448, 2004.
- [27] L. Braunschweig, A. K. Meyer, L. Wagenfuhr, and A. Storch, “Oxygen regulates proliferation of neural stem cells through Wnt/ β -catenin signalling,” *Molecular and Cellular Neurosciences*, vol. 67, pp. 84–92, 2015.
- [28] J. Milosevic, M. Maisel, F. Wegner et al., “Lack of hypoxia-inducible factor-1 alpha impairs midbrain neural precursor cells involving vascular endothelial growth factor signaling,” *The Journal of Neuroscience*, vol. 27, no. 2, pp. 412–421, 2007.
- [29] C. Krabbe, S. T. Bak, P. Jensen et al., “Influence of oxygen tension on dopaminergic differentiation of human fetal stem cells of midbrain and forebrain origin,” *PLoS One*, vol. 9, no. 5, p. e96465, 2014.
- [30] H. Felfly, A. C. Zambon, J. Xue et al., “Severe hypoxia: consequences to neural stem cells and neurons,” *Journal of Neurology Research*, vol. 1, no. 5, 2011.
- [31] N. Horie, K. So, T. Moriya et al., “Effects of oxygen concentration on the proliferation and differentiation of mouse neural stem cells in vitro,” *Cellular and Molecular Neurobiology*, vol. 28, no. 6, pp. 833–845, 2008.
- [32] M. Moreno, V. Fernandez, J. M. Monllau, V. Borrell, C. Lerin, and N. de la Iglesia, “Transcriptional profiling of hypoxic neural stem cells identifies calcineurin-NFATc4 signaling as a major regulator of neural stem cell biology,” *Stem Cell Reports*, vol. 5, no. 2, pp. 157–165, 2015.
- [33] Y. Liu, S. R. Cox, T. Morita, and S. Kourembanas, “Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells,” *Circulation Research*, vol. 77, no. 3, pp. 638–643, 1995.
- [34] G. L. Semenza, P. H. Roth, H. M. Fang, and G. L. Wang, “Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1,” *The Journal of Biological Chemistry*, vol. 269, no. 38, pp. 23757–23763, 1994.
- [35] G. L. Semenza and G. L. Wang, “A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation,” *Molecular and Cellular Biology*, vol. 12, no. 12, pp. 5447–5454, 1992.
- [36] C. Lange, M. Turrero Garcia, I. Decimo et al., “Relief of hypoxia by angiogenesis promotes neural stem cell differentiation by targeting glycolysis,” *The EMBO Journal*, vol. 35, no. 9, pp. 924–941, 2016.
- [37] S. Tomita, M. Ueno, M. Sakamoto et al., “Defective brain development in mice lacking the Hif-1alpha gene in neural cells,” *Molecular and Cellular Biology*, vol. 23, no. 19, pp. 6739–6749, 2003.
- [38] H. Mi, A. Muruganujan, D. Ebert, X. Huang, and P. D. Thomas, “PANTHER version 14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools,” *Nucleic Acids Research*, vol. 47, no. D1, pp. D419–D426, 2019.
- [39] The Gene Ontology Consortium, “Expansion of the Gene Ontology knowledgebase and resources,” *Nucleic Acids Research*, vol. 45, no. D1, pp. D331–D338, 2017.
- [40] A. Fischer, N. Schumacher, M. Maier, M. Sendtner, and M. Gessler, “The Notch target genes Hey1 and Hey2 are required for embryonic vascular development,” *Genes & Development*, vol. 18, no. 8, pp. 901–911, 2004.
- [41] V. K. Jordan, J. A. Rosenfeld, S. R. Lalani, and D. A. Scott, “Duplication of HEY2 in cardiac and neurologic development,” *American Journal of Medical Genetics. Part A*, vol. 167-A, no. 9, pp. 2145–2149, 2015.
- [42] C. Grimmer, N. Balbus, U. Lang et al., “Regulation of type II collagen synthesis during osteoarthritis by prolyl-4-hydroxylases: possible influence of low oxygen levels,” *The American Journal of Pathology*, vol. 169, no. 2, pp. 491–502, 2006.
- [43] K. Ullah, A.-H. Rosendahl, V. Izzi et al., “Hypoxia-inducible factor prolyl-4-hydroxylase-1 is a convergent point in the reciprocal negative regulation of NF- κ B and p53 signaling pathways,” *Scientific Reports*, vol. 7, no. 1, pp. 1–18, 2017.
- [44] G. Xiong, R. L. Stewart, J. Chen et al., “Collagen prolyl 4-hydroxylase 1 is essential for HIF-1 α stabilization and TNBC chemoresistance,” *Nature Communications*, vol. 9, no. 1, pp. 1–16, 2018.
- [45] E. C. Thoma, E. Wischmeyer, N. Offen et al., “Ectopic expression of neurogenin 2 alone is sufficient to induce differentiation of embryonic stem cells into mature neurons,” *PLoS One*, vol. 7, no. 6, article e38651, 2012.
- [46] K. C. Allan, L. R. Hu, M. A. Scavuzzo et al., “Non-canonical targets of HIF1a impair oligodendrocyte progenitor cell function,” *Cell Stem Cell*, vol. 28, no. 2, pp. 257–272.e11, 2021.
- [47] D. P. Kelenis, E. Hart, M. Edwards-Fligner, J. E. Johnson, and T. Y. Vue, “ASCL1 regulates proliferation of NG2-glia in the

- embryonic and adult spinal cord,” *Glia*, vol. 66, no. 9, pp. 1862–1880, 2018.
- [48] C. M. Cameron, F. Harding, H. Wei-Shou, and D. S. Kaufman, “Activation of hypoxic response in human embryonic stem cell-derived embryoid bodies,” *Experimental Biology and Medicine*, vol. 233, no. 8, pp. 1044–1057, 2008.
- [49] C. E. Forristal, K. L. Wright, N. A. Hanley, R. O. C. Oreffo, and F. D. Houghton, “Hypoxia inducible factors regulate pluripotency and proliferation in human embryonic stem cells cultured at reduced oxygen tensions,” *Reproduction (Cambridge, England)*, vol. 139, no. 1, pp. 85–97, 2010.
- [50] T. Kashiwagura, D. F. Wilson, and M. Erecińska, “Oxygen dependence of cellular metabolism: the effect of O₂ tension on gluconeogenesis and urea synthesis in isolated rat hepatocytes,” *Journal of Cellular Physiology*, vol. 120, no. 1, pp. 13–18, 1984.
- [51] J.-w. Kim, I. Tchernyshyov, G. L. Semenza, and C. V. Dang, “HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia,” *Cell Metabolism*, vol. 3, no. 3, pp. 177–185, 2006.
- [52] C. Maffezzini, J. Calvo-Garrido, A. Wredenberg, and C. Freyer, “Metabolic regulation of neurodifferentiation in the adult brain,” *Cellular and Molecular Life Sciences: CMLS*, vol. 77, no. 13, pp. 2483–2496, 2020.
- [53] W. G. Kaelin and P. J. Ratcliffe, “Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway,” *Molecular Cell*, vol. 30, no. 4, pp. 393–402, 2008.
- [54] P. Vaupel and G. Multhoff, “Revisiting the Warburg effect: historical dogma versus current understanding,” *The Journal of Physiology*, vol. 599, no. 6, pp. 1745–1757, 2021.
- [55] H. Zhang, P. Gao, R. Fukuda et al., “HIF-1 inhibits mitochondrial biogenesis and cellular respiration in VHL-deficient renal cell carcinoma by repression of C-MYC activity,” *Cancer Cell*, vol. 11, no. 5, pp. 407–420, 2007.
- [56] R. J. Appelhoff, Y.-M. Tian, R. R. Raval et al., “Differential function of the prolyl hydroxylases PHD1, PHD2, and PHD3 in the regulation of hypoxia-inducible factor,” *The Journal of Biological Chemistry*, vol. 279, no. 37, pp. 38458–38465, 2004.
- [57] O. Aprelikova, G. V. R. Chandramouli, M. Wood et al., “Regulation of HIF prolyl hydroxylases by hypoxia-inducible factors,” *Journal of Cellular Biochemistry*, vol. 92, no. 3, pp. 491–501, 2004.
- [58] E. Berra, E. Benizri, A. Ginouvès, V. Volmat, D. Roux, and J. Pouyssegur, “HIF prolyl-hydroxylase 2 is the key oxygen sensor setting low steady-state levels of HIF-1alpha in normoxia,” *The EMBO Journal*, vol. 22, no. 16, pp. 4082–4090, 2003.
- [59] J. Fu, K. Menzies, R. S. Freeman, and M. B. Taubman, “EGLN3 prolyl hydroxylase regulates skeletal muscle differentiation and myogenin protein stability,” *The Journal of Biological Chemistry*, vol. 282, no. 17, pp. 12410–12418, 2007.
- [60] J. H. Marxsen, P. Stengel, K. Doege et al., “Hypoxia-inducible factor-1 (HIF-1) promotes its degradation by induction of HIF-alpha-prolyl-4-hydroxylases,” *The Biochemical Journal*, vol. 381, Part 3, pp. 761–767, 2004.
- [61] P. Miikkulainen, H. Högel, F. Seyednasrollah, K. Rantanen, L. L. Elo, and P. M. Jaakkola, “Hypoxia-inducible factor (HIF)-prolyl hydroxylase 3 (PHD3) maintains high HIF2A mRNA levels in clear cell renal cell carcinoma,” *The Journal of Biological Chemistry*, vol. 294, no. 10, pp. 3760–3771, 2019.
- [62] L. del Peso, M. C. Castellanos, E. Temes et al., “The von Hippel Lindau/hypoxia-inducible factor (HIF) pathway regulates the transcription of the HIF-proline hydroxylase genes in response to low oxygen,” *The Journal of Biological Chemistry*, vol. 278, no. 49, pp. 48690–48695, 2003.
- [63] G. L. Semenza, “Hypoxia-inducible factor 1: control of oxygen homeostasis in health and disease,” *Pediatric Research*, vol. 49, no. 5, pp. 614–617, 2001.
- [64] E. Moroz, S. Carlin, K. Dyomina et al., “Real-time imaging of HIF-1 α stabilization and degradation,” *PLoS One*, vol. 4, no. 4, p. e5077, 2009.
- [65] F. Peña and J. M. Ramirez, “Hypoxia-induced changes in neuronal network properties,” *Molecular Neurobiology*, vol. 32, no. 3, pp. 251–284, 2005.
- [66] Z. Javed, K. Khan, Q. Raza et al., “Notch signaling and MicroRNA: the dynamic duo steering between neurogenesis and glioblastomas,” *Cellular and Molecular Biology*, vol. 67, no. 2, pp. 33–43, 2021.
- [67] X. Ruan, G. Liu, J. Zhou et al., “Zbed3 is indispensable for Wnt signaling regulation of cortical layers formation in developing brain,” *Cerebral Cortex*, vol. 31, no. 9, pp. 4078–4091, 2021.
- [68] M. C. Simon and B. Keith, “The role of oxygen availability in embryonic development and stem cell function,” *Nature Reviews. Molecular Cell Biology*, vol. 9, no. 4, pp. 285–296, 2008.
- [69] L.-L. Zhu, W. Li-Ying, D. T. Yew, and M. Fan, “Effects of hypoxia on the proliferation and differentiation of NSCs,” *Molecular Neurobiology*, vol. 31, no. 1-3, pp. 231–242, 2005.
- [70] S. Bansod, R. Kageyama, and T. Ohtsuka, “Hes5 regulates the transition timing of neurogenesis and gliogenesis in mammalian neocortical development,” *Development (Cambridge, England)*, vol. 144, no. 17, pp. 3156–3167, 2017.
- [71] L. S. Campos, A. J. Duarte, T. Branco, and D. Henrique, “mDll1 and mDll3 expression in the developing mouse brain: role in the establishment of the early cortex,” *Journal of Neuroscience Research*, vol. 64, no. 6, pp. 590–598, 2001.
- [72] S. L. Dunwoodie, D. Henrique, S. M. Harrison, and R. S. Beddington, “Mouse Dll3: a novel divergent delta gene which may complement the function of other delta homologues during early pattern formation in the mouse embryo,” *Development*, vol. 124, no. 16, pp. 3065–3076, 1997.
- [73] J. Hatakeyama, Y. Bessho, K. Katoh et al., “Hes genes regulate size, shape and histogenesis of the nervous system by control of the timing of neural stem cell differentiation,” *Development (Cambridge, England)*, vol. 131, no. 22, pp. 5539–5550, 2004.
- [74] E. Ladi, J. T. Nichols, W. Ge et al., “The divergent DSL ligand Dll3 does not activate Notch signaling but cell autonomously attenuates signaling induced by other DSL ligands,” *The Journal of Cell Biology*, vol. 170, no. 6, pp. 983–992, 2005.
- [75] T. Ohtsuka, M. Sakamoto, F. Guillemot, and R. Kageyama, “Roles of the basic helix-loop-helix genes Hes1 and Hes5 in expansion of neural stem cells of the developing brain,” *The Journal of Biological Chemistry*, vol. 276, no. 32, pp. 30467–30474, 2001.
- [76] C. Ramos, S. Rocha, C. Gaspar, and D. Henrique, “Two notch ligands, Dll1 and Jag1, are differently restricted in their range of action to control neurogenesis in the mammalian spinal cord,” *PLoS One*, vol. 5, no. 11, p. e15515, 2010.
- [77] P. Yuan, W. Han, L. Xie et al., “The implications of hippocampal neurogenesis in adolescent rats after status epilepticus: a novel role of notch signaling pathway in regulating

- epileptogenesis," *Cell and Tissue Research*, vol. 380, no. 3, pp. 425–433, 2020.
- [78] C. Knox, V. Camberos, L. Ceja et al., "Long-term hypoxia maintains a state of dedifferentiation and enhanced stemness in fetal cardiovascular progenitor cells," *International Journal of Molecular Sciences*, vol. 22, no. 17, p. 9382, 2021.
- [79] M. Sakamoto, H. Hirata, T. Ohtsuka, Y. Bessho, and R. Kageyama, "The basic helix-loop-helix genes *Hesr1/Hey1* and *Hesr2/Hey2* regulate maintenance of neural precursor cells in the brain," *The Journal of Biological Chemistry*, vol. 278, no. 45, pp. 44808–44815, 2003.
- [80] I. Lahmann, Y. Zhang, K. Baum, J. Wolf, and C. Birchmeier, "An oscillatory network controlling self-renewal of skeletal muscle stem cells," *Experimental Cell Research*, vol. 409, no. 2, p. 112933, 2021.
- [81] H. Shimojo, T. Ohtsuka, and R. Kageyama, "Oscillations in notch signaling regulate maintenance of neural progenitors," *Neuron*, vol. 58, no. 1, pp. 52–64, 2008.
- [82] C. Boucherie, C. Boutin, Y. Jossin et al., "Neural progenitor fate decision defects, cortical hypoplasia and behavioral impairment in *Celsr1*-deficient mice," *Molecular Psychiatry*, vol. 23, no. 3, pp. 723–734, 2018.
- [83] M.-H. Jang, M. A. Bonaguidi, Y. Kitabatake et al., "Secreted frizzled-related protein 3 regulates activity-dependent adult hippocampal neurogenesis," *Cell Stem Cell*, vol. 12, no. 2, pp. 215–223, 2013.
- [84] J. Kele, R. Emma, J. C. Andersson et al., "SFRP1 and SFRP2 dose-dependently regulate midbrain dopamine neuron development in vivo and in embryonic stem cells," *Stem Cells*, vol. 30, no. 5, pp. 865–875, 2012.
- [85] D. Kunke, V. Bryja, L. Mygland, E. Arenas, and S. Krauss, "Inhibition of canonical Wnt signaling promotes gliogenesis in P0-NSCs," *Biochemical and Biophysical Research Communications*, vol. 386, no. 4, pp. 628–633, 2009.
- [86] S. V. Ekholm and S. I. Reed, "Regulation of G₁ cyclin-dependent kinases in the mammalian cell cycle," *Current Opinion in Cell Biology*, vol. 12, no. 6, pp. 676–684, 2000.
- [87] J. W. Hofmann, T. McBryan, P. D. Adams, and J. M. Sedivy, "The effects of aging on the expression of Wnt pathway genes in mouse tissues," *Age (Dordrecht, Netherlands)*, vol. 36, no. 3, p. 9618, 2014.
- [88] P. Salles and U. de Andrade, "Expression of D-type cyclins in differentiating cells of the mouse spinal cord," *Genetics and Molecular Biology*, vol. 30, no. 3, pp. 702–708, 2007.
- [89] X. Jiang, J. Zhao, J. Lili et al., "Temporal expression patterns of insulin-like growth factor binding protein-4 in the embryonic and postnatal rat brain," *BMC Neuroscience*, vol. 14, no. 1, p. 132, 2013.
- [90] R. Herken, W. Götz, and K. H. Wattjes, "Initial development of capillaries in the neuroepithelium of the mouse," *Journal of Anatomy*, vol. 164, pp. 85–92, 1989.
- [91] I. Paredes, P. Himmels, and C. Ruiz de Almodóvar, "Neurovascular communication during CNS development," *Developmental Cell*, vol. 45, no. 1, pp. 10–32, 2018.
- [92] R. C. Bunn, W. D. King, M. K. Winkler, and J. L. Fowlkes, "Early developmental changes in IGF-I, IGF-II, IGF binding protein-1, and IGF binding protein-3 concentration in the cerebrospinal fluid of children," *Pediatric Research*, vol. 58, no. 1, pp. 89–93, 2005.
- [93] C.-H. Chen, P.-Y. Chen, Y.-Y. Lin et al., "Suppression of tumor growth via IGFBP3 depletion as a potential treatment in glioma," *Journal of Neurosurgery*, vol. 132, no. 1, pp. 168–179, 2020.
- [94] T. Sun, X. J. Wang, S. S. Xie et al., "A comparison of proliferative capacity and passaging potential between neural stem and progenitor cells in adherent and neurosphere cultures," *International Journal of Developmental Neuroscience*, vol. 29, no. 7, pp. 723–731, 2011.
- [95] V. Vukicevic, A. Jauch, T. C. Dinger et al., "Genetic instability and diminished differentiation capacity in long-term cultured mouse neurosphere cells," *Mechanisms of Ageing and Development*, vol. 131, no. 2, pp. 124–132, 2010.
- [96] M. Andang, J. Hjerling-Leffler, A. Moliner et al., "Histone H2AX-dependent GABA_A receptor regulation of stem cell proliferation," *Nature*, vol. 451, no. 7177, pp. 460–464, 2008.
- [97] M. Fukui, N. Nakamichi, M. Yoneyama et al., "Modulation of cellular proliferation and differentiation through GABAB receptors expressed by undifferentiated neural progenitor cells isolated from fetal mouse brain," *Journal of Cellular Physiology*, vol. 216, no. 2, pp. 507–519, 2008.
- [98] J. J. LoTurco, D. F. Owens, M. J. S. Heath, M. B. E. Davis, and A. R. Kriegstein, "GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis," *Neuron*, vol. 15, no. 6, pp. 1287–1298, 1995.
- [99] D. D. Wang and A. R. Kriegstein, "Defining the role of GABA in cortical development," *The Journal of Physiology*, vol. 587, no. 9, pp. 1873–1879, 2009.