


Combating Noncommunicable Diseases such as Cancer and Liver Toxicities Using Natural Products

Lead Guest Editor: Nikhil B. Ghate

Guest Editors: Sourav Panja and Abhishek Das





Combating Noncommunicable Diseases such as Cancer and Liver Toxicities Using Natural Products

**Combating Noncommunicable Diseases
such as Cancer and Liver Toxicities
Using Natural Products**

Lead Guest Editor: Nikhil B. Ghatе


Guest Editors: Sourav Panja and Abhishek Das

Chief Editor


Kaustubha Mohanty, India

Associate Editors

Mohammad Al-Ghouti, Qatar

Tingyue Gu , USA

Teodorico C. Ramalho , Brazil

Artur M. S. Silva , Portugal

Contents

Anti-Inflammatory Compounds from Vietnamese *Piper bavinum*

Viet Dung Hoang, Phi Hung Nguyen, Minh Thu Doan, Manh Hung Tran, Nhu Tuan Huynh, Huu Tung Nguyen , Byung Sun Min, and Dao Cuong To 

Research Article (7 pages), Article ID 1038636, Volume 2020 (2020)

Research Article

Anti-Inflammatory Compounds from Vietnamese *Piper bavinum*

Viet Dung Hoang,¹ Phi Hung Nguyen,² Minh Thu Doan,³ Manh Hung Tran,³
Nhu Tuan Huynh,⁴ Huu Tung Nguyen ,^{5,6} Byung Sun Min,⁷ and Dao Cuong To ^{5,6}

¹Institute of Pharmaceutical Training, Vietnam Military Medical University, 160 Phung Hung Street, Hadong District, Hanoi 151000, Vietnam

²Institute of Natural Products Chemistry, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet, Cau Giay District, Hanoi 122100, Vietnam

³Department of Biomedical Sciences, Institute for Research and Executive Education (VNUK), The University of Danang, 158A Le Loi Street, Hai Chau District, Da Nang 551000, Vietnam

⁴Faculty of Pharmacy, Dong A University, 33 Xo Viet Nghe Tinh, Hai Chau District, Da Nang 550000, Vietnam

⁵Faculty of Pharmacy, Phenikaa University, Yen Nghia, Ha Dong District, Hanoi 12116, Vietnam

⁶Phenikaa Research and Technology Institute (PRATI), A&A Green Phoenix Group JSC, 167 Hoang Ngan, Cau Giay District, Hanoi 11313, Vietnam

⁷College of Pharmacy, Drug Research and Development Center, Catholic University of Daegu, Gyeongsan 712-702, Republic of Korea

Correspondence should be addressed to Dao Cuong To; cuong.todao@phenikaa-uni.edu.vn

Received 11 September 2019; Revised 3 December 2019; Accepted 10 January 2020; Published 8 February 2020

Guest Editor: Nikhil B. Ghate

Copyright © 2020 Viet Dung Hoang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study reports the anti-inflammatory activity-guided fractionation of the aerial part of *Piper bavinum* C. CD. (Piperaceae) that led to the isolation of eight secondary metabolites (1–8). The chemical structures of 1–8 were established mainly by NMR and mass spectra. Compound 5 was isolated from *P. bavinum* for the first time. All the isolated compounds were evaluated against LPS-induced NO production in macrophage RAW 264.7 cells *in vitro*. Among them, compound 4 showed the most potent inhibitory activity against the LPS-induced NO production with an IC₅₀ value of 5.2 μ M followed by compound 5 that inhibited NO production with an IC₅₀ value of 13.5 μ M. In the protein levels, compound 4 suppressed LPS-induced COX-2 and iNOS expressions in a dose-dependent manner. The results suggested that *P. bavinum* and its constituents might exert anti-inflammatory effects.

1. Introduction

Inflammation is a protective response that occurs following trauma, infection, or tissue injury [1]. In this process, activated inflammatory cells include increased amounts of nitric oxide (NO) and prostaglandin E2 (PGE2). NO is a major product that is controlled by nitric oxide synthases (NOSs). NOSs included inducible nitric oxide synthase (iNOS) that is extremely expressed in macrophage cells, and the activation of iNOS generally leads to some autoimmune diseases [2]. PGE2 is another important inflammatory mediator, produced by arachidonic acid metabolites via the catalysis of cyclooxygenase-2 (COX-2) [2]. Lipopolysaccharide (LPS) plays an

important role in activating immune cells to upregulate to inflammatory states. The overproduction of NO by iNOS regulation has been concerned in the pathology of several inflammatory disorders, including septic shock, tissue damage, and rheumatoid arthritis [3–5]. NO and PGE2 production induced by LPS through iNOS and COX-2, respectively, can reflect the degree of inflammation.

During a screening program to discover inflammation inhibitors from natural sources, we have found that the *n*-hexane and ethyl acetate (EtOAc) fractions of Vietnamese *P. bavinum* exhibited appreciable inhibitory activity in LPS-induced NO production in macrophage RAW 264.7 cells. *P. bavinum* is a member of the *Piper* genus, which is the largest genus in the

Piperaceae family. The *Piper* genus included over 700 species and is identified in the tropical regions [6, 7]. *Piper* species have been used in traditional medicine to treat the gynecological ailment, gastrointestinal problem, and depression [7]. Some of these from India, Southeast Asia, and Africa are of high commercial, medicinal, and economic importance since they are used as spices and traditional medicines [8]. Otherwise, *Piper* species have also been found to possess biological activities, including antioxidative, antimicrobial, antiproliferative/anticancer, antiparasitic, and neuropharmacological activities [9]. Especially, *in vivo* and *in vitro* studies showed the *Piper* genus against inflammation such as *P. nigrum*, *P. crocatum*, *P. betle*, *P. umbellatum*, *P. gaudichaudianum*, *P. arboreum*, *P. umbellata*, *P. fuliginum*, *P. longum*, and *P. methysticum* [9]. The most widely recognized species of the genus *Piper* is *P. nigrum* followed by *P. longum*, *P. chaba*, *P. mullesia*, *P. umbellatum*, *P. hymenophyllum*, *P. argyrophyllum*, *P. attenuatum*, *P. colubrinum*, *P. galeatum*, and *P. bavinum* [10]. *P. bavinum* is a liana, distributed in forests of India, China, and Vietnam at altitudes of 1300–1700 m [11]. *P. bavinum* is known as an oriental medicinal plant and had been reported to possess various pharmacological activities such as antibacterial [11] and anticholinesterase [6]. Previous studies on chemical constituents of *P. bavinum* showed the presence of essential oil [11], terpenoids, phenolics, and flavonoids [6]. However, there has been no investigation in chemical constituents and inhibitory activity of NO production of *P. bavinum*. This paper describes the isolation and structural elucidation of the isolated compounds as well as evaluates their inhibitory activity on NO production.

2. Results and Discussion

2.1. Determination and Elucidation of Isolated Compounds. The methanol extract was partitioned with *n*-hexane, chloroform (CHCl₃), ethyl acetate (EtOAc), and *n*-butanol (BuOH) to obtain *n*-hexane, CHCl₃, EtOAc, and BuOH soluble fractions, respectively. Based on bioactivity-guided fractionation (data not are shown), the *n*-hexane and EtOAc fractions were subjected to column chromatography on a silica gel and C18-RP silica gel column to obtain eight compounds (**1–8**) (see Figure 1).

Compound **1** was obtained as a light yellow powder. The presence of two oxymethine groups (δ_{H} 4.82 (H-2)/ δ_{C} 83.2 (C-2), and δ_{H} 4.45 (H-2)/ δ_{C} 71.6 (C-3)) and a ketone at δ_{C} 197.7 (C-4) in the ¹H- and ¹³C-NMR spectra indicated **1** to be a flavanone (see Figure 1 and Supplementary Materials (available here)). The ¹H-NMR spectrum of **1** further showed signals of two tetrasubstituted benzene rings (δ_{H} 5.88 (1H, s, H-6), 5.91 (1H, s, H-8), and 6.52 (2H, s, H-2'/6')) [12]. The ¹³C-NMR spectrum exhibited fifteen signals due to twelve *sp*² carbons of two benzene rings, a ketone at δ_{C} 197.7 (C-4), and two oxymethine carbons (δ_{C} 83.2 (C-2) and 71.6 (C-3) (see Figure 1 and Supplementary Materials)). The HRESIMS spectrum of **1** showed the pseudomolecular ion at $[M + H]^+$ at m/z 321.0597, indicating the molecular formula C₁₅H₁₂O₈. The positive optical rotation $[\alpha]_{\text{D}}^{25} + 11.2^\circ$ (*c* 0.2, MeOH) hinted that **1** probably shared the (*R*, *R*) configuration. Based on this evidence and in comparison with the published data, compound **1** was identified as (2*R*, 3*R*)-3,5,7-

trihydroxy-2-(3,4,5-trihydroxyphenyl)-2,3-dihydrochromen-4-one (ampelopsin or dihydromyricetin) [6,13]. This compound was found to possess anticholinesterase [6], antiatherosclerosis (anti-inflammatory), antidiabetes, and antitumor activities and effects on cardioprotection, hepatoprotection, neuroprotection, and dermatoprotection [14].

Compound **2** was obtained as a yellow powder. The presence of a methine group (δ_{H} 6.54 (H-3)/ δ_{C} 102.6 (C-3)), an oxygenated carbon at δ_{C} 165.3 (C-2), and a carbonyl carbon at δ_{C} 182.6 (C-4) in the ¹H- and ¹³C-NMR spectra indicated **2** to be a flavone [12]. The ¹H-NMR of **2** showed signals the presence of 1,4-disubstituted benzene ring characterized with the A₂B₂ system (δ_{H} 77.91 (2H, d, *J* = 8.4 Hz, H-2'/6'), 6.89 (2H, d, d, *J* = 8.4 Hz, H-3'/5')) together with two sugar (δ_{H} 5.05 (1H, d, *J* = 9.6 Hz, H-1''), 5.16 (1H, br s, H-1''')) for the glucose and rhamnose moieties (see Figure 1 and Supplementary Materials). The ¹³C-NMR spectrum exhibited twenty-seven signals due to twelve *sp*² carbons of two benzene rings, a ketone, and twelve carbons of two sugar moieties. In the HMBC spectrum, the correlations of proton H-1'' at δ_{H} 5.05 (1H, d) to carbon signals at δ_{C} 109.1 (C-6), 157.6 (C-5), and 163.1 (C-7), as well as the correlations of proton H-1''' at δ_{H} 5.16 (1H, br s) to carbon signals at δ_{C} 101.8 (C-8), 155.9 (C-9), and 163.1 (C-7), were observed suggesting the glucose and rhamnose moieties were located at C-6 and C-8, respectively (see Figure 1 and Supplementary Materials). Therefore, compound **2** was identified as violanthin in comparison with literature data [6, 15]. Violanthin has been shown to possess antioxidant [16] and anticholinesterase activities [6].

Compounds **3** and **6** were isolated as a white amorphous powder. Their ¹H-NMR spectra displayed characteristic signals due to aromatic protons of two benzene rings and a carbinol methylene (2H-7'), while their ¹³C-NMR spectra revealed the signals of a carbonyl carbon (C-7), a carbinol methylene carbon (C-7'), and twelve *sp*² carbons of two benzene rings (see Figure 1 and Supplementary Materials). The above observation indicated these compounds were benzyl benzoate derivatives [17]. Detailed analysis of the ¹H- and ¹³C-NMR spectra revealed that **3** possessed ten aromatic protons of two benzene rings, a carbinol methylene (δ_{H} 5.39 (2H, s, H-7')/ δ_{C} 66.6 (C-7')), and a carbonyl carbon at δ_{C} 166.3 (C-7) (see Figure 1 and Supplementary Materials). Compound **6** also possessed one carbonyl carbon (C-7) and a carbinol methylene (C-7') but showed only nine aromatic protons and an additional one methoxy (see Figure 1). The HMBC correlated from H-4'/H-6'/H-7' and protons of a methoxy group (δ_{H} 3.86) to C-2' (δ_{C} 158.0), suggested the methoxy group was located at C-2' (see Figure 1 and Supplementary Materials). The HRESIMS spectrum of **6** showed the pseudomolecular ion at $[M - H]^-$ at m/z 241.1213, indicating the molecular formula C₁₅H₁₄O₃. Comparison of the ¹H- and ¹³C-NMR data of these compounds with those published in the literature led to the structural identification of compounds **3** and **6** to be benzyl benzoate [18] and 2-methoxybenzyl benzoate [19], respectively. Compound **3** (benzyl benzoate) has been found to


FIGURE 1: Structure of compounds 1–8 from *P. bavinum*.

exhibit antibacterial activity [19], and compound 6 (2-methoxybenzyl benzoate) exhibits inhibitory activity against the nuclear factor of activated T cells (NFAT) transcription factor [6].

Compound 4 was obtained as a colorless oil. The HRFABMS spectrum of 4 showed the pseudomolecular ion at $[M]^+$ at m/z 232.1829, indicating the molecular formula $C_{16}H_{24}O$. The 1H NMR data of 4 revealed the presence of an OH group (δ_H 5.91, 1H, br s) and 1,4-disubstituted benzene ring characterized by the A_2B_2 system, confirming the presence of a *para*-substituted phenol. The 1H NMR signals showed the presence of an OH group, a benzylic methylene (H-1'), a double bond (H-2' and H-3'), allylic methylene (H-4'), five methylenes (H-5' to H-9'), an A_2B_2 system of benzene ring, and an additional one methyl group (H-10'), indicating a decenyl chain (see Figure 1 and Supplementary Materials). The ^{13}C NMR and DEPT spectra revealed the signals of two olefinic carbons, six aromatic carbons, seven methylene carbons, and a methyl carbon (see Figure 1 and Supplementary Materials). The stereochemistry of C-2' was determined as *cis* by comparison of the ^{13}C NMR chemical shifts of the allylic carbons (δ_C 32.7 (C-1') and 27.2 (C-4')) in 4 with those of the *E* form in myricanene B (δ_C 34.3 (C-1') and 32.3 (C-4')). Analysis of these signals by the COSY, HMQC, and HMBC spectra led to the partial structures of 4 (see Figure 1). Therefore, compound 4 was identified as 4-(2'-(*Z*)-decenyl) phenol (bavinol A) in comparison with literature data [6]. Bavinol A was found to exhibit anticholinesterase activity [6].

Compound 5 was also obtained as a colorless oil. The 1H NMR spectrum of 5 displayed signals due to five methyls (H-16, H-17, H-18, H-19, and H-20), nine methylenes, an oxygenated methylene at δ_H 4.11 (H-1), three methines (H-7, H-11, and H-15), and an olefinic δ_H 5.37 (H-3) (see Figure 1 and Supplementary Materials). The ^{13}C NMR spectrum revealed the signals of two olefinic carbons (δ_C 123.3 (C-2), 140.5 (C-3)), five methyl carbon (δ_C 22.8 (C-16), 22.9 (C-17), 19.9 (C-18), 19.9 (C-19), and 16.4 (C-20)), nine methylene carbons, three methane carbons at δ_C 33.0 (C-7),

32.9 (C-11), and 28.2 (C-15), and an oxygenated methylene carbon at δ_C 59.6 (C-1) (see Figure 1 and Supplementary Materials). The correlations from H-20 to C-2, C-3, and C-4; H-2 to C-1, C-3, C-4, and C-20; and 2H-1 to C-2 and C-3 were observed in the HMBC. Further analysis of these signals by the COSY, HMQC and HMBC spectra led to the partial structures of 5 (see Figure 1). The stereochemistry of C-2 was determined as *trans* by comparison of the ^{13}C NMR chemical shifts. The positive optical rotation $[\alpha]_D^{25} + 9.7^\circ$ (*c* 0.05, MeOH) hinted that 5 probably shared the [R, R] configuration. Thus, the structure of compound 5 was assigned to be (2*E*, 7*R*, and 11*R*)-3,7,11,15-tetramethyl-2-hexadecen-1-ol (*trans*-phytol) [19] and isolated from *P. bavinum* for the first time. This compound was also isolated from *Piper kadsura* [20]. *Trans*-phytol was found to possess antinociceptive and antioxidant [21], cytotoxic [22], anti-schistosomal [23], and arthritis activities [24] and was also found to inhibit neutrophil migration [25].

Compound 7 was isolated as a colorless solid with negative optical rotation, $[\alpha]_D^{25} - 42.3^\circ$. The HRFABMS spectrum of 7 showed the pseudomolecular ion at $[M + H]^+$ at m/z 427.1393, indicating the molecular formula $C_{23}H_{22}O_8$. The 1H NMR spectrum of 7 displayed characteristic signals due to ten aromatic protons of two benzene rings, three oxygenated methines (δ_H 4.29 (H-2), 5.77 (H-3), and 5.68 (H-6)), two olefinic (δ_H 5.95 (H-4) and 5.87 (H-5)), an oxygenated methylene (δ_H 4.70 (H-7a) and 4.67 (H-7b)), an acetyl group (δ_H 2.02 (6-OAc)), and an OH group at δ_H 4.90 (1-OH) (see Figure 1 and Supplementary Materials). The ^{13}C NMR spectrum revealed the signals of two carbonyl carbons (C-7' and C-7''), two olefinic carbons (C-4 and C-5), twelve sp^2 carbons of two benzene rings, three oxygenated methines (C-2, C-3, and C-6), an oxygenated methylene (C-7), a quaternary carbon (C-1), and an acetyl group at δ_C 171.0 and 20.9 (see Figure 1 and Supplementary Materials). The long-range correlations between H-4 and C-7'', as well as between 2H7 and C-7', were observed in the HMBC spectrum, suggesting that two benzoyl groups were located at C-7 and C-4. In addition, the HMBC correlation from H-6 to

carbonyl carbon at δ_C 171.0 was also observed, indicating the acetyl group was located at C-6. Further analysis of these signals by the COSY, HMQC, and HMBC spectra led to the partial structures of **7** (see Figure 1). The CD Cotton effects $\Delta\epsilon_{236} - 6.24$ and $\Delta\epsilon_{282} + 4.32$ indicated that compound **7** probably shared the [S, S, R, S] configuration. Based on this evidence and in comparison with the published data, compound **7** was identified as (-)-6-acetylzeylenol [26]. This compound has been shown to exhibit cytotoxic activity [27].

Compound **8** was obtained as a white powder with positive optical rotation, $[\alpha]_D^{25} + 67.5^\circ$. The ^1H NMR data of **8** revealed the presence of three methyl groups (H-11, H-12, and H-13), two *trans* olefinic protons (H-7 and H-8), two methylene (H-2 and H-10), and two methine protons (H-4 and H-9) (see Figure 1 and Supplementary Materials). The ^{13}C NMR spectrum revealed the signals of a ketone (C-3), three methyl carbons (C-11, C-12, and C-13), four olefinic carbons (C-4, C-5, C-7, and C-8), two methylene carbons (C-2 and C-10), an methine (C-9), and two quaternary carbons (C-1 and C-6) (see Figure 1 and Supplementary Materials). The HRESIMS spectrum of **8** showed the pseudomolecular ion at $[\text{M} + \text{Na}]^+$ at m/z 263.1244, indicating the molecular formula $\text{C}_{13}\text{H}_{20}\text{O}_4$. The CD spectrum of **8** exhibited a positive Cotton effect ($\Delta\epsilon_{242} + 11.56$), indicating that the 6-position was shown to have an S-configuration. The absolute configuration at the 9-position was elucidated to be S when comparing with the ^1H , ^{13}C NMR, and the positive optical rotation value [27]. The 1D and 2D NMR spectra indicated this compound had megastigmane skeleton and identified as cucumegastigmane I in comparison with the literature [28].

2.2. NO Production Inhibition and the Cell Viability Assay.

In the first experiment, the cytotoxic assay was performed to determine the safe and nontoxic concentration of isolated compounds (**1–8**) for the next assay. Nontoxicity of the isolated compounds indicated by over 90% of cell viability by MTS assay [29]. The isolated compounds in the concentration of 100 μM were toxic toward RAW 264.7 cells; therefore, the respective concentration was not used for the treatments, and the concentrations of 30, 10, and 3 μM were chosen to further study (Figure 2).

To determine the effects of these compounds on the LPS-induced production of NO in RAW 264.7 cells, a cell culture medium was harvested, and the production of nitrite was measured using the Griess reaction [30–32]. As shown in Table 1, bavinol A (**4**) showed the strongest inhibitory on NO production among the isolated compounds with an IC_{50} value of 5.2 μM , followed by *trans*-phytol (**5**) exhibiting inhibitory effects with an IC_{50} value of 13.5 μM . Interestingly, the other compounds showed weak inhibitory activity with IC_{50} values over 30 μM (see Table 1). In this assay, celastrol, a natural secondary metabolite, was used as a positive inhibitor. This compound expressively withdrew LPS-induced NO production with an IC_{50} value of 1.0 μM [30, 31].

Neither LPS nor the samples were added to the control group. Thus, the inhibitory effects of these compounds on NO production were not attributable to any cytotoxic effect.

As shown in Figure 3, after LPS (1 $\mu\text{g}/\text{mL}$) stimulation, the NO production increased by approximately 13-fold after 24 h in the control. Compounds **4** and **5** reduced the NO production 24 h after LPS stimulation, in a dose-dependent manner (Figure 3).

It is well-known that COX-2 is induced by cytokines and other activators such as LPS, resulting in the release of large amounts of PGE2 in macrophage cells [2]. Among the tested compounds, bavinol A (**4**) exhibited the strongest inhibitory effect on the LPS-induced production of NO in macrophage RAW 264.7 cells. Therefore, western blot was performed to determine the inhibitory effects of bavinol A (**4**) on the modulation of iNOS and COX-2 expression [33] (Supplementary Materials). As shown in Figure 4, bavinol A (**4**) (0–30 μM) showed a dose-dependent reduction in LPS-induced iNOS and COX-2 expressions but did not change the alpha-tubulin expression (see Figure 4). The results showed that bavinol A (**4**) inhibited iNOS and COX-2 activities in LPS-stimulated RAW 264.7 cells at the transcription level.

Macrophage cells are principally concerned in acute and chronic inflammatory responses. Among the inflammation stimuli, LPS is known as lipoglycans and endotoxins that stimulated macrophages to induce the expression of iNOS protein to produce NO. In the meantime, NO plays as an activator of macrophages to kill microorganisms through signal transduction [34]. On the contrary, the overproduction of NO in the immune system might cause immune hypersensitivity reactions followed by tissue or cell injury [3–5]. NO is mostly synthesized by iNOS; however, the extensively higher amount of NO coactively activates inflammatory processes in conjunction with other inflammatory mediators [35]. Thus, the inhibition of iNOS activity or downregulation of iNOS expression could be beneficial to inhibit inflammatory responses. From the results, flavonoid glycosides (violanthin, **2**), a derivative of apigenin, did not significantly inhibit NO production at 30 μM , regardless of the aglycones (flavone) present and the glycoside linkages (C- or O-glycosides) (Figure 1). The glycosylation of apigenin to violanthin (**2**) resulted in a loss of nitrite inhibitory production in RAW 264.7 murine macrophages. In general, flavones showed strong inhibition of NO production. Previous studies showed that luteolin and apigenin (have a C-2,3 double bond) inhibited NO production via iNOS downregulation [36]. These results powerfully suggested that the C-2,3-double bond could be an essential factor for inhibiting NO production. It was found that flavanone derivatives (ampelopsin **1**), which do not have a C-2,3 double bond (Figure 1), were inactive up to 30 μM . These results also indicated that a planar ring system in the flavonoid molecule might be important for NO inhibition [37].

Lv et al. tested the effects of the megastigmane derivatives from *Lyonia ovalifolia* on NO production. The *in vitro* results indicated that megastigmane derivatives such as abscisic acid- β -D-glucopyranosyl ester and pisumionoside inhibited NO production from LPS-activated RAW 264.7 cells with inhibition rates of 54.5% and 83.3% at the concentration of 10^{-5} M, respectively. Pisumionoside was also found to possess better inhibitory activity than the other megastigmane derivatives tested [38]. Both of them revealed the presence of the




FIGURE 2: Effect on cell viability by LPS stimulation in the presence of compounds 1–8. Cell viability was determined by the MTS assay and expressed as a percentage of the control without the addition of indicated compounds 1–8.

TABLE 1: NO production inhibitory activity of isolated compounds 1–8.

Compound	IC ₅₀ value (μM) ^a
1	>30
2	>30
3	>30
4	5.2 ± 0.4
5	13.5 ± 1.5
6	>30
7	>30
8	>30
Celastrol ^b	1.0 ± 0.1

^aThe inhibitory effects are represented as the molar concentration (μM) giving 50% inhibition (IC₅₀) relative to the vehicle control. These data represent the average values of three repeated experiments (mean ± SD).

^bPositive control for NO.

conjugated double bond of the olefinic group with carbonyl carbon in the structures. Similarly, Trang et al. performed a study regarding the effect of megastigmane derivatives on NO production, and the results showed that these megastigmane derivatives were inactive [39]. In our experiment, cucumegastigmane I (8), a derivative of megastigmane, was also inactive on NO production, presumably because of the absence of the conjugated double bond of the olefinic group with carbonyl carbon in 8 (Figure 1).

Phenolics including alkenyl phenols are essential compounds for the suppression of inflammation among phytochemicals. Cuong et al. studied the anti-inflammatory effects of phenolic compounds, and the results showed a strong inhibitory on NO production of isolated compounds because of the presence of the conjugated double bond of the olefinic group, and the 3,4-hydroxylation(s) of benzene ring gave favorable results [31, 32, 40]. In our study, bavinol A (4), which belongs to the alkenyl phenol class, exhibited the strongest inhibitory on NO production among the isolated compounds. However, the inhibitory activities on NO production of bavinol A (4) were weaker than active




FIGURE 3: Inhibitory effect of compounds 4 and 5 on the LPS-induced NO production in RAW 264.7 cells. RAW 264.7 cells were pretreated with different concentrations (1, 3, 10, and 30 μM) of the tested compounds for 1 h and then with LPS (1 μg/mL) and incubated for 24 h. Control values were obtained in the absence of LPS and compounds. The data are expressed as the mean ± SD ($n = 3$). Statistical significance was assessed by two-tailed unpaired Student's t -test, and $P < 0.05$ was considered statistically significant.

compounds presumably because of the lack of 3-hydroxylation of the benzene ring [31, 32, 40]. Compound 5 (*trans*-phytol) containing the conjugated double bond of the olefinic group in the structure also exhibited a strong inhibitory on NO production. However, the inhibitory activities on NO production of *trans*-phytol (5) were significantly reduced presumably because of the absence of the benzene ring contained the 3,4-hydroxylation(s) (Figure 1). In opposite, other phenolic compounds (3, 6, and 7) were inactive presumably because of the absence of the conjugated double bond of the olefinic group as well as 3,4-hydroxylation(s) of benzene ring (Figure 1). Finally, our obtained data suggested that phenolic compounds, particularly the alkenyl phenols bearing the conjugated double bond as well as 3,4-hydroxylation(s) of the benzene ring, could be considered as




FIGURE 4: Inhibition of LPS-induced iNOS expression in RAW 264.7 cells by compound 4 (C4). RAW 264.7 cells were pretreated for 30 min at indicated concentrations of 0, 3, 10, and 30 μM , followed by stimulation with LPS ($1\mu\text{g/mL}$) for 24 h. Whole-cell lysates were blotted with the indicated antibodies. α -tubulin level was used as a loading control. The expression levels of iNOS and COX-2 were determined by the analysis of immunoblot.

new lead compounds for the development of agents against NO production. Moreover, the alkenyl phenol-enrich extracts may be applied as supplemental and/or functional foods having a beneficial effect against inflammation.

3. Conclusion

This study demonstrated for the first time the NO production inhibitory activities of isolated compounds from *P. bavinum*. NO production inhibitory guided fractionation led to the isolation of eight compounds, ampelopsin (1), violanthin (2), benzyl benzoate (3), bavinol A (4), *trans*-phytol (5), 2-methoxybenzyl benzoate (6), (–)-6-acetylzeylenol (7), and cucumegastigmane I (8), from the aerial part of *P. bavinum*. *Trans*-phytol (5) was isolated from *P. bavinum* for the first time. Bavinol A (4) and *trans*-phytol (5) showed inhibitory effects with IC_{50} values of 5.2 and 13.5 μM , respectively, but the others were inactive. In addition, bavinol A (4) showed a dose-dependent reduction in LPS-induced iNOS and COX-2 expressions but did not change the α -tubulin expression. The results showed that bavinol A (4) inhibited iNOS and COX-2 activities in LPS-stimulated RAW 264.7 cells at the transcription level. This result provides experimental evidence to support that bavinol A may serve as a useful anti-inflammation agent.

Data Availability

The data used to support the findings of this study are included within the article.

Conflicts of Interest

The authors declare that they have no conflicts of interest regarding the publication of this paper.

Acknowledgments

This work was supported by the Phenikaa University Foundation for Science and Technology Development.

Supplementary Materials

Supplement Material S1: materials and methods. It describes information about general experimental procedures, plant materials, extraction and isolation, physiochemical and NMR data of isolated compounds, NO production inhibitory and the cell viability assays, as well as immunoblot analysis. Figure S1: HR-ESI mass spectrum of compound 1. Figure S2: HR-FAB mass spectrum of compound 4. Figure S3: HR-ESI mass spectrum of compound 6. Figure S4: HR-ESI mass spectrum of compound 7. Figure S5: HR-ESI mass spectrum of compound 8. (*Supplementary Materials*)

References

- [1] S. Zedler and E. Faist, "The impact of endogenous triggers on trauma-associated inflammation," *Current Opinion in Critical Care*, vol. 12, no. 6, pp. 595–601, 2006.
- [2] A. Murakami and H. Ohigashi, "Targeting NOX, iNOS and COX-2 in inflammatory cells: chemoprevention using food phytochemicals," *International Journal of Cancer*, vol. 121, no. 11, pp. 2357–2363, 2007.
- [3] A. J. Farrell, D. R. Blake, R. M. Palmer, and S. Moncada, "Increased concentrations of nitrite in synovial fluid and serum samples suggest increased nitric oxide synthesis in rheumatic diseases," *Annals of the Rheumatic Diseases*, vol. 51, no. 11, pp. 1219–1222, 1992.
- [4] R. G. Kilbourn, A. Jubran, S. S. Gross et al., "Reversal of endotoxin-mediated shock by NG-methyl-L-arginine, an inhibitor of nitric oxide synthesis," *Biochemical and Biophysical Research Communications*, vol. 172, no. 3, pp. 1132–1138, 1990.
- [5] K. D. KRöncke, K. Fehsel, and V. Kolb-Bachofen, "Inducible nitric oxide synthase in human diseases," *Clinical & Experimental Immunology*, vol. 113, no. 2, pp. 147–156, 1998.
- [6] H. V. Dung, T. D. Cuong, N. M. Chinh et al., "Compounds from the aerial parts of *Piper bavinum* and their anti-cholinesterase activity," *Archives of Pharmacal Research*, vol. 38, no. 5, pp. 677–682, 2015.
- [7] J. Michel, R. E. Duarte, J. L. Bolton et al., "Medical potential of plants used by the Q'eqchi Maya of Livingston, Guatemala for the treatment of women's health complaints," *Journal of Ethnopharmacology*, vol. 114, no. 1, pp. 92–101, 2007.
- [8] B. Simpson and M. Ogorzaly, *Economic Botany: Plants in Our World*, McGraw-Hill, New York, NY, USA, 2nd edition, 1995.
- [9] B. Salehi, Z. A. Zakaria, R. Gyawali et al., "*Piper species*: a comprehensive review on their phytochemistry, biological activities and applications," *Molecules*, vol. 24, no. 7, p. 1364, 2019.
- [10] P. Chandra, R. Pandey, M. Srivastva, K. B. Rameshkumar, and B. Kumar, "Quantitative determination of chemical constituents of *Piper* spp. using UPLC-ESI-MS/MS," *Industrial Crops and Products*, vol. 76, pp. 967–976, 2015.
- [11] D. Lesueur, A. Bighelli, J. Casanova, T. M. Hoi, and T. H. Thai, "Composition of the essential oil of *Piper bavinum* C. DC. from Vietnam," *Journal of Essential Oil Research*, vol. 21, no. 1, pp. 16–18, 2009.
- [12] P. K. Agrawal, *Carbon-13 NMR of flavonoids. Studies in Organic Chemistry*, Vol. 39, Elsevier, Amsterdam, Netherlands, 1989.
- [13] Y. Zhang, S. Que, X. Yang, B. Wang, L. Qiao, and Y. Zhao, "Isolation and identification of metabolites from

- dihydromyricetin," *Magnetic Resonance in Chemistry*, vol. 45, no. 11, pp. 909–916, 2007.
- [14] J. Y. Zhang, Y. Chen, H. Q. Luo et al., "Recent update on the pharmacological effects and mechanisms of dihydromyricetin," *Frontiers in Pharmacology*, vol. 9, Article ID 1204, 2018.
- [15] G. Flamini, "Flavonoids and other compounds from the aerial parts of *Viola trusca*," *Chemistry & Biodiversity*, vol. 4, no. 2, pp. 139–144, 2007.
- [16] M. Skowrya, M. I. Calvo, M. G. Gallego, N. A. M. Azman, and M. P. Almajano, "Characterization of phytochemicals in petals of different colours from *Viola wittrockiana* gamms. and their correlation with antioxidant activity," *Journal of Agricultural Science*, vol. 6, no. 9, pp. 93–105, 2017.
- [17] P. V. Kiem, C. V. Minh, H. Huong et al., "Phenolic constituents with inhibitory activity against NFAT transcription from *Desmos chinensis*," *Archives of Pharmacal Research*, vol. 28, no. 12, pp. 1345–1349, 2005.
- [18] M. M. A. Hassan and J. S. Mossa, *Benzyl Benzoate, Analytical Profiles of Drug Substances*, Elsevier, Amsterdam, Netherlands, 1981.
- [19] J. Goutam, R. N. Kharwar, V. K. Tiwari, A. Mishra, and S. Singh, "Isolation and identification of antibacterial compounds isolated from endophytic fungus *emericella qau-drilineata* (RS-5)," *Natural Products Chemistry and Research*, vol. 4, no. 2, Article ID 1000205, 2016.
- [20] K. H. Kim, J. W. Choi, S. U. Choi et al., "The chemical constituents of *Piper kadsura* and their cytotoxic and anti-neuro inflammatory activities," *Journal of Enzyme Inhibition and Medicinal Chemistry*, vol. 26, no. 2, pp. 254–260, 2011.
- [21] C. C. M. P. Santos, M. S. Salvadori, V. G. Mota et al., "Antinociceptive and antioxidant activities of phytol in vivo and in vitro models," *Neuroscience Journal*, vol. 2013, Article ID 949452, 9 pages, 2013.
- [22] B. Pejic, V. Kojic, and G. Bogdanovic, "An insight into the cytotoxic activity of phytol at *in vitro* conditions," *Natural Product Research*, vol. 28, no. 22, pp. 2053–2056, 2014.
- [23] J. Moraes, R. N. Oliveira, J. P. Costa et al., "Phytol, a diterpene alcohol from chlorophyll, as a drug against neglected tropical disease schistosomiasis mansoni," *PLoS Neglected Tropical Diseases*, vol. 8, no. 1, Article ID e2617, 2014.
- [24] M. Shimizu and T. Tomoo, "Anti-inflammatory constituents of topically applied crude drugs. V. Constituents and anti-inflammatory effect of aoki, *aucuba japonica* THUNB," *Biological & Pharmaceutical Bulletin*, vol. 17, no. 5, pp. 665–667, 1994.
- [25] R. O. Silva, F. B. M. Sousa, S. R. B. Damasceno et al., "Phytol, a diterpene alcohol, inhibits the inflammatory response by reducing cytokine production and oxidative stress," *Fundamental & Clinical Pharmacology*, vol. 28, no. 4, pp. 455–464, 2014.
- [26] C. S. Philip, C. V. Nigel, and S. J. S. Monique, "Polyoxygenated cyclohexane derivatives and other constituents from *Kaempferia rotunda* L," *Phytochemistry*, vol. 68, no. 11, pp. 1579–1586, 2007.
- [27] T. Jaipetch, S. Hongthong, C. Kuhakarn et al., "Cytotoxic polyoxygenated cyclohexene derivatives from the aerial parts of *Uvaria cherrevensis*," *Fitoterapia*, vol. 137, Article ID 104182, 2019.
- [28] H. Kai, M. Baba, and T. Okuyama, "Two new megastigmanes from the leaves of *Cucumis sativus*," *Chemical & Pharmaceutical Bulletin*, vol. 55, no. 1, pp. 133–136, 2007.
- [29] K. Dewi, B. Widarto, P. Erawijantari, and W. Widowati, "In vitro study of *Myristica fragrans* seed (Nutmeg) ethanolic extract and quercetin compound as anti-inflammatory agent," *International Journal of Research in Medical Sciences*, vol. 3, no. 9, pp. 2303–2310, 2015.
- [30] K. S. Ahn, E. J. Noh, H. L. Zhao, S. H. Jung, S. S. Kang, and Y. S. Kim, "Inhibition of inducible nitric oxide synthase and cyclooxygenase II by Platycodon grandiflorum saponins via suppression of nuclear factor- κ B activation in RAW 264.7 cells," *Life Sciences*, vol. 76, no. 20, pp. 2315–2328, 2005.
- [31] T. D. Cuong, T. M. Hung, M. Na et al., "Inhibitory effect on NO production of phenolic compounds from *Myristica fragrans*," *Bioorganic & Medicinal Chemistry Letters*, vol. 21, no. 22, pp. 6884–6887, 2011.
- [32] T. D. Cuong, T. M. Hung, J. C. Kim et al., "Phenolic compounds from *Caesalpinia sappan* heartwood and their anti-inflammatory activity," *Journal of Natural Products*, vol. 75, no. 12, pp. 2069–2075, 2012.
- [33] J. K. Lee, N. R. Tae, J. J. Lee, T. Kim, and J. H. Lee, "Eupatolide inhibits lipopolysaccharide-induced COX-2 and iNOS expression in RAW264.7 cells by inducing proteasomal degradation of TRAF6," *European Journal of Pharmacology*, vol. 636, no. 1–3, pp. 173–180, 2010.
- [34] J. M. Olefsky and C. K. Glass, "Macrophages, inflammation, and insulin resistance," *Annual Review of Physiology*, vol. 72, no. 1, pp. 219–246, 2010.
- [35] C. Nathan, "Nitric oxide as a secretory product of mammalian cells," *The FASEB Journal*, vol. 6, no. 12, pp. 3051–3064, 1992.
- [36] Y. C. Liang, Y. T. Huang, and S. H. Tsai, "Suppression of inducible cyclooxygenase and inducible nitric oxide synthase by apigenin and related flavonoids in mouse macrophages," *Carcinogenesis*, vol. 20, no. 10, pp. 1945–1952, 1999.
- [37] H. K. Kim, B. S. Cheon, Y. H. Kim, S. Y. Kim, and H. P. Kim, "Effects of naturally occurring flavonoids on nitric oxide production in the macrophage cell line RAW 264.7 and their structure-activity relationships," *Biochemical Pharmacology*, vol. 58, no. 5, pp. 759–765, 1999.
- [38] X.-J. Lv, Y. Li, S.-G. Ma et al., "Bioactive megastigmane glucosides and monoterpenes from *Lyonia ovalifolia*," *Journal of Asian Natural Products Research*, vol. 21, no. 6, pp. 559–572, 2019.
- [39] T. T. T. Trang, T. D. Cuong, T. M. Hung et al., "Anti-inflammatory compounds from the aerial parts of *Aceriphyllum rossii*," *Chemical and Pharmaceutical Bulletin*, vol. 62, no. 2, pp. 185–190, 2014.
- [40] T. D. Cuong, T. M. Hung, J.-S. Lee, K.-Y. Weon, M. H. Woo, and B. S. Min, "Anti-inflammatory activity of phenolic compounds from the whole plant of *Scutellaria indica*," *Bioorganic & Medicinal Chemistry Letters*, vol. 25, no. 5, pp. 1129–1134, 2015.