

Review Article

Resistance Training Interventions for Lower Limb Tendinopathies: A Scoping Review of Resistance Training Reporting Content, Quality, and Scientific Implementation

Ian Burton ¹ and Aisling McCormack²

¹MSK Service, Fraserburgh Physiotherapy Department, Fraserburgh Hospital, NHS Grampian, Aberdeen, UK

²Aberdeen City Council, Aberdeen, UK

Correspondence should be addressed to Ian Burton; ianburton_10@hotmail.co.uk

Received 3 January 2022; Revised 26 January 2022; Accepted 11 February 2022; Published 10 March 2022

Academic Editor: Stephanie Georgina Dakin

Copyright © 2022 Ian Burton and Aisling McCormack. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The objectives of this scoping review were as follows: (1) to describe what exercises and intervention variables are used in resistance training interventions for lower limb tendinopathy, (2) to assess the completeness of reporting as assessed by the Consensus on Exercise Reporting Template (CERT) and the Toigo and Boutellier framework, and (3) to assess the implementation of scientific resistance training principles. We searched MEDLINE, CINAHL, AMED, Embase, SPORTDiscus, and Cochrane Library databases. Randomized controlled trials, cohort studies, case series, case reports, and observational studies that reported using resistance exercises for lower limb tendinopathies were considered for inclusion, with 194 studies meeting the inclusion criteria. Completeness of the reporting of exercise descriptors and programme variables was assessed by the CERT and the Toigo and Boutellier framework. Reporting of exercise descriptor items from the Toigo and Boutellier framework ranged from 0 to 13, with an average score of 9/13, with only 9 studies achieving a full 13/13. Reporting of items from the CERT ranged from 0 to 18, with an average score of 13/19. No study achieved a full 19/19; however, 8 achieved 18/19. Scoring for resistance training principles ranged from 1 to 10, with only 14 studies achieving 10/10. Eccentric heel-drops were the most common exercise (75 studies), followed by isotonic heel raises (38), and single-leg eccentric decline squats (27). The reporting of exercise descriptors and intervention content was high across studies, with most allowing exercise replication, particularly for Achilles and patellar tendinopathy. However, reporting for some tendinopathies and content items such as adherence was poor, limiting optimal translation to clinical practice.

1. Introduction

Tendinopathy represents a spectrum of potential changes to healthy tendons, leading to tendon damage and disease, with changes characterised by abnormal tendon composition and cellularity, ultimately leading to altered tendon microstructure [1]. In tendinopathic tendons, the normal arrangement of collagen fibres and organisation of tenocytes become altered by mechanical overuse, which leads to tendinopathy symptoms of pain, inflammation, swelling, and impaired physical function and performance [2]. Despite all healthy tendons having the capability to progress to tendinopathy, tendons of the lower limb may have an

increased risk for mechanical overload which causes tendon changes, such as disruption to the normal collagen matrix structure [3]. Although the aetiology of tendinopathy has yet to be fully elucidated, it is considered the result of a disrupted tendon healing process, with the hallmarks of collagen derangement, neovascularisation, altered tendon structure, and tissue calcification [4]. Tendinopathies account for up to 30% of all sports-related injuries with a range of extrinsic and intrinsic risk factors identified, suggesting that each individual pathogenesis of tendinopathy is multifactorial [5, 6]. The higher prevalence of patellar and Achilles tendinopathy found in athletes may be related to repetitive tendon microtrauma from repeated athletic

movements such as running, jumping, and landing [7]. Prevalence of Achilles and patellar tendinopathy have been reported to be as high as 23% and 45% in runners and jumping athletes, respectively. Plantar heel pain was reported to be found in up to 18% of running athletes in one cohort [8–10].

Despite a recent proliferation in clinical research investigating effectiveness of a range of treatment options for tendinopathy, it remains unclear which treatments are most effective, with exercise-based treatments such as resistance training currently being the most recommended due to consistent findings of efficacy [11, 12]. Common adjunctive treatments to exercise in clinical practice, which have conflicting findings of effectiveness include shockwave therapy, ultrasound, low-level laser therapy, manual therapy, and corticosteroid injections [13]. Isolated eccentric resistance training and heavy slow resistance training involving isotonic contractions have been shown to have favourable outcomes for common lower limb tendinopathies including Achilles tendinopathy, patellar tendinopathy, and plantar heel pain [14–17]. The high loads encountered during resistance training may stimulate tendon healing by counteracting structural tendon alteration, leading to reorganization and remodelling of collagen fibres, therefore improving the mechanical properties of tendons [18]. Despite positive outcomes, a limitation of some resistance training interventions in tendinopathy studies is that description, prescription, and progression of exercises and program variables may be poorly defined and reported, making translation to clinical practice difficult [19]. If the exercise dosage and training parameters prescribed clinically is insufficient, then, the mechanobiological stimulus may not be adequate to initiate tendon healing and positive outcomes from the intervention [20]. Despite the optimal dosage of resistance training for tendinopathy being unknown [21], research has shown an association between positive outcomes and higher exercise dosages in other musculoskeletal disorders [22].

In recent years, several guidelines or frameworks have been developed for reporting exercise interventions and specific exercise details within research studies in order to enhance reproducibility of exercise interventions and their translation to clinical practice [23]. The need to standardise reporting of components of exercise interventions has been highlighted by the development of the Consensus on Exercise Reporting Template (CERT), which advocates reporting detailed descriptions of exercises and their variables such as progression and tailoring, to allow clinical replication [24]. However, a limitation of the CERT is its omission of mechanobiological resistance training descriptors such as those included in the Toigo and Boutellier framework, such as rest intervals, time under tension, and relative load [25]. Holden et al. [26] recently highlighted how the poor reporting of exercise interventions in patellofemoral pain limits the clinical translation of exercise research findings in this population, with the authors recommending that future studies should use both the CERT and Toigo and Boutellier framework in conjunction as they report different aspects of exercise prescription and would therefore be

complementary. It is unclear if a similar issue exists within the interventional exercise literature in lower limb tendinopathies as no previous reviews have been conducted investigating the reporting of exercise descriptors using recommended frameworks. Although reporting of exercise interventions using the CERT has been recommended in tendinopathy to improve transparency and clinical translation, it is unclear if this recommendation has been widely adopted in research studies [27–29]. Both the CERT and Toigo and Boutellier framework are recommended templates and have been used in several review studies evaluating exercise descriptions and variables in rehabilitation for musculoskeletal disorders other than tendinopathy [30–32].

Although there has been a proliferation of clinical research examining resistance training treatment interventions for lower limb tendinopathies in recent years, it is unclear if these interventions have been sufficiently reported to allow clinical replication. Therefore, conducting a comprehensive scoping review of the current literature is warranted to investigate this question [29]. A search of MEDLINE, CINAHL, Cochrane Library, and PEDro identified no reviews with the objective of evaluating the reporting of resistance exercises and variables within interventions for lower limb tendinopathies using recommended frameworks such as the CERT or Toigo and Boutellier framework [33]. Therefore, the aims of this scoping review were to evaluate the reporting of exercise descriptors and programme variables used within resistance training interventions for treating lower limb tendinopathies. The scoping review was guided by addressing the following review questions on specific aspects of exercise reporting within lower limb tendinopathy resistance training interventions: (1) What exercises and program variables are used in resistance training interventions for lower limb tendinopathy? (2) How complete is the reporting of the exercise descriptors and programme variables as assessed by the CERT and the Toigo and Boutellier framework? (3) Do studies implement scientific resistance training principles?

2. Methods

Due to the exploratory nature of the research questions of this review, a scoping review was conducted as they are recommended for mapping key concepts, evidence gaps, and types of evidence within a particular field [34]. Scoping reviews can help guide future research and the possibility of conducting systematic reviews on the topic. As the aim of this review was to evaluate reporting across all study designs implementing resistance training for lower limb tendinopathies, rather than evaluate effectiveness, a scoping review was more appropriate than a systematic review. This scoping review was conducted in accordance with the Joanna Briggs Institute (JBI) methodology for scoping reviews [34]. The scoping review was reported in accordance with the Preferred Reporting Items for Systematic reviews and Meta-Analysis extension for Scoping Reviews (PRISMA-ScR) [34]. The findings allowed for dissemination of the parameters of research exercise interventions to clinical practitioners,

allowing increased likelihood of implementation in clinical practice [26]. The review also outlined future research and exercise reporting needs within lower limb tendinopathy resistance training interventions.

2.1. Study Identification. The inclusion criteria for this scoping review were guided by a modified PICO, which includes population, concept, and context (PCoCo), as recommended for scoping reviews [34]. The review included adults aged eighteen years or older with a diagnosis of a lower limb tendinopathy for any time duration. All lower limb tendinopathies were included, such as gluteal, hamstring, patellar, Achilles, tibialis posterior, and peroneal tendinopathy. Plantar heel pain was included as it is considered to have a similar pathophysiology to tendinopathy and should therefore be treated in accordance with other lower limb tendinopathies according to recent literature [19]. The concept of interest was resistance training for the treatment of lower limb tendinopathies, including any type or format of resistance training. The resistance training may be used as a first- or second-line intervention and may be delivered in isolation or combined with other treatments. Resistance training may be delivered across any healthcare or exercise setting, delivered by health or exercise professionals, in a supervised or unsupervised manner, using any methods for training progression and monitoring. This scoping review considered both experimental and quasi-experimental study designs including randomized controlled trials and nonrandomized controlled trials. In addition, prospective and retrospective cohort studies, case series, and case reports were considered for inclusion. Unpublished studies, reviews, or reports were not considered.

2.2. Search Strategy. A 3-step search strategy was implemented in this scoping review. It incorporated the following: (1) A limited search of MEDLINE and CINAHL using initial keywords, followed by analysis of the text words in the title or abstract and those used to describe articles to develop a full search strategy. (2) The full-search strategy was adapted to each database and applied to MEDLINE, CINAHL, AMED, Embase, SPORTDiscus, Cochrane Library (controlled trials and systematic reviews), and PEDro. The following trial registries were searched: ClinicalTrials.gov, ISRCTN, The Research Registry, EU-CTR (European Union Clinical Trials Registry), and ANZCTR (Australia and New Zealand Clinical Trials Registry). Databases were searched from inception to December 31, 2021. Although Stanish and Curwin [35] first published on the concept of eccentric resistance training for tendinopathy in 1986, it was only following the publication of the Alfredson protocol in 1998 [36] that resistance training became widespread in lower limb tendinopathy rehabilitation. Despite this, databases were searched from inception to ensure key articles and seminal studies on the topic published before 1998 were not omitted, which may influence overall findings. (3) For each article located in steps 1 and 2, a search of cited and citing articles using Scopus and hand-searching, where necessary,

was conducted. Studies published in a language other than English were only included if a translation was available as translation services were not available to the authors.

2.3. Study Selection. Following the search, all identified citations were collated and uploaded into RefWorks, with duplicates being removed. Titles and abstracts were then screened by two independent reviewers (IB and AM) for assessment against the inclusion criteria for the review. Potentially relevant studies were retrieved in full, and their citation details imported into Covidence (Veritas Health Innovation, Melbourne, Australia). Two independent reviewers (IB and AM) then assessed the full text of selected citations in detail against the inclusion criteria. Any disagreements that arose between the reviewers at each stage of the study selection process were resolved through discussion or by input from a third reviewer. The results of the search were reported in accordance with the PRISMA-ScR (Figure 1) [34].

2.4. Data Extraction. Data were extracted from sources included in the scoping review by one reviewer, with independent data extraction by a second reviewer for 10% of studies. The data were extracted using an excel spreadsheet developed specifically by the reviewers for each source type. The data extracted included specific details regarding the population, concept, context, study methods, and key findings relevant to the review questions. Any disagreements that arose between the reviewers were resolved through discussion. The data extracted included dimensions such as authors, year of publication, study type, purpose, population and sample size, methods, details of resistance training intervention, specific exercises, and outcome measures used. Details of the resistance training interventions included setting, mode of delivery, type, dosage, and methods used to progress and adjust the training stimulus. The contents and variables of the specific resistance training exercises were extracted using the 13-item Toigo and Boutellier framework for exercise mechanobiological determinants and included parameters such as repetitions, load magnitude, and time under tension. General information from the resistance training interventions such as exercise supervision and delivery methods were extracted using the CERT tool. An evaluation of the implementation of scientific resistance training principles was also conducted, by extracting data on the principles of specificity, overload, progression, individualisation, and adherence. The definitions and scoring criteria for these principles are provided in Tables 1 and 2. In accordance with guidance on conducting scoping reviews, critical appraisal was not conducted [34].

2.5. Data Analysis. The extracted data were analysed using descriptive statistics, with findings presented in tabular form as tables and figures, in a manner that aligns with the objective of this scoping review. A narrative summary accompanies the tabulated results and describes how the results relate to the review objective and questions.




FIGURE 1: PRISMA study flow diagram.

TABLE 1: Resistance training principles and training intervention criteria assessed.

Principle	Criteria for this review
Specificity: training and desired adaptations should be specific to the tendinopathy and relevant to desired outcomes	Appropriate population targeted and intervention designed to improve primary outcome
Progression: to allow for continuous adaptations, resistance or load must be increased providing a greater stress to the body	Training intervention was stated to be progressive with gradual increases in frequency, sets, repetitions, resistance, or loading throughout intervention
Overload: for the intervention to improve strength, greater than normal stress and training volume must occur above current training levels	Interventions included baseline strength testing or rationale that intervention was of sufficient intensity and volume relative to baseline capacity
Individualisation: training is tailored to the individual to allow for consideration of individual factors and training response	Training intervention considered methods to individually tailor exercises stimulus based on an individual's own factors or training response
Component of training	Description
Frequency	How many times per week or day
Intensity	Measurement methods: RM, %RM, RPE, pain level
Time	Duration of session
Sets	How many sets of each exercise
Repetitions	How many repetitions of each exercise or target number of repetitions
Exercise selection	Outline and description of specific exercises used in intervention
Adherence	Was adherence to the training intervention monitored and reported?

TABLE 2: Application and reporting of key training principles.

Principle/ criterion	Description	Score
Specificity	Design: have the authors designed the intervention to achieve desired outcomes? 1/10	Reporting: have the authors adequately described the intervention specificity? 1/10 2/10
Overload	Design: have the authors appropriately manipulated training variables to achieve desired outcomes? 1/10	Reporting: have the authors adequately described the intervention training variables? 1/10 2/10
Progression	Design: have the authors appropriately manipulated training variables to adequately progress the intervention? 1/10	Reporting: have the authors adequately described how intervention progression was achieved and assured? 1/10 2/10
Individualisation	Design: have the authors appropriately manipulated training variables to tailor the intervention adequately individually? 1/10	Reporting: have the authors adequately described how individually tailoring the intervention was achieved and assured? 1/10 2/10
Adherence	Design: have the authors appropriately designed and described methods for monitoring adherence? 1/10	Reporting: have the authors adequately reported individual adherence to training and training dose achieved? 1/10 2/10

Completeness of information regarding the resistance training interventions is presented as the number of complete items of the CERT, Toigo and Boutellier framework, and resistance training principles for each study. Resistance training exercises were categorised as eccentric, concentric, isometric, isotonic, heavy slow resistance training (HSRT), low-load blood flow restricted training (BFRT), isoinertial, manually resisted, hip strength, and general strength exercise.

3. Results

3.1. Study Characteristics. A total of 194 studies were included (Table 3), of which 109 were randomized controlled trials (RCTs), 32 were cohort studies, 15 were case series, 26 were case reports, with 12 being other designs, quasi-experimental (4), before-after design (2), case-control (4), and observational (2). The publication year ranged from 1989 to 2021, with 42 of the included studies being published since the year 2020, and 62 since 2019. The tendinopathy most frequently treated was Achilles (99), followed by patellar (58), plantar heel pain (16), gluteal (8), posterior tibial (6), hamstring (5), peroneal (1), extensor hallucis longus (1), and iliopsoas (1). The sample sizes of included studies ranged from 1 to 204 patients. The duration of included resistance training interventions ranged from a single session to 32 weeks, with 12 being the most common, implemented in 122 studies (63%). All studies evaluated intervention outcomes for at least one of pain and function, with the vast majority evaluating both outcomes. Pain was assessed by a visual analogue scale (VAS) in 90 (46%) studies, and pain numeric rating scale (NRS-P) in 21 (11%) studies. Pain and function were assessed by the Victorian Institute of Sport Assessment-Achilles (VISA-A) in 66 (34%) studies, Victorian Institute of Sport Assessment-Patellar (VISA-P) in 45 (23%) studies, Victorian Institute of Sport Assessment-Gluteal (VISA-G) in 5 (3%) studies, Foot Function Index (FFI) in 10 (5%) studies, and Lower Extremity Function Scale (LEFS) in 10 (5%) studies. Six (3%) studies also assessed the tendon structure through ultrasonography and four (2%) studies used quality of life outcome measures.

3.2. Content and Completeness of Exercise Description. Eccentric training was the most common type of resistance training, implemented in 130 (67%) studies, followed by general strength exercise in 35 (18%) studies, HSRT in 15 (8%) studies, isometric in 14 (7%) studies, isotonic in 11 (6%) studies, concentric in 6 (3%) studies, isoinertial in 4 (2%) studies, hip strength exercises in 4 (2%) studies, BFRT in 2 (1%) studies and manually resisted exercise in 1 (0.5%) study. In terms of specific resistance training exercises, the Alfredson eccentric heel-drop was the most common exercise with 75 (39%) studies implementing it, followed by isotonic heel raises in 38 (20%) studies, single-leg eccentric decline squats in 27 (14%) studies, knee extension in 16 (8%) studies, leg press in 13 (7%) studies, hip abduction in 10 (5%) studies, hip bridging in 9 (5%) studies, deadlifts in 8 (4%) studies, ankle inversion in 7 (4%) studies, plyometric jump exercises in 6 (3%) studies, and lunges in 5 (3%) studies.

The number of items that described the Toigo and Boutellier framework exercise descriptors ranged from 0 to 13 out of a possible 13, with an average score across the 194 studies of 9/13. Only 9 [19, 50, 79, 155, 175, 177, 201, 217, 221] (5%) studies achieved a full 13/13 for reporting items from the framework, with three of these from the same author [19, 217, 221]. Overall reporting across all studies for each item is reported in Figure 2. Only 3 items were reported by less than 80% of studies, rest between sets 47 (24%), time under tension 34 (18%), and volitional muscular failure 15 (8%). The item with the highest percentage of reporting at 97% was the contraction mode of the exercise employed in the intervention, reported in 188 studies. Of the 19 items included in the CERT, reporting among included studies ranged from 0 to 18, with an average score of 13/19. No study achieved a full score of 19, but 8 [15, 50, 79, 110, 111, 137, 177, 221] (4%) studies achieved a high score of 18/19. Four of these studies [50, 79, 177, 221] also achieved a full score of 13/13 for reporting exercise descriptors. Overall reporting for each item is presented in Figure 3. Most items were well-reported across studies, with seven items being reported in less than 70% of studies, progression rules 133 (69%), tailored how 134 (69%), adherence measures 94 (49%), exercise delivered

TABLE 3: Characteristics and reporting scores of the 194 included studies.

Author	Study design	Tendinopathy	Resistance training type	Resistance training exercise	TBF/13	CERT/19	RTP/10
Abdelkader et al. [37]	RCT	Achilles	ECCT	Alfredson heel-drop	11	11	2
Alfredson et al. [36]	RCT	Achilles	ECCT	Alfredson heel-drop	10	14	7
Balius et al. [38]	RCT	Achilles	ECCT	Alfredson heel-drop	8	10	4
Bell et al. [39]	RCT	Achilles	ECCT	Alfredson heel-drop	7	14	6
Beyer et al. [14]	RCT	Achilles	HSRT, ECCT	Heel raises	12	17	9
Boesen et al. [40]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	8
Brown et al. [41]	RCT	Achilles	ECCT	Alfredson heel-drop	1	1	1
Chester et al. [42]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	7
Choudhary et al. [43]	RCT	Achilles	ECCT	NR	8	12	7
De jonge et al. [44]	RCT	Achilles	ECCT	Alfredson heel-drop	10	14	8
De jonge et al. [45]	RCT	Achilles	ECCT	Alfredson heel-drop	6	11	5
De Vos et al. [46]	RCT	Achilles	ECCT	Alfredson heel-drop	6	11	5
De vos et al. [47]	RCT	Achilles	ECCT	Alfredson heel-drop	10	16	9
Gatz et al. [48]	RCT	Achilles	ECCT, ECCT + ISOM	Alfredson heel-drop	10	15	8
Habets et al. [49]	RCT	Achilles	ECCT, CONCT-ECCT	Alfredson heel-drop, heel raises	10	16	10
Hasani et al. [50]	RCT	Achilles	ISOT	Heel raises	13	18	9
Herrington et al. [51]	RCT	Achilles	ECCT	Alfredson heel-drop	10	16	8
Horstmann et al. [52]	RCT	Achilles	ECCT	Alfredson heel-drop	11	15	7
Kearney et al. [53]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	7
Kedia et al. [54]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	8
Knobloch et al. [55]	RCT	Achilles	ECCT	Alfredson heel-drop	10	11	2
Kozsalinski et al. [56]	RCT	Achilles	ECCT	Alfredson heel-drop	7	10	2
Mafi et al. [57]	RCT	Achilles	ECCT, CONCT	Alfredson heel-drop	10	15	7
Mansur et al. [58]	RCT	Achilles	ECCT	Alfredson heel-drop	10	12	4
McCormack et al. [59]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	5
Munteanu et al. [60]	RCT	Achilles	ECCT	Alfredson heel-drop	10	16	10
Niesen-Vertommen [61]	RCT	Achilles	ECCT, CONCT	Heel raises	10	17	9
Norregaard et al. [62]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	9
Notarnicola et al. [63]	RCT	Achilles	ECCT	NR	3	3	2
Pearson et al. [64]	RCT	Achilles	ECCT	Alfredson heel-drop	1	5	7
Petersen et al. [65]	RCT	Achilles	ECCT	Alfredson heel-drop	10	16	8
Praet et al. [66]	RCT	Achilles	ECCT	Alfredson heel-drop	10	17	10
Rabusin et al. [67]	RCT	Achilles	ECCT	Alfredson heel-drop	10	17	10
Romero-morales [68]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	8
Romero-morales [69]	RCT	Achilles	ECCT	Alfredson heel-drop	10	15	8
Rompe et al. [70]	RCT	Achilles	ECCT	Alfredson heel-drop	11	17	9
Rompe et al. [71]	RCT	Achilles	ECCT	Alfredson heel-drop	11	17	8
Rompe et al. [72]	RCT	Achilles	ECCT	Alfredson heel-drop	10	16	8
Roos et al. [73]	RCT	Achilles	ECCT	Alfredson heel-drop	10	16	9
Silbernagel et al. [74]	RCT	Achilles	General strength EX	Heel raises, plyometric heel raises	10	16	8
Silbernagel et al. [75]	RCT	Achilles	ECCT	Heel raises, plyometric heel raises	10	15	8
Solomons et al. [76]	RCT	Achilles	General strength EX	NR	1	11	6
Stefansson et al. [77]	RCT	Achilles	ECCT	Alfredson heel-drop	10	14	8
Stergioulas et al. [78]	RCT	Achilles	ECCT	Heel raises	11	16	10

TABLE 3: Continued.

Author	Study design	Tendinopathy	Resistance training type	Resistance training exercise	TBF/13	CERT/19	RTP/10
Stevens and Tan [79]	RCT	Achilles	ECCT	Alfredson heel-drop	13	18	9
Tumilty et al. [80]	RCT	Achilles	ECCT	Alfredson heel-drop	10	17	9
Tumilty et al. [81]	RCT	Achilles	ECCT	Alfredson heel-drop	10	17	9
Van der vlist et al. [82]	RCT	Achilles	ECCT	Heel raises, plyometric heel raises	12	17	9
Yelland et al. [83]	RCT	Achilles	ECCT	Alfredson heel-drop	10	17	8
Yu et al. [84]	RCT	Achilles	ECCT, CONCT	Heel raises, alfredson heel-drop	10	15	8
Zhang et al. [85]	RCT	Achilles	ECCT	Alfredson heel-drop	10	14	8
Kanniappan et al. [86]	Quasi	Achilles	ECCT, ISOM	Alfredson heel-drop, static plantar flexion	11	11	4
Stasinopoulos et al. [89]	Quasi	Achilles	ECCT	Alfredson heel-drop, heel raises	11	17	8
Van der vlist et al. [88]	Quasi	Achilles	ISOM, ISOT	Heel raises	12	17	8
De vos et al. [89]	OBS	Achilles	ECCT	Alfredson heel-drop	11	14	8
Masood et al. [90]	L-CC	Achilles	ECCT	Alfredson heel-drop	10	16	10
Abate et al. [91]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	16	9
Alfredson et al. [95]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	14	8
Alfredson et al. [96]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	14	7
Angermann et al. [94]	Cohort	Achilles	General strength EX	Heel raises	10	15	9
De jonge et al. [95]	Cohort	Achilles	ECCT	Alfredson heel-drop	3	2	2
Fahlstrom et al. [96]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	14	7
Jonsson et al. [97]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	14	8
Lagas et al. [98]	Cohort	Achilles	ECCT	NR	1	0	2
Maffulli et al. [103]	Cohort	Achilles	ECCT	Alfredson heel-drop	11	16	9
Mansur et al. [104]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	14	8
Mantovani et al. [110]	Cohort	Achilles	ISOM	Heel raises	12	18	8
O'Neill et al. [102]	Cohort	Achilles	ISOM	Dynamometer heel raise	10	11	4
Ohberg et al. [103]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	14	7
Ooi et al. [104]	Cohort	Achilles	ECCT	Alfredson heel-drop	1	2	2
Robinson et al. [110]	Cohort	Achilles	ECCT	Heel raises	8	12	2
Sancho et al. [111]	Cohort	Achilles	General strength EX	Heel raises, hip abduction and extension, hops, jumps	10	18	9
Sayana et al. [112]	Cohort	Achilles	ECCT	Alfredson heel-drop	11	16	9
Shalabi et al. [113]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	16	8
Von Wehren et al. [109]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	11	4
Wei et al. [110]	Cohort	Achilles	ECCT	Alfredson heel-drop	10	15	9
Wetke et al. [111]	Cohort	Achilles	ISOT	Heel raises	10	16	9
Wheeler et al. [112]	Cohort	Achilles	ECCT	NR	1	3	2
Langberg et al. [113]	CC	Achilles	ECCT	Alfredson heel-drop	10	15	10
Park et al. [114]	CC	Achilles	ECCT	Alfredson heel-drop	11	14	8
Ram et al. [115]	CC	Achilles	ECCT	Alfredson heel-drop	10	13	8
Benito et al. [116]	C. series	Achilles	ECCT	Alfredson heel-drop	10	13	6
Deans et al. [117]	C. series	Achilles	General strength EX	NR	1	4	2
Jayaseelan et al. [118]	C. series	Achilles	ECCT	Alfredson heel-drop	10	12	3
Pavone et al. [119]	C. series	Achilles	ECCT	NR	1	4	2
Silbernagel et al. [120]	C. series	Achilles	ECCT, plyometric EX	Heel raises, plyometric heel raises	10	14	7
Syverston et al. [121]	C. series	Achilles	ECCT	Alfredson heel-drop	10	11	4
Wheeler et al. [122]	C. series	Achilles	General strength EX	NR	1	5	2
Borda et al. [123]	C. report	Achilles	ECCT	Alfredson heel-drop	10	11	2
Cuddeford et al. [124]	C. report	Achilles	ECCT	Heel-drop on leg press	8	15	8

TABLE 3: Continued.

Author	Study design	Tendinopathy	Resistance training type	Resistance training exercise	TBF/ 13	CERT/ 19	RTP/ 10
Eckenrode et al. [125]	C. report	Achilles	ECCT	Alfredson heel-drop, bridging, hip abduction, squat	12	14	7
Francis et al. [126]	C. report	Achilles	ECCT	Alfredson heel-drop	10	15	8
Greene et al. [127]	C. report	Achilles	General strength EX	Heel raises, squats, leg pulls	8	14	7
McCormack et al. [128]	C. report	Achilles	ECCT	Alfredson heel-drop	10	14	7
Papa et al. [129]	C. report	Achilles	ECCT	Alfredson heel-drop	10	13	5
Ross et al. [130]	C. report	Achilles	General strength EX	Heel raises, plantarflexion, dorsiflexion, plyometric	10	16	8
Thompson et al. [131]	C. report	Achilles	General strength EX	Squats, pelvic thrust, deadlift, heel raises, band walks, lunges	11	14	7
Gardin et al. [132]	BAD	Achilles	ECCT	Alfredson heel-drop	10	10	4
Croisier et al. [133]	Cohort	Achilles, patellar	Isokinetic ECCT	Dynamometer heel raise, knee extension	10	16	8
Pinkelman et al. [134]	C. report	EHL	Manually resisted EX	Great toe extension	8	14	7
Clifford et al. [15]	RCT	Gluteal	ISOM, ISOT	Isometric and isotonic hip abduction exercises	12	18	9
Cowan et al. [135]	RCT	Gluteal	General strength EX	Isometric and isotonic hip exercises	10	17	9
Ganderton et al. [136]	RCT	Gluteal	General strength EX	Isometric and isotonic hip exercises	10	17	9
Mellor et al. [137]	RCT	Gluteal	General strength EX	Isometric and isotonic hip exercises	11	18	10
Ramon et al. [138]	RCT	Gluteal	General strength EX	Bridging, hip abduction and extension	10	12	2
Thompson et al. [139]	RCT	Gluteal	ECCT	Lunges, squats	6	10	5
Wheeler et al. [140]	RCT	Gluteal	General strength EX	Hip abduction, bridging, clams	7	13	7
Van Rooy et al. [141]	C. report	Gluteal	ECCT	Hip abduction, lunges bridging	10	12	4
Cacchio et al. [142]	RCT	Hamstring	General strength EX	Leg curls, lunge, squat, CM jumps, deadlift, hip strength exercises	8	7	4
Cushman et al. [143]	C. report	Hamstring	ECCT	Hip extension	10	12	4
Jayaseelan et al. [144]	C. report	Hamstring	General strength EX	Leg curl, deadlift, bridging, hip abduction	10	14	7
Krueger et al. [145]	C. report	Hamstring	HSRT	Squat, deadlift, hip thrust, leg curl, reverse lunge	11	15	8
McCormack et al. [146]	C. report	Hamstring	ECCT	Leg curl, hip extension, bridging, lunges Nordics, deadlift	10	13	4
Rauseo et al. [147]	C. report	Iliopsoas	ECCT	Hip flexion, bridging, squats, deadlift	10	15	8
Alvarez et al. [148]	RCT	P. tibial	General strength EX	Heel raises, plantarflexion, adduction, inversion	10	17	9
Houck et al. [149]	RCT	P. tibial	General strength EX	Heel raises, plantarflexion, adduction, inversion	11	17	9
Kulig et al. [150]	RCT	P. tibial	Isokinetic ECCT, CONCT	Resisted adduction with plantarflexion	12	17	10
Kulig et al. [151]	C. series	P. tibial	ECCT	Resisted adduction with plantarflexion	11	15	10
Robinson et al. [152]	C. series	P. tibial	General strength EX	Heel raises, short foot	8	11	2
Patla et al. [153]	C. report	P. tibial	General strength EX	Heel raises, pronation and supination	8	13	3
Abat et al. [154]	RCT	Patellar	ECCT	DSL squat	9	8	2
Agergaard et al. [155]	RCT	Patellar	HSRT, M-HSRT	Leg press and extension	13	17	10
Bahr et al. [156]	RCT	Patellar	ECCT	DSL squat	11	14	8
Biernat et al. [157]	RCT	Patellar	ECCT	DSL squat	10	14	7

TABLE 3: Continued.

Author	Study design	Tendinopathy	Resistance training type	Resistance training exercise	TBF/13	CERT/19	RTP/10
Breda et al. [158]	RCT	Patellar	HSRT, ECCT	DSL squat, leg press, knee extension, hip strength exercises	10	17	9
Cannell et al. [159]	RCT	Patellar	ECCT, ISOT	Drop squat, knee extension and curl	11	14	8
Da Cunha et al. [160]	RCT	Patellar	ECCT	DSL squat	10	14	8
Dimitrios et al. [161]	RCT	Patellar	ECCT	DSL squat	11	17	8
Dragoo et al. [162]	RCT	Patellar	ECCT	NR	1	5	2
Frohm et al. [163]	RCT	Patellar	ECCT	DSL squat	11	14	8
Holden et al. [164]	RCT	Patellar	ISOM, dynamic EX	Knee extension	12	13	5
Jensen et al. [165]	RCT	Patellar	Isokinetic ECCT	Dynamometer heel raise	11	16	8
Jonsson et al. [166]	RCT	Patellar	ECCT, CONCT	DSL squat	10	15	7
Kaux et al. [167]	RCT	Patellar	ECCT	Wall squat	11	13	5
Kongsgaard et al. [16]	RCT	Patellar	HSRT, ECCT	DSL squat, hack squat, leg press, squat	12	17	9
Lee et al. [168]	RCT	Patellar	ECCT	DSL squat	11	14	9
Lopez-royo et al. [169]	RCT	Patellar	ECCT	DSL squat	10	14	7
MacDonald et al. [170]	RCT	Patellar	ECCT, ECCT + hip	DSL squat, isotonic hip exercises	10	16	8
Olesen et al. [171]	RCT	Patellar	HSRT	Squat, leg press, knee extension, hack squat	10	14	7
Pietrosimone et al. [172]	RCT	Patellar	ISOM	Knee extension	12	12	4
Rio et al. [173]	RCT	Patellar	ISOM, ISOT	Knee extension	12	13	5
Rio et al. [174]	RCT	Patellar	ISOM, ISOT	Knee extension	12	16	9
Ruffino et al. [175]	RCT	Patellar	HSRT, isoinertial	Squat, leg press, knee extension, hack squat	13	17	9
Scott et al. [176]	RCT	Patellar	HSRT	NR	1	5	2
Sprague et al. [177]	RCT	Patellar	HSRT	Squat, leg press, knee extension, hack squat	13	18	9
Stasinopolous et al. [178]	RCT	Patellar	ECCT	DSL squat	10	14	7
Steunebrink et al. [179]	RCT	Patellar	ECCT	Alfredson heel-drop	10	15	10
Thijs et al. [180]	RCT	Patellar	ECCT	DSL squat	10	16	7
Van Ark et al. [181]	RCT	Patellar	ISOT, ISOM	Knee extension	12	16	8
Van Ark et al. [182]	RCT	Patellar	ISOM, ISOT	Knee extension	12	14	8
Van der Worp et al. [183]	RCT	Patellar	ECCT	DSL squat	9	16	8
Visnes et al. [184]	RCT	Patellar	ECCT	DSL squat	10	15	9
Wang et al. [185]	RCT	Patellar	ECCT	Quadriceps and hamstring strengthening	1	3	2
Warden et al. [186]	RCT	Patellar	ECCT	DSL squat	10	17	9
Young et al. [187]	RCT	Patellar	ECCT	DSL squat	10	16	10
Vander doelen et al. [188]	R chart	Patellar	General strength EX	Knee extension, leg press, squat, hack squat	11	14	5
Purdam et al. [189]	Quasi	Patellar	ECCT	DSL squat, squat	10	14	7
Abat et al. [190]	Cohort	Patellar	Isoinertial ECCT	NR	10	12	5
Abat et al. [191]	Cohort	Patellar	Isoinertial ECCT	Leg press	10	10	4
Basas et al. [192]	Cohort	Patellar	ISOT	NR	8	14	7
Kaux et al. [193]	Cohort	Patellar	ECCT	NR	3	4	2
Kaux et al. [194]	Cohort	Patellar	ECCT, ISOM	DSL squat	11	16	9
Kongsgaard et al. [195]	Cohort	Patellar	HSRT	Knee extension, leg press, hack squat, squat	12	15	7
Panni et al. [196]	Cohort	Patellar	General strength EX	NR	1	2	2
Bianco et al. [197]	C. series	Patellar	General strength EX	DSL squat, squat, drop squat, SL squat, jump downs, mini squat	8	14	5
Morton et al. [198]	C. series	Patellar	ECCT	DSL squat	1	5	2

TABLE 3: Continued.

Author	Study design	Tendinopathy	Resistance training type	Resistance training exercise	TBF/ 13	CERT/ 19	RTP/ 10
Munoz fernandez [199]	C. series	Patellar	General strength EX	Clams, bridging, DSL squat, squat, deadlift, short foot, hip abduction, pelvic drops	10	12	4
Romero-rodriguez et al. [200]	C. series	Patellar	Isoinertial ECCT	Flywheel	12	15	10
Skovlund et al. [201]	C. series	Patellar	LL-BFRT	Knee extension, leg press	13	17	9
Van ark et al. [202]	C. series	Patellar	General strength EX	Heel raises, squats, hip abduction, SL squat, lunges, step-downs, bridging, jumps	11	16	8
Cuddeford et al. [203]	C. report	Patellar	LL-BFRT	DSL squat, leg press	12	15	8
Dumont et al. [204]	C. report	Patellar	ECCT	Drop squats	10	16	9
Goldman et al. [205]	C. report	Patellar	ECCT	DSL squats, leg press, knee extension, leg curls, step-downs, heel taps	11	14	7
McCreesh et al. [206]	C. report	Patellar	ECCT	DSL squat	10	13	5
Rowan et al. [207]	C. report	Patellar	ECCT	DSL squats	8	10	2
Silva et al. [208]	C. report	Patellar	Hip strength EX	Hip extension, birddog, deadlift, drop jumps	11	14	8
Morgan et al. [209]	BAD	Patellar	General strength EX	NR	4	6	6
Hensley et al. [210]	C. report	Peroneal	General strength EX	Heel raises, inversion and eversion	10	13	5
Chesterton et al. [211]	RCT	Plantar	General strength EX	Foot, calf and hip strength exercises	2	14	6
Cil et al. [212]	RCT	Plantar	General strength EX	Foot, ankle and hip exercises	9	10	5
Johannsen et al. [213]	RCT	Plantar	General strength EX	Heel raises, inversion	4	8	4
Johannsen et al. [214]	RCT	Plantar	HSRT	Heel raises, inversion	3	5	2
Kamonseki et al. [215]	RCT	Plantar	Foot, hip strength EX	Foot, ankle and hip exercises	10	13	5
Rasenberg et al. [216]	RCT	Plantar	General strength EX	NR	1	0	3
Rathleff et al. [17]	RCT	Plantar	HSRT	Heel raises	11	14	5
Riel et al. [19]	RCT	Plantar	HSRT	Heel raises	13	17	9
Riel et al. [217]	RCT	Plantar	ISOM, ISOT	Heel raises	13	14	7
Ryan et al. [218]	RCT	Plantar	General strength EX	Inversion and eversion	6	11	3
Thong-On et al. [219]	RCT	Plantar	General strength EX	Heel raises, inversion and eversion, toe curls	10	17	9
Wheeler et al. [220]	RCT	Plantar	General strength EX	Heel raises, foot strength exercises	0	8	2
Riel et al. [221]	Cohort	Plantar	HSRT	Heel raise	13	18	9
Wheeler et al. [222]	Cohort	Plantar	General strength EX	Heel raises, IFM strength	2	5	2
Dos Santos et al. [223]	C. report	Plantar	Hip strength EX	Hip abduction, extension, adduction and flexion	5	9	2
Lee et al. [224]	C. report	Plantar	Hip strength EX	NR	1	4	2

CERT: Consensus on Exercise Reporting Template; TBF: Toigo and Boutellier Framework; RTP: resistance training principles; ECCT: eccentric training; CONCT: concentric training; ISOM: isometric; ISOT: isotonic; EX: exercise; LL-BFRT: low-load blood flow restriction training; HSRT: heavy slow resistance training; C. series: case series; C. report: case report; CC: case control; BAD: before-after design; L-CC: longitudinal case-control; OBS: observational; QUASI: quasiexperimental; P.tibial: Posterior tibial; NR: not reported; DSL: decline single-leg; R chart: retrospective chart review; FHL: flexor hallucis longus.




FIGURE 2: Percentage of studies (out of 194) with complete reporting for each item of the Toigo and Boutellier framework.

as planned 70 (36%), adverse events 55 (28%), fidelity measured 11 (6%), and motivation strategies 4 (2%), with the latter two items particularly poorly reported across the studies. Previous studies assessing the completeness of CERT items in musculoskeletal rehabilitation determined that reporting completeness of items could be regarded as high (>75%), moderate (60 to 74%), or low (<60%) [225, 226]. Based on this classification, 10 items can be rated as high, 4 as moderate, and 5 as low.

3.3. Application of Resistance Training Principles. An evaluation of the implementation of scientific resistance training principles was conducted by evaluating the design and reporting of the key principles of specificity, overload, progression, individualisation, and adherence (Table 1). One point each was given for the design and reporting of each of the 5 principles, with a maximum score of 10/10 available. The scoring system was based on scales used in previous reviews with the same objective [11, 227, 228]. Scoring for resistance training principles ranged from 1 to 10 across the 194 studies, with only 14 studies achieving a full score of 10/10 [49, 60, 66, 67, 78, 90, 113, 137, 150, 151, 155, 179, 187, 200]. Only one study did not implement and report the principle of specificity, whereas 193 (99%) studies implemented specificity by targeting the prescribed resistance training to the specific tendinopathy with the aim to improve pain and function. The principle of overload was not adequately implemented or reported in 45 studies, with 149 (77%) studies implementing overload by progressively increasing training resistance throughout the intervention. The principle of progression was not adequately implemented or reported in 57 studies, with 137 (71%) studies implementing progression, most commonly by increasing resistance through small increases in external weight. However, only 35 (18%) studies accurately

reported the exact amount of weight implemented in progression increments. Incremental increases in resistance ranged from 0.9 to 5 kg, with 5 kg being the most common, implemented in 27 (14%) studies. The principle of individualisation was not adequately implemented or reported in 59 studies, with 135 (70%) studies implementing individualisation, most commonly by adjusting training resistance based on pain response as implemented in 118 (61%) studies. Other reported methods for individually tailored training included exercise technique in 6 (3%) studies, as much volume as possible in 3 (2%) studies, increasing exercise difficulty in 8 (4%) studies, and level of fatigue in 7 (4%) studies, typically measured by using rating of perceived exertion (RPE) scales. The principle of adherence was not adequately implemented or reported in 68 studies, with 126 (65%) studies implementing adherence, most commonly by using an individual exercise diary as reported in 72 (37%) studies. However, only 46 (24%) studies reported the percentage of participants who achieved what each study authors determined to be an acceptable level of resistance training adherence, which ranged from 40 to 100%. Although the median value of adherence was reasonably high at 70%, it is concerning that only 24% of included studies reported adherence levels.

4. Discussion

The main findings from this scoping review are that the description and reporting of resistance training exercises and intervention parameters used in the rehabilitation of lower limb tendinopathies were generally well reported across studies. However, some reporting areas of weakness were identified, with items such as adherence, fidelity, and specific exercise progression parameters of interventions, being poorly reported. A broad range of resistance training




FIGURE 3: Percentage of studies (out of 194) with complete reporting for each item of the Consensus on Exercise Reporting Template (CERT).

types were implemented with eccentric training the most common at 67%, with Alfredson eccentric heel-drops (39%), isotonic heel raises (20%), and eccentric decline single-leg squats (14%) being the most implemented exercises. Most studies included sufficient information on exercise dosage (load, repetitions, sets, and frequency) to allow replication of the exercises in both research and clinical settings. However, not all studies provided sufficient detail to allow replication, suggesting there is room for improvement in future research, with items such as adherence and fidelity of interventions being poorly reported. Whereas the scientific resistance training principles of specificity and overload were well implemented and reported throughout the studies, the principals of progression, individualisation, and adherence had poorer reporting, preventing complete clinical replication of these principles in some studies. Despite these issues, the overall moderate to high quality of reporting was better for lower limb tendinopathies than for other musculoskeletal conditions as assessed in other reviews applying the CERT. For example, the quality of exercise content

reporting has been found to be considerably lower in exercise rehabilitation interventions for hamstring strains [225], groin injury [226], Achilles tendon ruptures [30], rotator cuff disorders [32], knee osteoarthritis [229, 230], patellofemoral pain [26], knee injuries [31], fibromyalgia [231], juvenile arthritis [232], hand osteoarthritis [233], pelvic floor dysfunction [234, 235], low back pain [236, 237], ACL injury [238], and femoral-acetabular impingement [239].

The only other review reporting an overall moderate–high quality for CERT reporting like this review, was for hip osteoarthritis, which reported an average CERT score of 13/19 [240]. Item 8 of the CERT relates to describing exercises to a level that allows replication, which was met by 85% of studies included in this review. In comparison, reporting of this item was much lower in the reviews for hamstring strains [225] (43%), knee osteoarthritis [230] (26%), rotator cuff disorders [32] (29%), groin injuries [226] (15%), and Achilles tendon ruptures [30] (26%). These findings highlight the comparative quality of exercise reporting for lower limb tendinopathies. The reasons

for exercise reporting quality being higher for lower limb tendinopathies compared to other musculoskeletal conditions are unclear but could be related to the fact that resistance training has been considered the gold standard first-line intervention which has been recommended for many years [11].

This review employed two common tools to describe resistance exercise implementation and reporting; the CERT which evaluates general information about the specific exercise intervention; and the Toigo and Boutellier framework which evaluates mechanobiological and exercise dosage descriptors, alongside an evaluation of five key scientific resistance training principles. The completeness of exercise reporting was high overall across the included studies, but the poor reporting of some items and key resistance training principles is concerning and limits the true translation of the findings regarding the resistance exercises to clinical practice. For example, the poor overall reporting on the specific loading employed during resistance exercise for progression, makes this principle difficult to replicate and translate to clinical practice. Although the Toigo and Boutellier framework is a well-accepted tool for reporting exercise descriptors in the literature, some of the items which were poorly reported in the included studies could theoretically be considered too detailed and impractical to properly implement both clinically and in research. The items, rest between sets, time under tension, and volitional muscular failure were all poorly reported; however, it could be argued these items are the least relevant for exercise replication. Despite this, their inclusion would allow for a complete replication of the resistance training interventions implemented in the studies, which may be a more optimal scenario for prescribing resistance exercise to patients to ensure complete translation of the protocols. As a tool that was developed in the sports science literature, it is not rehabilitation specific, so key aspects of resistance exercises in the rehabilitation setting are not accounted for such as the patients psychological state, level of pain, and tolerance to exercise. Similarly, while most items on the CERT were well reported, several items were very poorly reported such as motivational strategies, fidelity measures, adherence, adverse events, and if exercises were delivered as planned. Although the absence of these items does not prevent exercise replication, their inclusion would optimise replication in the clinical setting. It is concerning that only 2% of studies reported using motivational strategies, which may impact on exercise adherence, and overall reporting of adherence was lacking, despite many studies stating they employed adherence tracking measures such as exercise diaries. The very poor reporting of fidelity and adherence highlighted in this review, highlights the need for future studies to focus on practical implementation issues to ensure translation to clinical practice. If fidelity and adherence of the interventions are not monitored and reported, then, the quality of the exercise intervention reported in studies may be of less value. While the combination of both scales would allow exercise replication in a clinical setting, in

isolation, they would likely not be sufficient. Therefore, the development of a more rehabilitation specific scale for implementing and reporting resistance training interventions should be explored in future research to optimise clinical translation of research resistance exercise interventions.

4.1. Clinical Implications. For many years, progressive resistance training has been considered the gold standard intervention for rehabilitating lower limb tendinopathies. Optimising rehabilitation outcomes for patients with lower limb tendinopathies requires implementing the most effective evidence-based resistance training interventions. However, interventions shown to be efficacious in high-quality research must also be replicable and translatable to the clinical setting. To achieve this, research interventions must follow scientific resistance training principles and include enough detail to be reproducible. Many of the studies included in this review which have shown good outcomes also score highly for replicability on the scales employed, particularly those using eccentric heel raises for Achilles tendinopathy and eccentric decline squats for patellar tendinopathy. These protocols should remain the gold standard first-line intervention for clinicians as they have not only been found to be effective in the individual studies but can also be fully replicated clinically. The supplementary material for this review provides all the extracted data and key prescription content from the interventions and can help to guide clinicians in clinical practice. Several studies [50, 79, 177, 221] scored highly across all the reporting tools employed, so the authors recommend these as a starting point for clinicians requiring fully reproducible resistance training programs for implementing in rehabilitation for lower limb tendinopathies.

4.2. Implications for Research. In comparison to reviews on other musculoskeletal pathologies in physical therapy, which have evaluated exercise reporting, this review has found that reporting of resistance exercise for lower limb tendinopathies is generally of high quality, despite shortcomings in areas such as adherence reporting. This is in stark contrast to reviews on other common musculoskeletal disorders where the reporting was considered poor overall. Indeed, the average scores found in this review for the CERT and Toigo and Boutellier framework are higher than for all the other pathologies previously listed. However, this review has highlighted potential weaknesses in the delivery of exercise, with poor reporting of adherence, fidelity, exercise motivational strategies, and specific progression parameters. The implementation of these items should be addressed in future research to optimise clinical translation and outcomes for resistance training interventions in lower limb tendinopathies. Until a rehabilitation specific exercise reporting scale is validated, and available, future studies should continue to design and implement resistance training protocols using scales such as the CERT and Toigo and Boutellier framework, to ensure they are clinically reproducible.

4.3. Strengths and Limitations. This scoping review has included a broad range of study designs from RCTs to individual case reports, with a broad range of interventions, so there is therefore vast heterogeneity in findings across all the studies, so findings should be interpreted with caution. However, determining effectiveness of interventions through meta-analysis techniques was not the objective of the review, with the aims focused on the description, reporting, and implementation of resistance training in interventions for lower limb tendinopathies. Only studies available in English language were included, which may introduce language bias. To provide a comprehensive analysis and state of the art review on reporting of resistance training interventions for lower limb tendinopathies, all primary study designs were considered. Despite all primary designs being included, this review did not consider review papers or clinical practice guidelines, which may have included detailed exercise reporting. Databases were searched from inception, and there was no limitation on sample size or intervention duration. Although many studies included were published before the publication of the CERT (2016) and Toigo and Boutellier framework (2006), there was no obvious reporting discrepancies from earlier to more recent studies, despite the culture of reporting becoming more widespread in recent years [23, 26]. Both scales are transparent and contain sufficient exercise details to allow 100% replication if fully followed, despite not being rehabilitation or tendinopathy specific. Most of the studies included in this review were for Achilles and patellar tendinopathies which also had the highest quality reporting, with other lower limb tendinopathies poorly represented and with comparatively poorer overall reporting quality. Therefore, the findings of this review cannot be generalised to all lower limb tendinopathies, with future research required to address the dearth of resistance training interventions for non-Achilles and patellar lower limb tendinopathies.

5. Conclusion

Resistance training interventions and specific exercises are generally well reported across all primary study designs for treating lower limb tendinopathies, particularly eccentric training for Achilles and patellar tendinopathy. However, certain reporting items and training principles related to the delivery of exercise were poorly reported and implemented, including adherence, fidelity, exercise motivational strategies, and specific progression parameters. While most studies provided sufficient details to allow clinical exercise replication, the weaknesses highlighted must be addressed in future research to allow resistance training interventions and exercises to be fully clinically reproducible to enhance rehabilitation outcomes.

Data Availability

All data relevant to the study are included in the article or are available in the supplementary files or appendices.

Disclosure

A preprint version of this manuscript has previously been published on the SportRxiv preprint server [241]. No patients were involved in this study.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

IB conceptualised the work and developed the methods, search strategy, and framework for the review. IB and AM contributed to the development of the research questions and the study design. All authors developed the first and subsequent drafts of the manuscript and reviewed and approved the manuscript.

Supplementary Materials

S1: MEDLINE search strategy for the review. S2: study characteristics and reporting scores of included studies. S3: application of resistance training principles in included studies. S4: Toigo and Boutellier framework exercise descriptors reporting for each study. S5: Consensus on Exercise Reporting Template (CERT) items reporting for each study. (*Supplementary Materials*)

References

- [1] N. L. Millar, K. G. Silbernagel, and K. Thorborg, "Nature reviews disease primers," *Tendinopathy. Nat Rev Dis Primers*, vol. 7, 2021.
- [2] C. C. Skjong, A. K. Meininger, and S. S. W. Ho, "Tendinopathy treatment: where is the evidence?" *Clinics in Sports Medicine*, vol. 31, no. 2, pp. 329–350, 2012.
- [3] S. P. Magnusson and M. Kjaer, "The impact of loading, unloading, ageing and injury on the human tendon," *The Journal of Physiology*, vol. 597, no. 5, pp. 1283–1298, 2019.
- [4] U. Fredberg and K. Stengaard-Pedersen, "Chronic tendinopathy tissue pathology, pain mechanisms, and etiology with a special focus on inflammation," *Scandinavian Journal of Medicine & Science in Sports*, vol. 18, no. 1, pp. 3–15, 2008.
- [5] A. L. Seitz, P. W. McClure, S. Finucane, N. D. Boardman 3rd, and L. A. Michener, "Mechanisms of rotator cuff tendinopathy: intrinsic, extrinsic, or both?" *Clinical Biomechanics*, vol. 26, no. 1, pp. 1–12, 2011.
- [6] S. Steinmann, C. G. Pfeifer, C. Brochhausen, and D. Docheva, "Spectrum of tendon pathologies: triggers, trails and end-state," *International Journal of Molecular Sciences*, vol. 21, 2020.
- [7] J. Zwerver, S. W. Bredeweg, and I. van den Akker-Scheek, "Prevalence of jumper's knee among nonelite athletes from different sports," *The American Journal of Sports Medicine*, vol. 39, no. 9, pp. 1984–1988, 2011.
- [8] M. J. Arnold and A. L. Moody, "Common running injuries: evaluation and management," *American Family Physician*, vol. 97, pp. 510–516, 2018.
- [9] I. Janssen, H. van der Worp, S. Hensing, and J. Zwerver, "Investigating achilles and patellar tendinopathy prevalence in elite athletics," *Research in Sports Medicine*, vol. 26, no. 1, pp. 1–12, 2018.

- [10] A. L. Sprague, A. H. Smith, P. Knox, R. T. Pohlig, and K. Grävare Silbernagel, "Modifiable risk factors for patellar tendinopathy in athletes: a systematic review and meta-analysis," *British Journal of Sports Medicine*, vol. 52, no. 24, pp. 1575–1585, 2018.
- [11] I. Burton and A. McCormack, "The implementation of resistance training principles in exercise interventions for lower limb tendinopathy: a systematic review," *Physical Therapy in Sport*, vol. 50, pp. 97–113, 2021.
- [12] P. Malliaras, C. J. Barton, N. D. Reeves, and H. Langberg, "Achilles and patellar tendinopathy loading programmes," *Sports Medicine*, vol. 43, no. 4, pp. 267–286, 2013.
- [13] T. B. Cardoso, T. Pizzari, R. Kinsella, D. Hope, and J. L. Cook, "Current trends in tendinopathy management," *Best Practice & Research Clinical Rheumatology*, vol. 33, no. 1, pp. 122–140, 2019.
- [14] R. Beyer, M. Kongsgaard, B. Hougs Kjær, T. Øhlenschläger, M. Kjær, and S. P. Magnusson, "Heavy slow resistance versus eccentric training as treatment for achilles tendinopathy," *The American Journal of Sports Medicine*, vol. 43, no. 7, pp. 1704–1711, 2015.
- [15] C. Clifford, L. Paul, G. Syme, and N. L. Millar, "Isometric versus isotonic exercise for greater trochanteric pain syndrome: a randomised controlled pilot study," *BMJ open sport & exercise medicine*, vol. 5, Article ID e000558, 2019.
- [16] M. Kongsgaard, V. Kovanen, P. Aagaard et al., "Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy," *Scandinavian Journal of Medicine & Science in Sports*, vol. 19, no. 6, pp. 790–802, 2009.
- [17] M. S. Rathleff, C. M. Mølgaard, U. Fredberg et al., "High-load strength training improves outcome in patients with plantar fasciitis: a randomized controlled trial with 12-month follow-up," *Scandinavian Journal of Medicine & Science in Sports*, vol. 25, pp. e292–300, 2015.
- [18] S. Bohm, F. Mersmann, and A. Arampatzis, "Human tendon adaptation in response to mechanical loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults," *Sports Medicine-Open*, vol. 1, no. 1, p. 7, 2015.
- [19] H. Riel, M. B. Jensen, J. L. Olesen, B. Vicenzino, and M. S. Rathleff, "Self-dosed and pre-determined progressive heavy-slow resistance training have similar effects in people with plantar fasciopathy: a randomised trial," *Journal of Physiotherapy*, vol. 65, no. 3, pp. 144–151, 2019.
- [20] K. M. Khan and A. Scott, "Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair," *British Journal of Sports Medicine*, vol. 43, no. 4, pp. 247–252, 2009.
- [21] P. Malliaras, R. Johnston, G. Street et al., "The efficacy of higher versus lower dose exercise in rotator cuff tendinopathy: a systematic review of randomized controlled trials," *Archives of Physical Medicine and Rehabilitation*, vol. 101, no. 10, pp. 1822–1834, 2020.
- [22] B. Østerås, H. Østerås, T. A. Torstensen, and O. Vasseljen, "Dose-response effects of medical exercise therapy in patients with patellofemoral pain syndrome: a randomised controlled clinical trial," *Physiotherapy*, vol. 99, no. 2, pp. 126–131, 2013.
- [23] S. Holden and C. J. Barton, "What should I prescribe?: time to improve reporting of resistance training programmes to ensure accurate translation and implementation," *British Journal of Sports Medicine*, vol. 53, no. 5, pp. 264–265, 2019.
- [24] S. C. Slade, C. E. Dionne, M. Underwood, and R. Buchbinder, "Consensus on exercise reporting template (CERT): explanation and elaboration statement," *British Journal of Sports Medicine*, vol. 50, no. 23, pp. 1428–1437, 2016.
- [25] M. Toigo and U. Boutellier, "New fundamental resistance exercise determinants of molecular and cellular muscle adaptations," *European Journal of Applied Physiology*, vol. 97, no. 6, pp. 643–663, 2006.
- [26] S. Holden, M. S. Rathleff, M. B. Jensen, and C. J. Barton, "How can we implement exercise therapy for patellofemoral pain if we don't know what was prescribed? A systematic review," *British Journal of Sports Medicine*, vol. 52, 2018.
- [27] S. M. Auliffe, V. Korakakis, R. Hilfiker, R. Whiteley, and K. O'Sullivan, "Participant characteristics are poorly reported in exercise trials in tendinopathy: a systematic review," *Physical Therapy in Sport*, vol. 48, pp. 43–53, 2021.
- [28] J. Naunton, G. Street, C. Littlewood, T. Haines, and P. Malliaras, "Effectiveness of progressive and resisted and non-progressive or non-resisted exercise in rotator cuff related shoulder pain: a systematic review and meta-analysis of randomized controlled trials," *Clinical Rehabilitation*, vol. 34, no. 9, pp. 1198–1216, 2020.
- [29] M. H. Ross, M. D. Smith, R. Mellor, and B. Vicenzino, "Exercise for posterior tibial tendon dysfunction: a systematic review of randomised clinical trials and clinical guidelines," *BMJ open sport & exercise medicine*, vol. 4, Article ID e000430, 2018.
- [30] M. Christensen, J. A. Zellers, I. L. Kjær, K. G. Silbernagel, and M. S. Rathleff, "Resistance exercises in early functional rehabilitation for achilles tendon ruptures are poorly described: a scoping review," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 50, no. 12, pp. 681–690, 2020.
- [31] A. J. Goff, W. S. Page, and N. C. Clark, "Reporting of acute programme variables and exercise descriptors in rehabilitation strength training for tibiofemoral joint soft tissue injury: a systematic review," *Physical Therapy in Sport*, vol. 34, pp. 227–237, 2018.
- [32] D. H. Major, Y. Røe, M. Grotle et al., "Content reporting of exercise interventions in rotator cuff disease trials: results from application of the consensus on exercise reporting template (CERT)," *BMJ open sport & exercise medicine*, vol. 5, Article ID e000656, 2019.
- [33] Y. R. Dos Santos Franco, G. C. Miyamoto, K. F. M. Franco, R. R. de Oliveira, and C. M. N. Cabral, "Exercise therapy in the treatment of tendinopathies of the lower limbs: a protocol of a systematic review," *Systematic Reviews*, vol. 8, 2019.
- [34] A. C. Tricco, E. Lillie, W. Zarin et al., "PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation," *Annals of Internal Medicine*, vol. 169, no. 7, pp. 467–473, 2018.
- [35] W. D. Stanish, R. M. Rubinovich, and S. Curwin, "Eccentric exercise in chronic tendinitis," *Clinical Orthopaedics and Related Research*, vol. 208, no. &NA;, pp. 65–68, 1986.
- [36] H. Alfredson, T. Pietilä, P. Jonsson, and R. Lorentzon, "Heavy-load eccentric calf muscle training for the treatment of chronic achilles tendinosis," *The American Journal of Sports Medicine*, vol. 26, no. 3, pp. 360–366, 1998.
- [37] N. A. Abdelkader, M. N. K. Helmy, N. A. Fayaz, and E. S. B. Saweeres, "Short- and intermediate-term results of extracorporeal shockwave therapy for noninsertional achilles tendinopathy," *Foot & Ankle International*, vol. 42, no. 6, pp. 788–797, 2021.

- [38] R. Balius, G. Álvarez, F. Baró et al., “A 3-arm randomized trial for achilles tendinopathy: eccentric training, eccentric training plus a dietary supplement containing mucopolysaccharides, or passive stretching plus a dietary supplement containing mucopolysaccharides,” *Current Therapeutic Research*, vol. 78, pp. 1–7, 2016.
- [39] K. J. Bell, M. L. Fulcher, D. S. Rowlands, and N. Kerse, “Impact of autologous blood injections in treatment of mid-portion achilles tendinopathy: double blind randomised controlled trial,” *BMJ*, vol. 346, p. f2310, 2013.
- [40] A. P. Boesen, R. Hansen, M. I. Boesen, P. Malliaras, and H. Langberg, “Effect of high-volume injection, platelet-rich plasma, and sham treatment in chronic midportion achilles tendinopathy: a randomized double-blinded prospective study,” *The American Journal of Sports Medicine*, vol. 45, no. 9, pp. 2034–2043, 2017.
- [41] R. Brown, J. Orchard, M. Kinchington, A. Hooper, and G. Nalder, “Aprotinin in the management of achilles tendinopathy: a randomised controlled trial,” *British Journal of Sports Medicine*, vol. 40, no. 3, pp. 275–279, 2006.
- [42] R. Chester, M. L. Costa, L. Shepstone, A. Cooper, and S. T. Donell, “Eccentric calf muscle training compared with therapeutic ultrasound for chronic Achilles tendon pain-A pilot study,” *Manual Therapy*, vol. 13, no. 6, pp. 484–491, 2008.
- [43] A. Choudhary, S. Sahu, A. Vasudeva et al., “Comparing effectiveness of combination of collagen peptide type-1, low molecular weight chondroitin sulphate, sodium hyaluronate, and vitamin-C versus oral diclofenac sodium in achilles tendinopathy: a prospective randomized control trial,” *Cureus*, vol. 13, p. 19737, 2021.
- [44] S. de Jonge, R. J. de Vos, H. T. M. Van Schie, J. A. N. Verhaar, A. Weir, and J. L. Tol, “One-year follow-up of a randomised controlled trial on added splinting to eccentric exercises in chronic midportion achilles tendinopathy,” *British Journal of Sports Medicine*, vol. 44, no. 9, pp. 673–677, 2010.
- [45] S. de Jonge, R. J. de Vos, A. Weir et al., “One-year follow-up of platelet-rich plasma treatment in chronic achilles tendinopathy,” *The American Journal of Sports Medicine*, vol. 39, no. 8, pp. 1623–1630, 2011.
- [46] R. J. De Vos, A. Weir, H. T. M. van Schie et al., “Platelet-Rich plasma injection for chronic achilles tendinopathy,” *JAMA*, vol. 303, no. 2, pp. 144–149, 2010.
- [47] R. J. de Vos, A. Weir, R. J. Visser, T. de Winter, and J. L. Tol, “The additional value of a night splint to eccentric exercises in chronic midportion achilles tendinopathy: a randomised controlled trial,” *British Journal of Sports Medicine*, vol. 41, 2007.
- [48] M. Gatz, M. Betsch, T. Dirrichs et al., “Eccentric and isometric exercises in achilles tendinopathy evaluated by the VISA-A score and shear wave elastography,” *Sport Health: A Multidisciplinary Approach*, vol. 12, no. 4, pp. 373–381, 2020.
- [49] B. Habets, R. E. H. van Cingel, F. J. G. Backx, H. J. van Elten, P. Zuithoff, and B. M. A. Huisstede, “No difference in clinical effects when comparing alfredson eccentric and silbernagel combined concentric-eccentric loading in achilles tendinopathy: a randomized controlled trial,” *Orthopaedic journal of sports medicine*, vol. 9, no. 10, 2021.
- [50] F. Hasani, T. Haines, S. E. Munteanu, P. Schoch, B. Vicenzino, and P. Malliaras, “LOAD-intensity and time-under-tension of exercises for men who have achilles tendinopathy (the LOADIT trial): a randomised feasibility trial,” *BMC Sports Science, Medicine and Rehabilitation*, vol. 13, no. 1, p. 57, 2021.
- [51] L. Herrington and R. McCulloch, “The Role of Eccentric Training in the Management of Achilles tendinopathy: a pilot study,” *Physical Therapy in Sport*, vol. 8, 2007.
- [52] T. Horstmann, H. M. Jud, V. Fröhlich, A. Mündermann, and S. Grau, “Whole-body vibration versus eccentric training or a wait-and-see approach for chronic achilles tendinopathy: a randomized clinical trial,” *Journal of Orthopaedic & Sports Physical Therapy*, vol. 43, no. 11, pp. 794–803, 2013.
- [53] R. S. Kearney, N. Parsons, and M. L. Costa, “Achilles tendinopathy management,” *Bone & Joint Research*, vol. 2, no. 10, pp. 227–232, 2013.
- [54] M. Kedia, M. Williams, L. Jain et al., “The effects of conventional physical therapy and eccentric strengthening for insertional achilles tendinopathy,” *International Journal of Sports Physical Therapy*, vol. 9, pp. 488–497, 2014.
- [55] K. Knobloch, L. Schreibermueller, U. G. Longo, and P. M. Vogt, “Eccentric exercises for the management of tendinopathy of the main body of the achilles tendon with or without the AirHeel brace. A randomized controlled trial. A: effects on pain and microcirculation,” *Disability & Rehabilitation*, vol. 30, no. 20-22, pp. 1685–1691, 2008.
- [56] A. Koszaliniski, T. Flynn, M. Hellman, and J. A. Cleland, “Trigger point dry needling, manual therapy and exercise versus manual therapy and exercise for the management of achilles tendinopathy: a feasibility study,” *Journal of Manual & Manipulative Therapy*, vol. 28, no. 4, pp. 212–221, 2020.
- [57] N. Mafi, R. Lorentzon, and H. Alfredson, “Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic achilles tendinosis,” *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 9, no. 1, pp. 42–47, 2001.
- [58] N. S. B. Mansur, F. T. Matsunaga, O. L. Carrazzone et al., “Shockwave therapy plus eccentric exercises versus isolated eccentric exercises for achilles insertional tendinopathy,” *Journal of Bone and Joint Surgery*, vol. 103, no. 14, pp. 1295–1302, 2021.
- [59] J. R. McCormack, F. B. Underwood, E. J. Slaven, and T. A. Cappaert, “Eccentric exercise versus eccentric exercise and soft tissue treatment (astym) in the management of insertional achilles tendinopathy,” *Sport Health: A Multidisciplinary Approach*, vol. 8, no. 3, pp. 230–237, 2016.
- [60] S. E. Munteanu, L. A. Scott, D. R. Bonanno et al., “Effectiveness of customised foot orthoses for achilles tendinopathy: a randomised controlled trial,” *British Journal of Sports Medicine*, vol. 49, no. 15, pp. 989–994, 2015.
- [61] S. L. Nisen-Vertommen, J. E. Taunton, D. B. Clement, and R. E. Mosher, “The effect OF eccentric versus concentric exercise IN the management OF achilles tendonitis,” *Clinical Journal of Sport Medicine*, vol. 2, pp. 109–113, 1992.
- [62] J. Nørregaard, C. C. Larsen, T. Bieler, and H. Langberg, “Eccentric exercise in treatment of achilles tendinopathy,” *Scandinavian Journal of Medicine & Science in Sports*, vol. 17, no. 2, pp. 133–138, 2007.
- [63] A. Notarnicola, G. Maccagnano, S. Tafuri, M. I. Forcignanò, A. Panella, and B. Moretti, “CHELT therapy in the treatment of chronic insertional achilles tendinopathy,” *Lasers in Medical Science*, vol. 29, no. 3, pp. 1217–1225, 2014.
- [64] J. Pearson, D. Rowlands, and R. Highet, “Autologous blood injection to treat achilles tendinopathy? A randomized controlled trial,” *Journal of Sport Rehabilitation*, vol. 21, no. 3, pp. 218–224, 2012.

- [65] W. Petersen, R. Welp, and D. Rosenbaum, "Chronic achilles tendinopathy," *The American Journal of Sports Medicine*, vol. 35, no. 10, pp. 1659–1667, 2007.
- [66] S. F. E. Praet, C. R. Purdam, M. Welvaert et al., "Oral supplementation of specific collagen peptides combined with calf-strengthening exercises enhances function and reduces pain in achilles tendinopathy patients," *Nutrients*, vol. 11, 2019.
- [67] C. L. Rabusin, H. B. Menz, J. A. McClelland et al., "Efficacy of heel lifts versus calf muscle eccentric exercise for mid-portion achilles tendinopathy (HEALTHY): a randomised trial," *British Journal of Sports Medicine*, vol. 55, no. 9, pp. 486–492, 2021.
- [68] C. Romero-Morales, P. J. Martín-Llantino, C. Calvo-Lobo et al., "Effectiveness of eccentric exercise and a vibration or cryotherapy program in enhancing rectus abdominis muscle thickness and inter-rectus distance in patients with chronic mid-portion achilles tendinopathy: a randomized clinical trial," *International Journal of Medical Sciences*, vol. 15, no. 14, pp. 1764–1770, 2018.
- [69] C. Romero-Morales, P. J. Martín-Llantino, C. Calvo-Lobo et al., "Vibration increases multifidus cross-sectional area versus cryotherapy added to chronic non-insertional achilles tendinopathy eccentric exercise," *Physical Therapy in Sport*, vol. 42, pp. 61–67, 2020.
- [70] J. D. Rompe, J. Furia, and N. Maffulli, "Eccentric loading compared with shock wave treatment for chronic insertional achilles tendinopathy," *Journal of Bone and Joint Surgery American Volume*, vol. 90, no. 1, pp. 52–61, 2008.
- [71] J. D. Rompe, J. Furia, and N. Maffulli, "Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy," *The American Journal of Sports Medicine*, vol. 37, no. 3, pp. 463–470, 2009.
- [72] J. D. Rompe, B. Nafe, J. P. Furia, and N. Maffulli, "Eccentric loading, shock-wave treatment, or a wait- and-see policy for tendinopathy of the main body of tendo achillis," *The American Journal of Sports Medicine*, vol. 35, no. 3, pp. 374–383, 2007.
- [73] E. M. Roos, M. Engstrom, A. Lagerquist, and B. Soderberg, "Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy - a randomized trial with 1-year follow-up," *Scandinavian Journal of Medicine & Science in Sports*, vol. 14, no. 5, pp. 286–295, 2004.
- [74] K. G. Silbernagel, R. Thomeé, B. I. Eriksson, and J. Karlsson, "Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with achilles tendinopathy," *The American Journal of Sports Medicine*, vol. 35, no. 6, pp. 897–906, 2007.
- [75] K. Grävare Silbernagel, R. Thomeé, P. Thomeé, and J. Karlsson, "Eccentric overload training for patients with chronic Achilles tendon pain - a randomised controlled study with reliability testing of the evaluation methods," *Scandinavian Journal of Medicine & Science in Sports*, vol. 11, no. 4, pp. 197–206, 2001.
- [76] L. Solomons, J. J. Y. Lee, M. Bruce, L. D. White, and A. Scott, "Intramuscular stimulation vs sham needling for the treatment of chronic midportion achilles tendinopathy: a randomized controlled clinical trial," *PLoS One*, vol. 15, Article ID e0238579, 2020.
- [77] S. H. Stefansson, S. Brandsson, H. Langberg, and A. Arnason, "Using pressure massage for achilles tendinopathy: a single-blind, randomized controlled trial comparing a novel treatment versus an eccentric exercise protocol," *Orthopaedic journal of sports medicine*, vol. 7, 2019.
- [78] A. Stergioulas, M. Stergioula, R. Aarskog, R. A. B. Lopes-Martins, and J. M. Bjordal, "Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy," *The American Journal of Sports Medicine*, vol. 36, no. 5, pp. 881–887, 2008.
- [79] M. Stevens and C.-W. Tan, "Effectiveness of the alfredson protocol compared with a lower repetition-volume protocol for midportion achilles tendinopathy: a randomized controlled trial," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 44, no. 2, pp. 59–67, 2014.
- [80] S. Tumilty, R. Mani, and G. D. Baxter, "Photobiomodulation and eccentric exercise for achilles tendinopathy: a randomized controlled trial," *Lasers in Medical Science*, vol. 31, no. 1, pp. 127–135, 2016.
- [81] S. Tumilty, S. McDonough, D. A. Hurley, and G. D. Baxter, "Clinical effectiveness of low-level laser therapy as an adjunct to eccentric exercise for the treatment of achilles' tendinopathy: a randomized controlled trial," *Archives of Physical Medicine and Rehabilitation*, vol. 93, no. 5, pp. 733–739, 2012.
- [82] A. C. van der Vlist, R. F. van Oosterom, P. L. J. van Veldhoven et al., "Effectiveness of a high volume injection as treatment for chronic achilles tendinopathy: randomised controlled trial," *BMJ*, vol. 370, 2020.
- [83] M. J. Yelland, K. R. Sweeting, J. A. Lyftogt, S. K. Ng, P. A. Scuffham, and K. A. Evans, "Prolotherapy injections and eccentric loading exercises for painful achilles tendinosis: a randomised trial," *British Journal of Sports Medicine*, vol. 45, no. 5, pp. 421–428, 2011.
- [84] J. Yu, D. Park, and G. Lee, "Effect of eccentric strengthening on pain, muscle strength, endurance, and functional fitness factors in male patients with achilles tendinopathy," *American Journal of Physical Medicine & Rehabilitation*, vol. 92, no. 1, pp. 68–76, 2013.
- [85] B.-M. Zhang, L.-W. Zhong, S.-W. Xu, H.-R. Jiang, and J. Shen, "Acupuncture for chronic achilles tendinopathy: a randomized controlled study," *Chinese Journal of Integrative Medicine*, vol. 19, no. 12, pp. 900–904, 2013.
- [86] V. Kannappan and A. M. Sathosh, "To compare the effect of eccentric exercises and isometric exercises for achilles tendinitis in skaters," *Journal of Lifestyle Medicine*, vol. 10, no. 1, pp. 49–54, 2020.
- [87] D. Stasinopoulos and P. Manias, "Comparing two eccentric exercise programmes for the management of achilles tendinopathy. A pilot trial," *Journal of Bodywork and Movement Therapies*, vol. 17, no. 3, pp. 309–315, 2013.
- [88] A. C. van der Vlist, P. L. J. van Veldhoven, R. F. van Oosterom, J. A. N. Verhaar, and R. J. de Vos, "Isometric exercises do not provide immediate pain relief in Achilles tendinopathy: a quasi-randomized clinical trial," *Scandinavian Journal of Medicine & Science in Sports*, vol. 30, no. 9, pp. 1712–1721, 2020.
- [89] R. J. de Vos, M. P. Heijboer, H. Weinans, J. A. N. Verhaar, and H. T. M. van Schie, "Tendon structure's lack of relation to clinical outcome after eccentric exercises in chronic midportion achilles tendinopathy," *Journal of Sport Rehabilitation*, vol. 21, no. 1, pp. 34–43, 2012.
- [90] T. Masood, K. Kalliokoski, S. P. Magnusson, J. Bojsen-Moller, and T. Finni, "Effects of 12-wk eccentric calf muscle training on muscle-tendon glucose uptake and SEMG in patients with chronic achilles tendon pain," *Journal of Applied Physiology*, vol. 117, pp. 105–111, 1985.

- [91] M. Abate, L. Di Carlo, A. Belluati, and V. Salini, "Factors associated with positive outcomes of platelet-rich plasma therapy in achilles tendinopathy," *European Journal of Orthopaedic Surgery and Traumatology*, vol. 30, no. 5, pp. 859–867, 2020.
- [92] H. Alfredson and R. Lorentzon, "Intratendinous glutamate levels and eccentric training in chronic achilles tendinosis: a prospective study using microdialysis technique," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 11, no. 3, pp. 196–199, 2003.
- [93] H. Alfredson, P. Nordström, T. Pietilä, and R. Lorentzon, "Bone mass in the calcaneus after heavy loaded eccentric calf-muscle training in recreational athletes with chronic achilles tendinosis," *Calcified Tissue International*, vol. 64, no. 5, pp. 450–455, 1999.
- [94] P. Angermann and D. Hovgaard, "Chronic achilles tendinopathy in athletic individuals: results of nonsurgical treatment," *Foot & Ankle International*, vol. 20, no. 5, pp. 304–306, 1999.
- [95] S. de Jonge, J. L. Tol, A. Weir, J. H. Waarsing, J. A. N. Verhaar, and R.-J. de Vos, "The tendon structure returns to asymptomatic values in nonoperatively treated achilles tendinopathy but is not associated with symptoms," *The American Journal of Sports Medicine*, vol. 43, no. 12, pp. 2950–2958, 2015.
- [96] M. Fahlström, P. Jonsson, R. Lorentzon, and H. Alfredson, "Chronic achilles tendon pain treated with eccentric calf-muscle training," *Knee Surgery, Sports Traumatology, Arthroscopy: Official Journal of the ESSKA*, vol. 11, pp. 327–333, 2003.
- [97] P. Jonsson, H. Alfredson, K. Sunding, M. Fahlstrom, and J. Cook, "New regimen for eccentric calf-muscle training in patients with chronic insertional achilles tendinopathy: results of a pilot study," *British Journal of Sports Medicine*, vol. 42, no. 9, pp. 746–749, 2008.
- [98] I. F. Lagas, A. C. van der Vlist, R. F. van Oosterom et al., "Victorian institute of sport assessment-achilles (VISA-A) questionnaire-minimal clinically important difference for active people with midportion achilles tendinopathy: a prospective cohort study," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 51, no. 10, pp. 510–516, 2021.
- [99] N. Maffulli, G. Walley, M. K. Sayana, U. G. Longo, and V. Denaro, "Eccentric calf muscle training in athletic patients with achilles tendinopathy," *Disability & Rehabilitation*, vol. 30, no. 20-22, pp. 1677–1684, 2008.
- [100] N. S. B. Mansur, T. Baumfeld, F. Villalon et al., "Shockwave therapy associated with eccentric strengthening for achilles insertional tendinopathy: a prospective study," *Foot & Ankle Specialist*, vol. 12, no. 6, pp. 540–545, 2019.
- [101] L. Mantovani, L. Maestroni, F. Bettariga, M. Gobbo, N. F. Lopomo, and S. McLean, "Does isometric exercise improve leg stiffness and hop pain in subjects with Achilles tendinopathy? A feasibility study," *Physical Therapy in Sport*, vol. 46, pp. 234–242, 2020.
- [102] S. O'Neill, J. Radia, K. Bird et al., "Acute sensory and motor response to 45-s heavy isometric holds for the plantar flexors in patients with Achilles tendinopathy," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 27, no. 9, pp. 2765–2773, 2019.
- [103] L. Ohberg and H. Alfredson, "Effects on neovascularisation behind the good results with eccentric training in chronic mid-portion achilles tendinosis?" *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 12, pp. 465–470, 2004.
- [104] C. C. Ooi, M. Schneider, P. Malliaras et al., "Real-time sonoelastography evaluation of the achilles tendon following ultrasound-guided platelet-rich plasma injection and eccentric exercise for the treatment of refractory achilles tendinopathy," *Ultrasound*, vol. 27, no. 3, pp. 138–147, 2019.
- [105] D. M. Robinson, C. O. Tan, and A. S. Tenforde, "Functional gains using radial and combined shockwave therapy in the management of achilles tendinopathy," *Journal of Foot & Ankle Surgery*, vol. 76, 2021.
- [106] I. Sancho, D. Morrissey, R. W. Willy, C. Barton, and P. Malliaras, "Education and exercise supplemented by a pain-guided hopping intervention for male recreational runners with midportion achilles tendinopathy: a single cohort feasibility study," *Physical Therapy in Sport*, vol. 40, pp. 107–116, 2019.
- [107] M. K. Sayana and N. Maffulli, "Eccentric calf muscle training in non-athletic patients with achilles tendinopathy," *Journal of Science and Medicine in Sport*, vol. 10, no. 1, pp. 52–58, 2007.
- [108] A. Shalabi, M. Kristoffersen-Wilberg, L. Svensson, P. Aspelin, and T. Movin, "Eccentric training of the gastrocnemius-soleus complex in chronic achilles tendinopathy results in decreased tendon volume and intratendinous signal as evaluated by MRI," *The American Journal of Sports Medicine*, vol. 32, no. 5, pp. 1286–1296, 2004.
- [109] L. von Wehren, K. Pokorny, F. Blanke, J. Sailer, and M. Majewski, "Injection with autologous conditioned serum has better clinical results than eccentric training for chronic achilles tendinopathy," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 27, no. 9, pp. 2744–2753, 2019.
- [110] M. Wei, Y. Liu, Z. Li, and Z. Wang, "Comparison of clinical efficacy among endoscopy-assisted radio-frequency ablation, extracorporeal shockwaves, and eccentric exercises in treatment of insertional achilles tendinosis," *Journal of the American Podiatric Medical Association*, vol. 107, no. 1, pp. 11–16, 2017.
- [111] E. Wetke, F. Johannsen, and H. Langberg, "Achilles tendinopathy: a prospective study on the effect of active rehabilitation and steroid injections in a clinical setting," *Scandinavian Journal of Medicine & Science in Sports*, vol. 25, pp. e392–9, 2015.
- [112] P. C. Wheeler and C. Tattersall, "Novel interventions for recalcitrant achilles tendinopathy: benefits seen following high-volume image-guided injection or extracorporeal shockwave therapy-A prospective cohort study," *Clinical Journal of Sport Medicine: Official Journal of the Canadian Academy of Sport Medicine*, vol. 30, pp. 14–19, 2020.
- [113] H. Langberg, H. Ellingsgaard, T. Madsen et al., "Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with achilles tendinosis," *Scandinavian Journal of Medicine & Science in Sports*, vol. 17, pp. 61–66, 2007.
- [114] Y. H. Park, W. Kim, J. Y. Kim, G. W. Choi, and H. J. Kim, "Clinical impact of metabolic syndrome on eccentric exercises for chronic insertional achilles tendinopathy," *Journal of Foot & Ankle Surgery*, vol. 34, 2021.
- [115] R. Ram, W. Meeuwisse, C. Patel, D. A. Wiseman, and J. P. Wiley, "The limited effectiveness of a home-based eccentric training for treatment of achilles tendinopathy," *Clinical and Investigative Medicine*, vol. 36, no. 4, pp. 197–206, 2013.
- [116] E. Benito, "Physiotherapy's protocol to approach the insertional achilles tendinopathy," *Journal of Human Sport and Exercise*, vol. 11, pp. 358–366, 2016.

- [117] V. M. Deans, A. Miller, and J. Ramos, "A prospective series of patients with chronic achilles tendinopathy treated with autologous-conditioned plasma injections combined with exercise and therapeutic ultrasonography," *Journal of Foot & Ankle Surgery*, vol. 51, no. 6, pp. 706–710, 2012.
- [118] D. J. Jayaseelan, M. Kecman, D. Alcorn, and J. D. Sault, "Manual therapy and eccentric exercise in the management of achilles tendinopathy," *Journal of Manual & Manipulative Therapy*, vol. 25, no. 2, pp. 106–114, 2017.
- [119] V. Pavone, L. Cannavo, A. Di Stefano, G. Testa, L. Costarella, and G. Sessa, "Low-energy extracorporeal shock-wave therapy in the treatment of chronic insertional achilles tendinopathy: a case series," *BioMed Research International*, vol. 65, 2016.
- [120] K. G. Silbernagel, A. Brorsson, and M. Lundberg, "The majority of patients with achilles tendinopathy recover fully when treated with exercise alone," *The American Journal of Sports Medicine*, vol. 39, no. 3, pp. 607–613, 2011.
- [121] P. Syvertson, E. Dietz, M. Matocha et al., "A treatment-based classification algorithm to treat achilles tendinopathy: an exploratory case series," *Journal of Sport Rehabilitation*, vol. 26, no. 3, pp. 260–268, 2017.
- [122] P. C. Wheeler, "Extracorporeal shock wave therapy plus rehabilitation for insertional and noninsertional achilles tendinopathy shows good results across a range of domains of function," *Journal of Foot & Ankle Surgery*, vol. 58, no. 4, pp. 617–622, 2019.
- [123] J. Borda and M. Selhorst, "The use of compression tack and flossing along with lacrosse ball massage to treat chronic achilles tendinopathy in an adolescent athlete: a case report," *Journal of Manual & Manipulative Therapy*, vol. 25, no. 1, pp. 57–61, 2017.
- [124] T. Cuddeford, J. Houck, D. Palmer, J. Beilstein, and J. Visser, "What maximum ankle torque is appropriate for training patients with non-insertional achilles tendinopathy?" *Faculty Publications-College of Physical Therapy*, vol. 100, 2013, https://digitalcommons.georgefox.edu/pt_fac/100.
- [125] B. J. Eckenrode and S. K. Stackhouse, "Improved pressure pain thresholds and function following noxious electrical stimulation on a runner with chronic achilles tendinopathy: a case report," *International journal of sports physical therapy*, vol. 10, no. 3, pp. 354–362, 2015.
- [126] P. Francis, I. Thornley, A. Jones, and M. I. Johnson, "Pain and function in the runner a ten (din) uous link," *Medicina (Kaunas, Lithuania)*, vol. 56, 2020.
- [127] B. L. Greene, "Physical therapist management of fluoroquinolone-induced achilles tendinopathy," *Physical Therapy*, vol. 82, no. 12, pp. 1224–1231, 2002.
- [128] J. R. McCormack, "The management of mid-portion achilles tendinopathy with astym(R) and eccentric exercise: a case report," *Int J Sports Phys Ther*, vol. 7, no. 6, pp. 672–677, 2012.
- [129] J. A. Papa, "Conservative management of achilles tendinopathy: a case report," *Journal of the Canadian Chiropractic Association*, vol. 56, no. 3, pp. 216–224, 2012.
- [130] G. Ross, C. Macfarlane, and B. Vaughan, "Combined osteopathy and exercise management of achilles tendinopathy in an athlete," *The Journal of Sports Medicine and Physical Fitness*, vol. 58, pp. 106–112, 2018.
- [131] R. C. Thompson, "Managing plantar fasciopathy and tendinopathy through conditioning and load monitoring within elite sport," *J. Aust. Strength Cond.* vol. 25, pp. 85–103, 2017.
- [132] A. Gärdin, T. Movin, L. Svensson, and A. Shalabi, "The long-term clinical and MRI results following eccentric calf muscle training in chronic achilles tendinosis," *Skeletal Radiology*, vol. 39, no. 5, pp. 435–442, 2010.
- [133] J.-L. Croisier, B. Forthomme, M. Foidart-Dessalle, B. Godon, and J.-M. Crielaard, "Treatment of recurrent tendinitis by isokinetic eccentric exercises," *Isokinetics and Exercise Science*, vol. 9, no. 2-3, pp. 133–141, 2001.
- [134] K. Pinkelman and J. F. Schilling, "Treatment of extensor hallucis longus tendinosis in a track athlete," *International Journal of Athletic Therapy & Training*, vol. 17, no. 4, pp. 4–7, 2012.
- [135] R. M. Cowan, C. L. Ganderton, J. Cook, A. I. Semciw, D. M. Long, and T. Pizzari, "Does menopausal hormone therapy, exercise, or both improve pain and function in postmenopausal women with greater trochanteric pain syndrome? A 2 x 2 factorial randomized clinical trial," *The American Journal of Sports Medicine*, vol. 23, 2021.
- [136] C. Ganderton, A. Semciw, J. Cook, E. Moreira, and T. Pizzari, "Gluteal loading versus sham exercises to improve pain and dysfunction in postmenopausal women with greater trochanteric pain syndrome: a randomized controlled trial," *Journal of Women's Health*, vol. 27, no. 6, pp. 815–829, 2018.
- [137] R. Mellor, K. Bennell, A. Grimaldi et al., "Education plus exercise versus corticosteroid injection use versus a wait and see approach on global outcome and pain from gluteal tendinopathy: prospective, single blinded, randomised clinical trial," *British Journal of Sports Medicine*, vol. 52, no. 22, pp. 1464–1472, 2018.
- [138] S. Ramon, S. Russo, F. Santoboni et al., "Focused shockwave treatment for greater trochanteric pain syndrome," *Journal of Bone and Joint Surgery*, vol. 102, no. 15, pp. 1305–1311, 2020.
- [139] G. Thompson and J. F. Pearson, "No attributable effects of PRP on greater trochanteric pain syndrome," *New Zealand Medical Journal*, vol. 132, pp. 22–32, 2019.
- [140] P. C. Wheeler, C. Dudson, R. Calver et al., "Three sessions of radial extracorporeal shockwave therapy gives No additional benefit over "minimal-dose" radial extracorporeal shockwave therapy for patients with chronic greater trochanteric pain syndrome: a double-blinded, randomized, controlled trial," *Clinical Journal of Sport Medicine*, vol. 1, no. 32, pp. 7–18, 2022.
- [141] J. C. Van Rooy, "A different approach to the management of greater trochanter pain syndrome," *South African Journal of Physiotherapy*, vol. 65, 2009.
- [142] A. Cacchio, J. D. Rompe, J. P. Furia, P. Susi, V. Santilli, and F. De Paulis, "Shockwave therapy for the treatment of chronic proximal hamstring tendinopathy in professional athletes," *The American Journal of Sports Medicine*, vol. 39, no. 1, pp. 146–153, 2011.
- [143] D. Cushman and M. E. Rho, "Conservative treatment of subacute proximal hamstring tendinopathy using eccentric exercises performed with a treadmill: a case report," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 45, no. 7, pp. 557–562, 2015.
- [144] D. J. Jayaseelan, N. Moats, and C. R. Ricardo, "Rehabilitation of proximal hamstring tendinopathy utilizing eccentric training, lumbopelvic stabilization, and trigger point dry needling: 2 case reports," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 44, no. 3, pp. 198–205, 2014.
- [145] K. Krueger, N. B. Washmuth, and T. D. Williams, "The management of proximal hamstring tendinopathy in a competitive powerlifter with heavy slow resistance training - a case report," *International Journal of Sports Physical Therapy*, vol. 15, no. 5, pp. 814–822, 2020.

- [146] J. R. McCormack, "The management of bilateral high hamstring tendinopathy with ASTYM treatment and eccentric exercise: a case report," *Journal of Manual & Manipulative Therapy*, vol. 20, no. 3, pp. 142–146, 2012.
- [147] C. Rausedo, "The rehabilitation of a runner with iliopsoas tendinopathy using an eccentric-biased exercise—a case report," *International Journal of Sports Physical Therapy*, vol. 12, no. 7, pp. 1150–1162, 2017.
- [148] R. G. Alvarez, A. Marini, C. Schmitt, and C. L. Saltzman, "Stage I and II posterior tibial tendon dysfunction treated by a structured nonoperative management protocol: an orthosis and exercise program," *Foot & Ankle International*, vol. 27, no. 1, pp. 2–8, 2006.
- [149] J. Houck, C. Neville, J. Tome, and A. Flemlister, "Randomized controlled trial comparing orthosis augmented by either stretching or stretching and strengthening for stage II tibialis posterior tendon dysfunction," *Foot & Ankle International*, vol. 36, no. 9, pp. 1006–1016, 2015.
- [150] K. Kulig, S. F. Reischl, A. B. Pomrantz et al., "Nonsurgical management of posterior tibial tendon dysfunction with orthoses and resistive exercise: a randomized controlled trial," *Physical Therapy*, vol. 89, no. 1, pp. 26–37, 2009.
- [151] K. Kulig, E. S. Lederhaus, S. Reischl, S. Arya, and G. Bashford, "Effect of eccentric exercise program for early tibialis posterior tendinopathy," *Foot & Ankle International*, vol. 30, no. 9, pp. 877–885, 2009.
- [152] D. Robinson, M. Mitchkash, L. Wasserman, and A. S. Tenforde, "Nonsurgical approach in management of tibialis posterior tendinopathy with combined radial shockwave and foot core exercises: a case series," *Journal of Foot & Ankle Surgery*, vol. 59, no. 5, pp. 1058–1061, 2020.
- [153] C. Patla, J. Lwin, L. Smith, and E. Chaconas, "Cuboid manipulation and exercise in the management of posterior tibialis tendinopathy: a case report," *International journal of sports physical therapy*, vol. 10, pp. 363–370, 2015.
- [154] F. Abat, J. L. Sánchez-Sánchez, A. M. Martín-Nogueras et al., "Randomized controlled trial comparing the effectiveness of the ultrasound-guided galvanic electrolysis technique (USGET) versus conventional electro-physiotherapeutic treatment on patellar tendinopathy," *Journal of Experimental Orthopaedics*, vol. 3, no. 1, p. 34, 2016.
- [155] A.-S. Agergaard, R. B. Svensson, N. M. Malmgaard-Clausen et al., "Clinical outcomes, structure, and function improve with both heavy and moderate loads in the treatment of patellar tendinopathy: a randomized clinical trial," *The American Journal of Sports Medicine*, vol. 49, no. 4, pp. 982–993, 2021.
- [156] R. Bahr, B. Fossan, S. Løken, and L. Engebretsen, "Surgical treatment compared with eccentric training for patellar tendinopathy (Jumper's knee)," *The Journal of Bone & Joint Surgery*, vol. 88, no. 8, pp. 1689–1698, 2006.
- [157] R. Biernat, Z. Trzaskoma, L. Trzaskoma, and D. Czuprowski, "Rehabilitation protocol for patellar tendinopathy applied among 16- to 19-year old volleyball players," *The Journal of Strength & Conditioning Research*, vol. 28, no. 1, pp. 43–52, 2014.
- [158] S. J. Breda, E. H. G. Oei, J. Zwerver et al., "Effectiveness of progressive tendon-loading exercise therapy in patients with patellar tendinopathy: a randomised clinical trial," *British Journal of Sports Medicine*, vol. 55, no. 9, pp. 501–509, 2021.
- [159] L. J. Cannell, J. E. Taunton, D. B. Clement, C. Smith, and K. M. Khan, "A randomised clinical trial of the efficacy of drop squats or leg extension/leg curl exercises to treat clinically diagnosed jumper's knee in athletes: pilot study," *British Journal of Sports Medicine*, vol. 35, no. 1, pp. 60–64, 2001.
- [160] R. A. D. Cunha, "Comparative study of two protocols of eccentric exercise on knee pain and function in athletes with patellar tendinopathy: randomized controlled study," *Revista Brasileira de Medicina do Esporte*, vol. 18, 2012.
- [161] S. Dimitrios, M. Pantelis, and S. Kalliopi, "Comparing the effects of eccentric training with eccentric training and static stretching exercises in the treatment of patellar tendinopathy. A controlled clinical trial," *Clinical Rehabilitation*, vol. 26, no. 5, pp. 423–430, 2012.
- [162] J. L. Drago, A. S. Wasterlain, H. J. Braun, and K. T. Nead, "Platelet-Rich plasma as a treatment for patellar tendinopathy," *The American Journal of Sports Medicine*, vol. 42, no. 3, pp. 610–618, 2014.
- [163] A. Frohm, T. Saartok, K. Halvorsen, and P. Renström, "Eccentric treatment for patellar tendinopathy: a prospective randomised short-term pilot study of two rehabilitation protocols," *British Journal of Sports Medicine*, vol. 41, p. e7, 2007.
- [164] S. Holden, K. Lyng, T. Graven-Nielsen et al., "Isometric exercise and pain in patellar tendinopathy: a randomized crossover trial," *Journal of Science and Medicine in Sport*, vol. 23, no. 3, pp. 208–214, 2020.
- [165] K. Jensen and R. P. Di Fabio, "Evaluation of eccentric exercise in treatment of patellar tendinitis," *Physical Therapy*, vol. 69, no. 3, pp. 211–216, 1989.
- [166] P. Jonsson and H. Alfredson, "Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: a prospective randomised study," *British Journal of Sports Medicine*, vol. 39, no. 11, pp. 847–850, 2005.
- [167] J. F. Kaux, S. Bornheim, N. Dardenne et al., "Comparison between platelet-rich plasma injections and hyaluronic acid injections in the treatment of patellar tendinopathies: a randomized trial," *Muscles, Ligaments and Tendons Journal*, vol. 9, pp. 322–327, 2019.
- [168] W. C. Lee, G. Y. Ng, Z. J. Zhang, P. Malliaras, L. Masci, and S. N. Fu, "Changes on tendon stiffness and clinical outcomes in athletes are associated with patellar tendinopathy after eccentric exercise," *Clinical Journal of Sport Medicine: Official Journal of the Canadian Academy of Sport Medicine*, vol. 30, pp. 25–32, 2020.
- [169] M. P. López-Royo, J. Ríos-Díaz, R. M. Galán-Díaz, P. Herrero, and E. M. Gómez-Trullén, "A comparative study of treatment interventions for patellar tendinopathy: a randomized controlled trial," *Archives of Physical Medicine and Rehabilitation*, vol. 102, no. 5, pp. 967–975, 2021.
- [170] K. MacDonald, "Effect of eccentric exercises at the knee with hip muscle strengthening to treat patellar tendinopathy in active duty military personnel: A randomized pilot," *Orthopaedic Practice*, vol. 31, 2019.
- [171] J. L. Olesen, M. Hansen, I. F. Turtumoygard et al., "No treatment benefits of local administration of insulin-like growth factor-1 in addition to heavy slow resistance training in tendinopathic human patellar tendons: a randomized, double-blind, placebo-controlled trial with 1-year follow-up," *The American Journal of Sports Medicine*, vol. 49, no. 9, pp. 2361–2370, 2021.
- [172] L. S. Pietrosimone, J. T. Blackburn, E. A. Wikstrom et al., "Landing biomechanics are not immediately altered by a single-dose patellar tendon isometric exercise protocol in male athletes with patellar tendinopathy: a single-blinded randomized cross-over trial," *Physical Therapy in Sport*, vol. 46, pp. 177–185, 2020.

- [173] E. Rio, D. Kidgell, C. Purdam et al., "Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy," *British Journal of Sports Medicine*, vol. 49, no. 19, pp. 1277–1283, 2015.
- [174] E. Rio, M. van Ark, S. Docking et al., "Isometric contractions are more analgesic than isotonic contractions for patellar tendon pain," *Clinical Journal of Sport Medicine*, vol. 27, no. 3, pp. 253–259, 2017.
- [175] D. Ruffino, P. Malliaras, S. Marchegiani, and V. Campana, "Inertial flywheel vs heavy slow resistance training among athletes with patellar tendinopathy: a randomised trial," *Physical Therapy in Sport*, vol. 52, pp. 30–37, 2021.
- [176] A. Scott, R. F. LaPrade, K. G. Harmon et al., "Platelet-rich plasma for patellar tendinopathy: a randomized controlled trial of leukocyte-rich PRP or leukocyte-poor PRP versus saline," *The American Journal of Sports Medicine*, vol. 47, no. 7, pp. 1654–1661, 2019.
- [177] A. L. Sprague, C. Couppé, R. T. Pohlig, L. Snyder-Mackler, and K. G. Silbernagel, "Pain-guided activity modification during treatment for patellar tendinopathy: a feasibility and pilot randomized clinical trial," *Pilot and feasibility studies*, vol. 7, no. 1, p. 58, 2021.
- [178] D. Stasinopoulos and I. Stasinopoulos, "Comparison of effects of exercise programme, pulsed ultrasound and transverse friction in the treatment of chronic patellar tendinopathy," *Clinical Rehabilitation*, vol. 18, no. 4, pp. 347–352, 2004.
- [179] M. Steunebrink, J. Zwerver, R. Brandsema, P. Groenenboom, I. v. d. Akker-Scheek, and A. Weir, "Topical glyceryl trinitrate treatment of chronic patellar tendinopathy: a randomised, double-blind, placebo-controlled clinical trial," *British Journal of Sports Medicine*, vol. 47, no. 1, pp. 34–39, 2013.
- [180] K. M. Thijs, J. Zwerver, F. J. G. Backx et al., "Effectiveness of shockwave treatment combined with eccentric training for patellar tendinopathy," *Clinical Journal of Sport Medicine*, vol. 27, no. 2, pp. 89–96, 2017.
- [181] M. van Ark, J. L. Cook, S. I. Docking et al., "Do isometric and isotonic exercise programs reduce pain in athletes with patellar tendinopathy in-season? A randomised clinical trial," *Journal of Science and Medicine in Sport*, vol. 19, no. 9, pp. 702–706, 2016.
- [182] M. van Ark, E. Rio, J. Cook et al., "Clinical improvements are not explained by changes in tendon structure on ultrasound tissue characterization after an exercise program for patellar tendinopathy," *American Journal of Physical Medicine & Rehabilitation*, vol. 97, no. 10, pp. 708–714, 2018.
- [183] H. van der Worp, J. Zwerver, M. Hamstra, I. van den Akker-Scheek, and R. L. Diercks, "No difference in effectiveness between focused and radial shockwave therapy for treating patellar tendinopathy: a randomized controlled trial," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 22, no. 9, pp. 2026–2032, 2014.
- [184] H. v. Visnes, A. Hoksrud, J. Cook, and R. Bahr, "No effect of eccentric training on Jumper's knee in volleyball players during the competitive season," *Clinical Journal of Sport Medicine*, vol. 15, no. 4, pp. 227–234, 2005.
- [185] C.-J. Wang, J.-Y. Ko, Y.-S. Chan, L.-H. Weng, and S.-L. Hsu, "Extracorporeal shockwave for chronic patellar tendinopathy," *The American Journal of Sports Medicine*, vol. 35, no. 6, pp. 972–978, 2007.
- [186] S. J. Warden, B. R. Metcalf, Z. S. Kiss et al., "Low-intensity pulsed ultrasound for chronic patellar tendinopathy: a randomized, double-blind, placebo-controlled trial," *Rheumatology*, vol. 47, pp. 467–471, 2008.
- [187] M. A. Young, J. L. Cook, C. R. Purdam, Z. S. Kiss, and H. Alfredson, "Eccentric decline squat protocol offers superior results at 12 months compared with traditional eccentric protocol for patellar tendinopathy in volleyball players," *British Journal of Sports Medicine*, vol. 39, no. 2, pp. 102–105, 2005.
- [188] T. Vander Doelen and A. Scott, "Multimodal management of patellar tendinopathy in basketball players: a retrospective chart review pilot study," *Journal of Bodywork and Movement Therapies*, vol. 24, no. 3, pp. 267–272, 2020.
- [189] C. R. Purdam, P. Jonsson, H. Alfredson, R. Lorentzon, J. L. Cook, and K. M. Khan, "A pilot study of the eccentric decline squat in the management of painful chronic patellar tendinopathy," *British Journal of Sports Medicine*, vol. 38, no. 4, pp. 395–397, 2004.
- [190] F. Abat, W. J. Diesel, P. E. Gelber, F. Polidori, J. C. Monllau, and J. M. Sanchez-Ibanez, "Effectiveness of the Intratissue Percutaneous Electrolysis (EPI) technique and isoinertial eccentric exercise in the treatment of patellar tendinopathy at two years follow-up," *Muscles, Ligaments and Tendons Journal*, vol. 4, pp. 188–193, 2014.
- [191] F. Abat, P. E. Gelber, F. Polidori, J. C. Monllau, and J. M. Sanchez-Ibanez, "Clinical results after ultrasound-guided intratissue percutaneous electrolysis (EPI) and eccentric exercise in the treatment of patellar tendinopathy," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 23, no. 4, pp. 1046–1052, 2015.
- [192] Á. Basas, J. Cook, M. A. Gómez et al., "Effects of a strength protocol combined with electrical stimulation on patellar tendinopathy: 42 months retrospective follow-up on 6 high-level jumping athletes," *Physical Therapy in Sport*, vol. 34, pp. 105–112, 2018.
- [193] J. F. Kaux, O. Bruyere, J. L. Croisier, B. Forthomme, C. Le Goff, and J. M. Crielaard, "One-year follow-up of platelet-rich plasma infiltration to treat chronic proximal patellar tendinopathies," *Acta Orthopaedica Belgica*, vol. 81, pp. 251–256, 2015.
- [194] J.-F. o. Kaux, B. Forthomme, and M. H. Namurois, "Description of a standardized rehabilitation program based on sub-maximal eccentric following a platelet-rich plasma infiltration for jumpers knee," *Muscle, Ligaments and Tendons Journal*, vol. 4, pp. 85–89, 2014.
- [195] M. Kongsgaard, K. Qvortrup, J. Larsen et al., "Fibril morphology and tendon mechanical properties in patellar tendinopathy," *The American Journal of Sports Medicine*, vol. 38, no. 4, pp. 749–756, 2010.
- [196] A. S. Panni, M. Tartarone, and N. Maffulli, "Patellar tendinopathy in athletes," *The American Journal of Sports Medicine*, vol. 28, no. 3, pp. 392–397, 2000.
- [197] L. Bianco, J. M. May, S. Fermin, and S. W. Cheatham, "The effect of positional release therapy on intercollegiate male basketball athletes classified with patella tendinopathy," *International Journal of Athletic Therapy & Training*, vol. 24, pp. 1–24, 2019.
- [198] S. Morton, O. Chan, and J. King, "High volume image-guided injections for patellar tendinopathy: a combined retrospective and prospective case series," *Muscles, Ligaments and Tendons Journal*, vol. 4, pp. 214–219, 2014.
- [199] A. C. M. Fernández, C. B. Carballar, J. H. Villafañe et al., "A new ultrasound-guided percutaneous electrolysis and exercise treatment in patellar tendinopathy: three case reports," *Frontiers in Bioscience*, vol. 26, no. 11, pp. 1166–1175, 2021.

- [200] D. Romero-Rodriguez, G. Gual, and P. A. Tesch, "Efficacy of an inertial resistance training paradigm in the treatment of patellar tendinopathy in athletes: a case-series study," *Physical Therapy in Sport*, vol. 12, no. 1, pp. 43–48, 2011.
- [201] S. V. Skovlund, P. Aagaard, P. Larsen, R. B. Svensson, M. Kjaer, and S. P. Magnusson, "The effect of low-load resistance training with blood flow restriction on chronic patellar tendinopathy — a case series," *Translational Sports Medicine*, vol. 3, 2020.
- [202] M. van Ark, I. van den Akker-Scheek, L. T. B. Meijer, and J. Zwerver, "An exercise-based physical therapy program for patients with patellar tendinopathy after platelet-rich plasma injection," *Physical Therapy in Sport*, vol. 14, no. 2, pp. 124–130, 2013.
- [203] T. Cuddeford and J. Brumitt, "In-season rehabilitation program using blood flow restriction therapy for two de-athletes with patellar tendinopathy: a case report," *International Journal of Sports Physical Therapy*, vol. 15, no. 6, pp. 1184–1195, 2020.
- [204] T. L. Dumont, D. L. MacIntyre, and S. R. Harris, "Effects of a six-week eccentric exercise program on patients with patellar tendinopathy: single-subject research study," *Physiotherapie Canada*, vol. 58, no. 2, pp. 130–147, 2006.
- [205] R. B. Goldman and T. Lentz, "The use of eccentric overloading exercise for the treatment of patellar tendinosis in an Olympic-style weightlifter: a case report," *Orthopaedic practice*, vol. 22, p. 10, 2010.
- [206] K. M. McCreesh, S. J. Riley, and J. M. Crotty, "Neovascularity in patellar tendinopathy and the response to eccentric training: a case report using power Doppler ultrasound," *Manual Therapy*, vol. 18, no. 6, pp. 602–605, 2013.
- [207] T. L. Rowan and J. L. Drouin, "A multidisciplinary approach including the use of platelet-rich plasma to treat an elite athlete with patellar tendinopathy - a case report," *Journal of the Canadian Chiropractic Association*, vol. 57, pp. 301–309, 2013.
- [208] R. S. Silva, A. L. G. Ferreira, T. H. Nakagawa, J. E. M. Santos, and F. V. Serrão, "Rehabilitation of patellar tendinopathy using hip extensor strengthening and landing-strategy modification: case report with 6-month follow-up," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 45, no. 11, pp. 899–909, 2015.
- [209] S. Morgan and F. F. Coetzee, "Proposing a patellar tendinopathy screening tool following a systematic review," *South African Journal of Physiotherapy*, vol. 74, no. 1, p. 454, 2018.
- [210] C. P. Hensley and A. J. Emerson Kavchak, "Novel use of a manual therapy technique and management of a patient with peroneal tendinopathy: a case report," *Manual Therapy*, vol. 17, no. 1, pp. 84–88, 2012.
- [211] L. S. Chesterton, M. J. Thomas, and G. Hendry, "Self-management advice, exercise and foot orthoses for plantar heel pain: the TREADON pilot and feasibility randomised trial," *Pilot Feasibility Stud*, vol. 92, 2021.
- [212] E. T. Çil, U. Şaylı, and F. Subaşı, "Outpatient vs home management protocol results for plantar fasciitis," *Foot & Ankle International*, vol. 40, no. 11, pp. 1295–1303, 2019.
- [213] F. Johannsen, L. Konradsen, R. Herzog, and M. R. Krogsgaard, "Endoscopic fasciotomy for plantar fasciitis provides superior results when compared to a controlled non-operative treatment protocol: a randomized controlled trial," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 28, no. 10, pp. 3301–3308, 2020.
- [214] F. E. Johannsen, R. B. Herzog, N. M. Malmgaard-Clausen, M. Hoegberget-Kalisz, S. P. Magnusson, and M. Kjaer, "Corticosteroid injection is the best treatment in plantar fasciitis if combined with controlled training," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 27, no. 1, pp. 5–12, 2019.
- [215] D. H. Kamonseki, G. A. Gonçalves, L. C. Yi, and I. L. Júnior, "Effect of stretching with and without muscle strengthening exercises for the foot and hip in patients with plantar fasciitis: a randomized controlled single-blind clinical trial," *Manual Therapy*, vol. 23, pp. 76–82, 2016.
- [216] N. Rasenberg, S. M. A. Bierma-Zeinstra, L. Fuit et al., "Custom insoles versus sham and GP-led usual care in patients with plantar heel pain: results of the STAP-study - a randomised controlled trial," *British Journal of Sports Medicine*, vol. 55, no. 5, pp. 272–278, 2021.
- [217] H. Riel, B. Vicenzino, M. B. Jensen, J. L. Olesen, S. Holden, and M. S. Rathleff, "The effect of isometric exercise on pain in individuals with plantar fasciopathy: a randomized crossover trial," *Scandinavian Journal of Medicine & Science in Sports*, vol. 25, 2018.
- [218] M. Ryan, J. Hartwell, S. Fraser, R. Newsham-West, and J. Taunton, "Comparison of a Physiotherapy program versus dexamethasone injections for plantar fasciopathy in prolonged standing workers," *Clinical Journal of Sport Medicine*, vol. 24, no. 3, pp. 211–217, 2014.
- [219] S. Thong-On, S. Bovonsunthonchai, R. Vachalathiti, W. Intiravaranont, S. Suwannarat, and R. Smith, "Effects of strengthening and stretching exercises on the temporospatial gait parameters in patients with plantar fasciitis: a randomized controlled trial," *Annals of Rehabilitation Medicine*, vol. 43, no. 6, pp. 662–676, 2019.
- [220] P. C. Wheeler, "The addition of a tension night splint to a structured home rehabilitation programme in patients with chronic plantar fasciitis does not lead to significant additional benefits in either pain, function or flexibility: a single-blinded randomised controlled trial," *BMJ open sport & exercise medicine*, vol. 3, Article ID e000234, 2017.
- [221] H. Riel, J. L. Olesen, M. B. Jensen, B. Vicenzino, and M. S. Rathleff, "Heavy-slow resistance training in addition to an ultrasound-guided corticosteroid injection for individuals with plantar fasciopathy: a feasibility study," *Pilot and Feasibility Studies*, vol. 5, no. 1, p. 105, 2019.
- [222] P. C. Wheeler and C. Dudson, "Similar benefits seen after radial extracorporeal shockwave therapy or autologous blood injection in patients with chronic plantar fasciitis-A retrospective cohort study," *Clinical Journal of Sport Medicine*, vol. 24, 2021.
- [223] B. D. Santos, L. A. Corrêa, L. Teixeira Santos, N. A. M. Filho, T. Lemos, and L. A. C. Nogueira, "Combination of hip strengthening and manipulative therapy for the treatment of plantar fasciitis: a case report," *Journal of Chiropractic Medicine*, vol. 15, no. 4, pp. 310–313, 2016.
- [224] J. H. Lee, J. H. Park, and W. Y. Jang, "The effects of hip strengthening exercises in a patient with plantar fasciitis: a case report," *Medicine*, vol. 98, Article ID e16258, 2019.
- [225] R. Breed, D. Opar, R. Timmins, N. Maniar, H. Banyard, and J. Hickey, "Poor reporting of exercise interventions for hamstring strain injury rehabilitation: a scoping review of reporting quality and content in contemporary applied research," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 45, pp. 1–32, 2021.
- [226] P. C. Charlton, M. K. Drew, B. F. Mentiplay, A. Grimaldi, and R. A. Clark, "Exercise interventions for the prevention

- and treatment of groin pain and injury in athletes: a critical and systematic review,” *Sports Medicine*, vol. 47, no. 10, pp. 2011–2026, 2017.
- [227] C. M. Fairman, P. N. Hyde, and B. C. Focht, “Resistance training interventions across the cancer control continuum: a systematic review of the implementation of resistance training principles,” *British Journal of Sports Medicine*, vol. 51, no. 8, pp. 677–685, 2017.
- [228] C. Minshull and N. Gleeson, “Considerations of the principles of resistance training in exercise studies for the management of knee osteoarthritis: a systematic review,” *Archives of Physical Medicine and Rehabilitation*, vol. 98, no. 9, pp. 1842–1851, 2017.
- [229] C. Bartholdy, S. M. Nielsen, S. Warming, D. J. Hunter, R. Christensen, and M. Henriksen, “Poor replicability of recommended exercise interventions for knee osteoarthritis: a descriptive analysis of evidence informing current guidelines and recommendations,” *Osteoarthritis and Cartilage*, vol. 27, no. 1, pp. 3–22, 2019.
- [230] J. O’Neil, D. McEwen, M. J. Del Bel et al., “Assessment of the content reporting for therapeutic exercise interventions among existing randomized controlled trials on knee osteoarthritis,” *Clinical Rehabilitation*, vol. 32, pp. 980–984, 2018.
- [231] D. Jo, M. J. Del Bel, D. McEwen et al., “A study of the description of exercise programs evaluated in randomized controlled trials involving people with fibromyalgia using different reporting tools, and validity of the tools related to pain relief,” *Clinical Rehabilitation*, vol. 33, no. 3, pp. 557–563, 2019.
- [232] T.-R. Kattackal, S. Cavallo, L. Brosseau et al., “Assessing the reporting quality of physical activity programs in randomized controlled trials for the management of juvenile idiopathic arthritis using three standardized assessment tools,” *Pediatric Rheumatology*, vol. 18, no. 1, p. 41, 2020.
- [233] J. O’Neil, D. McEwen, B. K. Kang et al., “Intervention reporting and dissemination of information for the management of hand osteoarthritis,” *Journal of Hand Therapy*, vol. 34, no. 3, pp. 362–368, 2021.
- [234] M. Charette, M. È. Bérubé, K. Brooks, J. O’Neil, L. Brosseau, and L. McLean, “How well do published randomized controlled trials on pelvic floor muscle training interventions for urinary incontinence describe the details of the intervention? A review,” *Neurourology and Urodynamics*, vol. 39, no. 1, pp. 35–44, 2020.
- [235] S. Giagio, T. Innocenti, S. Salvioli et al., “Completeness of exercise reporting among randomized controlled trials on pelvic floor muscle training for women with pelvic organ prolapse: a systematic review,” *Neurourology and Urodynamics*, vol. 40, no. 6, pp. 1424–1432, 2021.
- [236] B. S. d. Barros, A. M. Imoto, J. O’Neil et al., “The management of lower back pain using pilates method: assessment of content exercise reporting in RCTs,” *Disability & Rehabilitation*, vol. 44, pp. 1–9, 2020.
- [237] S. R. E. Davidson, S. J. Kamper, R. Haskins et al., “Exercise interventions for low back pain are poorly reported: a systematic review,” *Journal of Clinical Epidemiology*, vol. 139, pp. 279–286, 2021.
- [238] A. Vlok, N. van Dyk, D. Coetzee, and H. Grindem, “Exercise descriptors that determine muscle strength gains are missing from reported anterior cruciate ligament reconstruction rehabilitation programs: a scoping review of 117 exercises in 41 studies,” *Journal of Orthopaedic & Sports Physical Therapy*, vol. 24, pp. 1–40, 2021.
- [239] M. P. Reiman, J. Boyd, N. Ingel, A. Reichert, M. Westhoven, and S. Peters, “There is limited and inconsistent reporting of postoperative rehabilitation for femoroacetabular impingement syndrome: a scoping review of 169 studies,” *Journal of Orthopaedic & Sports Physical Therapy*, vol. 50, no. 5, pp. 252–258, 2020.
- [240] L. C. Burgess, T. W. Wainwright, K. A. James, J. von Heiden, and M. D. Iversen, “The quality of intervention reporting in trials of therapeutic exercise for hip osteoarthritis: a secondary analysis of a systematic review,” *Trials*, vol. 22, no. 1, p. 388, 2021.
- [241] I. Burton and A. McCormack, “Resistance training interventions for lower limb tendinopathy: a scoping review of resistance training reporting content, quality, and scientific implementation,” 2022.