Nitrogen Deposition in the Greater Tehran Metropolitan Area

Ali Salahi^{1,*}, Shirin Geranfar², and
Soudabeh A.A. Korori¹

¹Research Institute of Forests and Rangelands (RIFR),
Alborz Research Center, P.O. Box 31585-343, Karaj, Iran,
Fax: +98-21-6603482; ²RIFR, Fax: +98-261-8008406

An investigation of air pollution in the Tehran metropolitan area between 1992–2000 indicated that there are significant amounts of nitrate ion (NO_3^-), over 30 kg/ha/year, deposited as wet deposition, compared to 13 kg/ha/year in the Chitgar Parkland near the Tehran metropolitan area. The amount of NO_3^- in warm seasons is twofold that of cold seasons (see Fig. 1), and there was a significant difference between cold and warm seasons (Table 1). Annual wet deposition of ammonia (NH_3) was 10 kg/ha/year in the Chitgar Parkland[1].

KEY WORDS: nitrate ion (NO_3^-), photochemical reaction, ammonia (NH_3), wet deposition, acidity (pH), suspended particulated matter (SPM)

DOMAINS: plant processes, environmental chemistry, environmental monitoring, analytical chemistry, plant sciences, atmospheric systems, soil systems

INTRODUCTION

Human activities have recently caused the fall of acid rain and polluted precipitation, containing nitrogen oxides (NO_x) and sulfur dioxide (SO_2) which result from the consumption of fossil fuels, condensation of mineral materials, and oil refineries, combined with the atmospheric acids. Acid rain has destructive ef-

fects on forests, causing a decrease in organic matters of soils and increasing soil erosion as well as loss of underground water reservoirs. Acid deposition on vegetation causes their higher susceptibility to pests and pathogen factors[2].

Investigation of acid rain in the area south of Tehran indicates that annual deposition of nitrate ion (NO_3^-) in some southern areas of Tehran is more than 30 kg/ha/year[3], and sulfate ion (SO_4^{2-}) deposition in some southern areas of Tehran is approximately 300 kg/ha/year[4], compared to 80 kg/ha/year in some sampling stations of Chitgar Parkland (Fig. 2). Such huge amounts of air pollutants come down with snow and rain[5].

EXPERIMENTAL METHOD/PROCEDURES

This experiment and study used a method of wet deposition by precipitation. For measurement and sampling, specific instruments were designed and established in different stations south of Tehran, and also on the east and west sides of Tehran. Samples were analyzed in the Analytical Chemistry Laboratory using a flame photometer for potassium and sodium, a spectrophotometer for SO_4^{2-} , and a titration for NO_3^- .

RESULTS AND DISCUSSION

Monthly rainfall indicates that, in spite of the production of NO_x in the cooling season, total nitrate is decreased over winter (Fig. 1). The reason is that production of NO_3^- originates from photochemical processes, and the conditions for photochemical

* Corresponding author.

E-mails: ali.salahi@rifr-ac.org; alisalahi@hotmail.com;

shiringeranfar@hotmail.com

© 2001 with author.

FIGURE 1. Annual wet deposition of NO_3^- in kg/ha/year inside Chitgar Parkland in west greater Tehran metropolitan area during warm and cold seasons.

TABLE 1
Dependent Variable: pH; Independent Variable : NO_3^-

Regression Analysis					
Parameter	Estimate	St. Error	T-Value	Prob.	
Intercept	6.048	0.199	30.457	0.0000	
Slope	0.071	0.035	2.017	0.0522	
Analysis of Variance					
Source	S.S.	D.F.	M.F.	F	Prob.
Model	2.240	1	2.240	4.07	0.0966
Residual	17.620	32	0.551		

Note: $R = 0.34^*$

reactions are not provided in winter. Therefore, production of NO_3^- is decreased (Tables 2 and 3). Amounts of NO_3^- as wet deposition in a number of sampling stations in Tehran was over 30 kg/ha/year, and 13 kg/ha/year for the Chitgar Parkland, 15 km west of the Tehran metropolitan area. The amount of NO_3^- in warm seasons was more than twofold that of cold seasons (Fig. 1). Seasonal sampling of NO_3^- showed that there was significant difference ($p = 0.01$) between cold and warm seasons (Table 3)[1]. In the cold period, increased production of NO_3^- does not parallel the increased consumption of fossil fuels. This means that conditions for photochemical reactions do not occur during winter and cold seasons. Ammonia (NH_3) is formed as a result of the

decomposition of most nitrogenous organic materials used as fertilizer and as a chemical intermediate. Evaporation from animal manure accounts for most of the emissions of NH_3 to the atmosphere. It is, however, the surplus of N in the farming cycle that lies at the bottom of the problem, produced as a result of industrialized farming methods based on the use of artificial fertilizer[6].

Annual wet deposition of NH_3 was 9 kg/ha/year (Fig. 3). Acidity (pH) of precipitation is neutralized by suspended particulate matter (SPM). However, many samples of precipitation showed acid rain (pH = 4) in Tehran. This minimum measured acidity in Tehran is 50 times as acidic as atmospheric background

FIGURE 2. Average amount of SO_4^{2-} wet deposition as kg/ha/year inside precipitation of Chitgar Parkland.

TABLE 2
Analysis of Variance for NO_3^- Related to Precipitation in Chitgar Parkland

K Value	Source	Degrees of Freedom	Sum of Squares	Mean Square	F Value	Prob.
1	Replication	5	213.816	42.763	1.6973	0.1721
2	Factor A	2	50.378	25.189	0.9998	
4	Factor B	1	130.721	130.721	5.1884	0.0316
6	AB	2	99.369	49.684	1.9720	0.1602
-7	Error	25	629.868	25.195		
	Total	35	1124.152			

TABLE 3
Duncan's Multiple Range Test
Related to NO_3^- for Different
Seasons (time) of Precipitation Sampling

Warm seasons	7.551	a
Cold seasons	3.789	b

acidity. Analysis of variance for acidity (pH) related to precipitation in Chitgar Parkland showed that there was significant difference ($p = 0.01$) for different blocks of precipitation sampling (Tables 4 and 5). The regression equation for the independent variable of NO_3^- and dependent variable of acidity (pH) was com-

puted as $Y = 6.048 + 0.071 X$ (Fig. 4). The coefficient of determination was 0.34 ($r = 0.34$). The coefficient of determination for correlation between NO_3^- and SO_4^{2-} was 0.64, between NO_3^- and electro-conductivity (EC) was 0.74, and between NO_3^- and Na^+ was 0.57[7].

FIGURE 3. Average amount of NH_3 inside precipitation of Chitgar Parkland (kg/ha/year).

TABLE 4
Analysis of Variance for Acidity (pH) Related to Precipitation in Chitgar Parkland

K Value	Source	Degrees of Freedom	Sum of Squares	Mean Square	F Value	Prob.
1	Replication	5	4.836	0.967	2.0430	0.1070
2	Factor A	2	6.286	3.143	6.6390	0.0049
4	Factor B	1	0.912	0.912	1.9265	0.1774
6	AB	2	5.097	2.548	5.3830	0.0114
-7	Error	25	11.835	0.473		
	Total	35	28.966			

TABLE 5
Duncan's Multiple Range Test
Related to pH for Different
Treatment of Precipitation Sampling

Near freeway	6.788	a
Center of Parkland	6.488	ab
Iner part of Parkland	5.791	b

FIGURE 4. Regression line for pH (dependent variable) and NO_3^- (independent variable) studied south of Tehran.

CONCLUSION

The main contributor to the concentration of NO_x in urban surroundings is usually road traffic, although in some cities combustion plants make a significant contribution. Animal manure consequently becomes a waste product on farms with intensive stock raising, while those concentrating on grain growing have to make up for a shortage of nutrients by using artificial fertilizer. Since so much of the feed stuff production is now dependent on the use of artificial fertilizer, it can be said that the NH_3 that emanates from animal manure has its origin in the artificial additives. Where farming operations are thus unbalanced, N is continually being added in the form of artificial fertilizer, and leaks out into air and water in great quantities. Hence, the joint strategy puts more emphasis on cutting down NO_x emissions, and more research on health effects is needed.

ACKNOWLEDGEMENTS

The author would like to thank Mrs. Shirin Geranfer for close cooperation, Mr. Hamid Yarmand, former head of the Research Division of Forests and Rangelands Health and Protection, and Professor Mahmoud Shariat, Head of the Environmental Health Group, Tehran University. The author also wishes to thank Mr. Vahid Moniri, Mr. Rasoul Omid, and Mr. Ebrahim Aziz-Khani, colleagues who encouraged me and helped me during research works.

REFERENCES

1. Kojoure, A.S. (2000) Investigation on Air Pollution Affecting Chitgar Parkland Vegetation Cover. Final report. Research Institute of Forests and Rangelands, Karaj, Iran.
2. Smith, W.H. (1990) *Air Pollution and Forests*. 2nd ed. Springer-Verlag, New York.
3. Kojoure, A.S. (1995) *Acid Rain and Their Neutralizing Agents in South of Tehran*. 5th Intl. Conf. Acidic Deposition, 26–30 June 1995, Goteborg, Sweden. Kluwer Academic Publishers, Dordrecht, the Netherlands. 343 p.
4. Kojoure, A.S. (1997) Environmental aspects of acid precipitation quality. *Acid Snow and Rain*. International Congress of ICASR, 6–8 October 1997, Niigata, Japan. pp. 293–298.
5. Fraser, G.A. (1985) The Potential Impacts of the Long Range Transboundary Air Pollutants on Canadian Forests, Information Report E-X-36. Economic Branch, Canadian Forestry Service.
6. Johannesson, M. (1998) Ammonia Also Enters the Picture. *Acid News*, No. 2. The Swedish Society for Conservation of Nature. pp. 20–21.
7. Kojoure, A.S. (1993) *Acid Rain in South of Tehran and Their Intensity and Alternation* [Thesis]. Azad University, Tehran, Iran.
8. Johnson, A.H. (1983) Acid deposition and forest decline. *Environ. Sci. Technol.* **17**(7), 294–305.
9. Kamari, J. (1990) *The Rains Model of Acidification*. Kluwer Academic Publishers, Dordrecht, the Netherlands. pp. 225–231.

This article should be referenced as follows:

Salahi, A., Geranfar, S., and Korori, S.A.A. (2001) Nitrogen deposition in the greater Tehran metropolitan area. In *Optimizing Nitrogen Management in Food and Energy Production and Environmental Protection: Proceedings of the 2nd International Nitrogen Conference on Science and Policy*. *TheScientificWorld* **1**(S2), 261–265.

Received:	July	28, 2001
Revised:	November	11, 2001
Accepted:	November	12, 2001
Published:	December	8, 2001