

Research Article

Mutation at the Human D1S80 Minisatellite Locus

Kuppareddi Balamurugan,¹ Martin L. Tracey,² Uwe Heine,³
George C. Maha,³ and George T. Duncan⁴

¹ School of Criminal Justice, University of Southern Mississippi, 118 College Drive # 5127, Hattiesburg, MS 39406, USA

² Department of Biology, Florida International University, University Park Campus, Miami, FL 33199, USA

³ DNA Identification Testing Division, Laboratory Corporation of America, 1440 York Court Extension, Burlington, NC 27215, USA

⁴ Broward County Sheriff's Office, Forensic Laboratory DNA Unit, Fort Lauderdale, FL 33301, USA

Correspondence should be addressed to Kuppareddi Balamurugan, kuppareddi.balamurugan@usm.edu

Received 30 October 2011; Accepted 5 January 2012

Academic Editors: P. Momigliano Richiardi, P. Y. Woon, and N. Zhang

Copyright © 2012 Kuppareddi Balamurugan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Little is known about the general biology of minisatellites. The purpose of this study is to examine repeat mutations from the D1S80 minisatellite locus by sequence analysis to elucidate the mutational process at this locus. This is a highly polymorphic minisatellite locus, located in the subtelomeric region of chromosome 1. We have analyzed 90,000 human germline transmission events and found seven (7) mutations at this locus. The D1S80 alleles of the parentage trio, the child, mother, and the alleged father were sequenced and the origin of the mutation was determined. Using American Association of Blood Banks (AABB) guidelines, we found a male mutation rate of 1.04×10^{-4} and a female mutation rate of 5.18×10^{-5} with an overall mutation rate of approximately 7.77×10^{-5} . Also, in this study, we found that the identified mutations are in close proximity to the center of the repeat array rather than at the ends of the repeat array. Several studies have examined the mutational mechanisms of the minisatellites according to infinite allele model (IAM) and the one-step stepwise mutation model (SMM). In this study, we found that this locus fits into the one-step mutation model (SMM) mechanism in six out of seven instances similar to STR loci.

1. Introduction

The human genome can be grossly partitioned into three categories: nonrepetitive (single copy sequences), moderately repetitive (families of retroposon-like sequences), and highly repetitive ("classical" satellite) DNA [1]. Minisatellites represent a class of tandem repeats with repeat units ranging from 6 to 100 bp and fall into the class of highly repetitive satellites. Total array sizes range from 0.5 to 30 kb. Minisatellite loci are found throughout the genome and are estimated to number in the thousands with a strong telomeric and strand bias [2]. They are generally clustered in subtelomeric and centromeric regions; however, it has been suggested that minisatellites show no obvious predisposition for accumulation in subtelomeric or centromeric regions but are associated with rates of high recombination and mutation in these regions [2–9].

Studies have shown that minisatellites may mutate by unexpectedly complex conversion-like events [10]. Higher

mutational polar bias occurs at the termini at some, if not most minisatellites, as well as microsatellite loci [11]. Minisatellite loci generally are variable near one end of the array or in some cases both ends. It has been suggested that this implies a mutational mechanism driven by cis-acting elements and thus the existence of a localized mutation hotspot at one end of the array involving complex processes of gene conversion [12–14]. One caveat is that the control regions have not been identified and only associated flanking region markers have been found.

Little is known about the general biology of minisatellites [15]. The purposes of the study are (1) to examine the mutational rate and (2) to examine mutations from the D1S80 minisatellite locus by sequence analysis to elucidate the mutational process, as an example of a particular type of minisatellite array. The highly polymorphic minisatellite locus D1S80, (gene location: 1p36.32, GenBank sequence Accession # D28507) was first described by Nakamura in 1989. It is a minisatellite locus composed of short repeat units,

TABLE 1: Nucleotide sequences of observed repeat units. The consensus sequence represents the most common nucleotide observed in each position of the repeats. Twenty variations based on the consensus sixteen base repeat unit are tabulated. Each repeat unit is assigned a letter code. Dots (·) represent a match to the consensus sequence as represented by type H repeat unit. Letters represent nucleotide differences when compared to the consensus sequence and correspond to A, G, C, and T nucleotides. A (-) represents a missing nucleotide in the repeat unit A.

Repeat type		Sequence															
Type A	T	C	A	·	C	·	·	·	·	A	·	·	·	·	·	·	·
Type B	A	C	A	·	·	·	·	·	·	A	·	·	·	·	·	·	·
Type C	·	·	·	·	·	·	·	·	·	·	·	·	A	·	·	·	·
Type D	·	·	A	·	·	·	·	·	·	·	·	·	A	·	·	·	·
Type E	·	·	A	·	·	·	·	·	·	A	·	·	·	·	·	·	·
Type F	·	·	·	·	·	·	·	·	·	A	·	·	·	·	·	·	·
Type G	·	·	A	·	·	·	·	·	·	·	·	·	·	·	·	·	·
Type H	G	A	G	G	A	C	C	A	C	C	G	G	C	A	A	G	(Consensus)
Type I	·	·	·	·	·	·	·	·	·	·	A	·	G	·	·	·	·
Type J	·	·	·	A	·	·	·	·	·	·	A	·	G	·	·	·	·
Type K	·	·	·	·	·	·	·	·	·	·	A	·	·	·	·	·	·
Type L	·	·	·	·	·	·	·	·	·	T	·	·	·	·	·	·	·
Type M	·	·	·	·	·	·	·	·	·	·	A	·	G	·	G	·	·
Type N	·	·	·	·	·	·	·	·	·	G	·	·	·	·	·	·	·
Type O	·	·	·	·	·	·	·	·	·	·	·	·	·	G	·	·	·
Type P	·	·	A	·	·	·	·	·	·	·	·	·	A	·	G	·	·
Type Q	·	·	A	·	·	·	·	·	G	·	·	·	A	·	·	·	·
Type R	·	G	·	A	·	·	·	·	·	·	A	·	G	·	·	·	·
Type S	·	·	A	·	·	·	·	·	·	·	A	·	G	·	·	·	·
Type T	·	·	A	·	·	·	·	·	·	·	A	·	·	·	·	·	·
Type U	·	·	A	·	·	·	·	·	·	·	A	·	A	·	·	·	·




FIGURE 1: The general structure of the D1S80 locus is that of a monomeric type VNTR consisting of four 5' repeat units (A-B-C-D) and seven 3' repeat units (H-I-J-I-I-L-G) that are constant (motifs), with variable number of repeats in between. These two motifs are essentially identical in almost all alleles sequenced in all population groups so far (data not shown). There is a 132 base pair 5' flanking region which includes the forward PCR primer (P1) and a 32-base pair 3' flanking region which includes the reverse PCR primer (P2) sequence. The two-flanking restriction site polymorphism, *Hinf*I (G↓ANTC), 58 base pairs from the 5' end of the primer, and the *Fnu*4HI (GC↓NGC), the first base after the last repeat are shown by arrows.

typically less than or equal to 16 base pairs per unit [16, 17] (Table 1, Figure 1). It is located in the sub-telomeric region of chromosome 1 (chr1:2,390,716 - 2,391,193) roughly 2.4M bases from the p-telomere [18, 19]. The total range of repeat lengths (approx. 0.2 to 1kb) and the relatively small number of repeats (generally 14 to >40) within the unit make this locus suitable for PCR analysis [15, 20] (Figure 1). The observed heterozygosity has been reported as high as 90.5% and as low as 24% with greater than 27 alleles [21]. In most human populations, 18 and 24 repeat units are the most common and possibly the primordial alleles at this locus [22]. Because of the complexity of the repeat array structure, differences between the alleles become more evident when

the arrays of repeat units are divided into common motifs. Six motifs ranging from three to nine repeat units have been identified. Allele frequencies and motifs characterized by the 18 and 24 structural alleles were described earlier [23, 24]. The D1S80 locus was previously used for forensic analysis due to its small size; however, this locus has recently been used as an associative tool in the elucidation of the chromosome 1p36 deletion syndrome which is one of the most common deletion syndromes located near a chromosomal terminus (one in 5000 births) [15, 25]. Recent and past work with respect to this locus has included the use of two single nucleotide polymorphisms (SNPs). These included rs16824398, which is a SNP that

involves a *Hinf*I restriction site at a nucleotide 58 bases downstream from the forward primer and another SNP in the 3' flanking region that involves an *Fnu*4HI restriction site, at the base next to the last repeat. Interestingly, all 18-repeat alleles to date have been found to be associated with *Hinf*I(+) and *Fnu*4HI(-) restriction site polymorphisms at the 5' and 3' ends, respectively (Figure 1). On the other hand, allele 24 is associated with *Hinf*I(-) and *Fnu*4HI(+) polymorphism. Of the alleles tested, 98.5% exhibits linkage disequilibrium between two specific SNPs. If an allele is positive for *Hinf*I, then it is negative for *Fnu*4HI, and if an allele is negative for *Hinf*I, it is then positive for *Fnu*4HI. There is very strong linkage disequilibrium between the two SNPs [23, 26, 27]. In this study, we report the mutation rate and a nonpolarized mutational mechanism at the D1S80 minisatellite locus.

2. Materials and Methods

2.1. Sample Collection and Parentage Analysis. Buccal epithelial cell samples from 45,000 trios for paternity analysis were collected in quadruplicate from each individual using cotton-tipped swabs and air dried at ambient temperature. Each paternity short tandem repeat (STR) assay was done using duplicate independently prepared DNA extractions and assayed using proprietary inhouse STR multiplexes. Each multiplex included a shared locus (generally D1S80) with other multiplexes to confirm sample identity. D1S80 was assayed in combination with two to three other STR loci in multiplex reactions. The other STRs were selected to produce lower molecular weight fragments that would not interfere with analysis of the larger D1S80 amplified products. Amplified products were analyzed using nondenaturing high-resolution polyacrylamide gel electrophoresis with silver stain detection. Alleles were manually determined by comparison with allelic ladders containing all common D1S80 alleles. All STR-based parentage testing was performed with full knowledge and consent of the tested individuals or authorized parent/guardian. Appropriate institutional review board approval was obtained for sequence analysis of the samples.

2.2. Parentage and Mutation Analysis. To fully understand the implications and delineate the mutational mechanisms that the D1S80 locus undergoes, we examined the number of parent child allele transfers analyzed at this locus for the year 1996 from Laboratory Corporation of America, Burlington, NC. Observed mutations were confirmed as coming from concordant father or mother by analyzing short tandem repeat (STR) data from samples collected from alleged father, mother, and child using Ampf/STR Identifier human identification kit (Applied Biosystems, Foster City, CA). This was essential to confirm that the mutation occurred in the D1S80 locus and was not the result of nonpaternity.

Several factors were used in the elucidation of the origin of the mutation between the alleged father and mother of the child. The first factor was the phase of the SNPs at the 5'- and 3'-flanking sequence of the repeat array. Another factor was a match of the repeat units of the alleles most likely

to be associated between father and mother and finally we assumed that the minimal size change was the most probable one.

A mutation rate was then estimated by the number of samples sequenced that contain the mutation and compared to the number of trios sampled that year at Laboratory Corporation of America. Parentage calculations and mutation rate analysis were performed following guidelines put forth by the American Association of Blood Banks (AABB) [28].

2.3. DNA Analysis and Sequencing. Total genomic DNA was isolated from the trios that contained the mutation by standard organic extraction with phenol/chloroform/isoamyl alcohol followed by purification and concentration using Amicon Ultracel centrifugal filters (Millipore Corporation, Billerica, MA, USA). Quantitation of total genomic DNA was performed using Quantifiler human DNA quantitation kit (Applied Biosystems, Foster City, CA, USA). Paternity was confirmed by analyzing a set of fifteen short tandem repeat (STR) markers using the Identifier human identification kit (Applied Biosystems, Foster City, CA, USA) as per manufacturer's recommendations. The samples were run on an automated 310 Genetic Analyzer and analyzed with GeneMapper ID software version 3.2 (Applied Biosystems, Foster City, CA, USA). The D1S80 locus was amplified using primers described by Kasai et al. [16]. Alleles were separated by 5% polyacrylamide gels and stained with ethidium bromide. Individual allelic bands were excised from the gel using disposable razor blades and suspended in 100 μ L of Tris-EDTA (TE) buffer overnight. One to five μ L of the gel extract was used for amplification of single alleles using the same primers described above. The alleles were sequenced using big dye terminator chemistry and ABI 310 genetic analyzer (Applied Biosystems, Foster City, CA, USA). Homozygous samples or samples that did not yield clean sequence were cloned into pBluescript plasmid vector. The DH5 α competent cells were transformed using the recombinant vector and the cells were grown in LB/ampicillin agar plates. The resulting colonies were screened for the presence of the right size insert by directly sequencing the plasmid DNA isolated from minipreps of individual colonies. Sequencing of the cloned alleles was performed using big dye terminator chemistry and the sequence was analyzed using Applied Biosystems DNA sequencing analysis software version 5.2.

3. Results and Discussion

Paternity calculations were performed using AABB guidelines [28] and a residual likelihood ratio from a low of 180,848 to a high of 190,926,913,150 was observed. Mutations were identified as non-Mendelian inheritance of alleles differing in size in seven (7) out of a total of 45,000 trios (Table 2). Most of the mutations (6/7) involved a change of one repeat unit (Table 2). Although one mutation involved a two-repeat deletion, our observations support the strict step-wise replication slippage mutation (SSM) model according to which the majority of the mutational events at microsatellite loci involve gain or loss of one-repeat unit rather than the complex mutational events as illustrated by

TABLE 2: DNA sequence characteristics of seven trios illustrating the mutational event. The standard paternity test, called a trio, involves the child, mother, and alleged father. Allele represents the number of 16 base pairs repeat units making up the entire allele. The alphabets in each sample represent the arrangement of the repeat types making up each allele. The alphabet code for the repeat types and the corresponding bases is given in Table 1. HinfI and Fnu4HI are the two restriction site polymorphisms located on the 5' and 3' flanking regions of the repeats, respectively, and are concordant between the progenitor and the mutated allele. The star (*) represents the mutated allele and aligned with the progenitor wherever possible. The progenitors are identified based on the sequence characteristic, SNP polymorphism, and the smallest allele difference between the parent and child. For the four indeterminate mutations, the mutated allele is aligned with both the possible progenitors.

	Sample	Allele	HinfI	Repeat types	Fnu4HI	Comments
1Trio						
1	Mother-18 allele 1	18	Positive	A B C D D E C H H [*]	I I H I J I I L G	Negative Possible progenitor
1	* Child-17 allele 1	17	Positive	A B C D D E C H	I I H I J I I L G	Negative Deletion of one repeat unit
1	Father-18 allele 1	18	Positive	A B C D D E C H H [*]	I I H I J I I L G	Negative Possible progenitor
1	Child-18 allele 2	18	Positive	A B C D D E C H H	I I H I J I I L G	Negative
1	Mother-18 allele 2	18	Positive	A B C D D E C H H	I I H I J I I L G	Negative
1	Father-22 allele 2	22	Positive	A B C D D E C H H H I	J I I H I J I I L G	Negative
2Trio						
3	Father-18 allele 1	18	Positive	A B C D D E C H H	I I H I J I I L G	Negative Possible progenitor
3	* Child-19 allele 2	19	Positive	A B C D D D [*]	E C H H I I H I J I I L G	Negative Duplication of one repeated unit
3	Mother-18 allele 1	18	Positive	A B C D D E C H H	I I H I J I I L G	Negative Possible progenitor
3	Child-18 allele 1	18	Positive	A B C D D E C H H	I I H I J I I L G	Negative
3	Mother-25 allele 2	25	Negative	A B C D D D E F C G H	I I I J H I J I I L G	Positive
3	Father-31 allele 2	31	Positive	A B C D D E F C G H	I I H I I K I K I I I L G	Negative
3Trio						
7	* Child-19 allele 1	19	Negative	A B C D E F C G H	I I J H I J I I L G	Positive Deletion of one repeated unit
7	Father-20 allele 1	20	Negative	A B C D E F C G H	I I I I I I L G	Positive Progenitor
7	Child-24 allele 2	24	Negative	A B C D D E F C G H	I I I J H I J I I L G	Positive
7	Mother-24 allele 1	24	Negative	A B C D D E F C G H	I I I J H I J I I L G	Positive Donor, child allele 2
7	Father-24 allele 2	24	Negative	A B C D D E F C G H	I I I J H I J I I L G	Positive
7	Mother-31 allele 2	31	Positive	A B C D D E F C G H	I I H K I K H K I K I I I L G	Negative
4Trio						
8	* Child-21 allele 1	21	Negative	A B C D E F F [*]	C G H I I I J H I J I I L G	Positive Duplication of one repeated unit
8	Mother-20 allele 1	20	Negative	A B C D E F	C G H I I I J H I J I I L G	Positive Progenitor
8	Child-31 allele 2	31	Positive	A B C D D E F C G H	I I H I I K I K I I I L G	Negative
8	Father-31 allele 2	31	Positive	A B C D D E F C G H	I I H I I K I K I I I L G	Negative Donor, child allele 2
8	Father-26 allele 1	26	Positive	A B C D D E F C G H	I I H I I S J I I L G	Negative
8	Mother-31 allele 2	31	Positive	A B C D D E F C G H	I I H K I K H K I K I I I L G	Negative

TABLE 2: Continued.

Sample					Repeat types			Fnu4HI		Comments
	Sample	Allele	HinfI	Repeat types						
5Trio	10	Mother-28 allele 2	28	Positive	A B C D D E C G H	I K K I I [¶] H [¶]	I I H I J I H I J I I L G	Negative	Possible progenitor	
	10	*Child-26 allele 1	26	Positive	A B C D D E C G H	I K K I	I I H I J I H I J I I L G	Negative	Deletion of two repeated units	
	10	Father-28 allele 1	28	Positive	A B C D D E C G H	I K K I I [¶] H [¶]	I I H I J I H I J I I L G	Negative	Possible progenitor	
	10	Child-28 allele 2	28	Positive	A B C D D E C G H	I K K I I H I I H I J I H I J I I L G	Negative			
	10	Mother-24 allele 1	24	Positive	A B C D D E C G H	I K K I I K I I H I J I I L G	Negative	Mother's sequence does not match child 26		
	10	Father-31 allele 2	31	Positive	A B C D D E F C G H	I I H K I K H K I K I K I I H I J I I L G	Negative			
	6Trio	11	Mother-24 allele 2	24	Negative	A B C D D E F C G H [¶]	I I I J H I J H I J I I L G	Positive	Possible progenitor	
11		*Child-23 allele 1	23	Negative	A B C D D E F C G	I I I J H I J H I J I I L G	Positive	Deletion of one repeated unit		
11		Father-24 allele 1	24	Negative	A B C D D E F C G H [¶]	I I I J H I J H I J I I L G	Positive	Possible progenitor		
11		Child-24 allele 2	24	Negative	A B C D D E F C G H	I I I J H I J H I J I I L G	Positive			
11		Mother-18 allele 1	18	Positive	A B C D D E C H I	I H I J I I L G	Negative			
11		Father-31 allele 2	31	Positive	A B C D D E F C G H	I I H K I K H K I K I K I I H I J I I L G	Negative			
7Trio		12	*Child-19 allele 1	19	Positive	A B C D D E C H H	I I H I J I [¶] I I L G	Negative	Duplication of one repeated unit	
	12	Father-18 allele 1	18	Positive	A B C D D E C H H	I I H I J I I L G	Negative	Progenitor		
	12	Child-20 allele 2	20	Negative	A B C D E F C G H	I I I J H I J I I L G	Positive			
	12	Mother-20 allele 2	20	Negative	A B C D E F C G H	I I I J H I J I I L G	Positive	Donor, child allele 2		
	12	Mother-18 allele 1	18	Positive	A B C D D E C H I	I H I J I I L G	Negative			
	12	Father-31 allele 2	31	Positive	A B C D D E F C G H	I I H I H I I K I K I K I I H I J I I L G	Negative			

[¶] Represent the mutated region of the parental and child allele.

other minisatellites. We assumed that the smallest size change was most probable, thus the most parsimonious changes were assumed in our analysis of the mutational events [29, 30]. The phase of the *HinfI* and *Fnu4HI* restriction site polymorphisms of all mutated alleles of the children was concordant to the phase of the polymorphisms of the progenitor. Alec Jeffreys, who has examined these mutational events in detail, has stated that replication slippage and unequal crossover events which are intrinsic to the tandem repeat array are not primarily the mechanisms of mutation. Instead, he postulated that germline repeat instability which is regulated by cis-acting elements somewhere near the array may be responsible [31]. In our data, in four trio sets, it was not possible to ascertain which parent was the progenitor of the mutation. In the remaining three trios, we were able to ascertain two mutations to the father and one to the mother. Other studies have emphasized the origin of most mutations from paternal lineage due to more numbers of meiotic events [32–34], but we cannot make this prediction from our data since four of the seven mutations are indeterminate. The advantage of samples like this permits the analysis of true *in vivo* germline events rather than somatic mutational events.

Only two studies are reported that have dealt with the mutational events within this locus [35–37]. The first study [35] reported three mutations involving the loss or gain of three repeats to as much as a loss of six repeats. Of these events, two could be attributed to a maternal origin and the remaining event could not be discerned between mother and father. The data of our study does not support these findings where most mutational events involved one repeat loss or gain. Their conclusion was that paternal events were most prevalent in microsatellite loci, while in minisatellite loci maternal events were most prevalent. The second study dealt with computer simulations and a stepwise mutation model [37]. Jeffreys et al. found that minisatellites do not mutate by replication slippage and concluded that most mutations were exhibited in the male germline [14].

Mutation rates for minisatellites have been estimated to range from 0.5% to greater than 20% per generation in some studies and in another study the rate was reported as 0.53×10^{-3} to 1.53×10^{-3} [35]. Only one study up to this time has examined the D1S80 locus specifically and found a mutation rate of 3 out of 6153 meioses or 5×10^{-4} [35]. We found 7 mutations out of 90,000 meioses (45,000 paternal and 45,000 maternal meioses). This then represented an effective overall mutation rate of approximately 7.77×10^{-5} . The mutation rate of 7.77×10^{-5} represents the overall mutation rate rather than a rate specific to the male or female germline. Only one event could be narrowed down to the mother and two events attributable to the father. In the remaining four trios, we could not choose the progenitor because of the similarities of the maternal and paternal sequences. For the four indeterminate mutations, the ABB prorate them based on known mutation rates [28]. In this case, one adds 2.667 to the males and 1.333 to the female for estimated mutation rates of 1.04×10^{-4} (4.667/45,000) and 5.18×10^{-5} (2.333/45,000), respectively. Some mutations among the trios might not appear as Mendelian errors in the analysis since the scoring was based on a size difference between

alleles (difference in the number of repeats between parent and child), and single base changes would not be detected, and the actual mutation rates could be slightly higher.

Several studies have examined mutational mechanisms according to infinite allele model (IAM) and the one-step stepwise mutation model (SMM). They have found that minisatellites sometimes fit both mechanisms; however, short tandem repeat loci (STR) were most similar to the simulation results under the SMM model of mutation. In this study, we found that this locus fit into the one-step stepwise mutation model (SMM) mechanism in six out of seven instances similar to STR loci [31, 37, 38]. Jeffreys, when using single-molecule polymerase chain reaction (SP-PCR) for elucidation of the minisatellite MS1, found an abundance of mutational events in sperm consisting of the deletion or addition of one-repeat unit [38].

Jeffreys et al. [31] reported polarized mutational events for the three minisatellite loci MS32, MS205, and MS31A. In this study, the authors reported that 93% of mutants lay within 20 repeats of the progenitor allele and 90% of the mutant alleles which had gained repeats showed extreme polarity, with repeat segments being added in the 5' region of the progenitor allele. This finding is in contrast to our results in this study. All the mutated alleles have deletion or duplication in close proximity to the center of the repeats, while four repeats in the 5' end and seven repeats in the 3' end are relatively constant in both the progenitor and the mutated alleles.

In addition to the differences in the repeat regions of the progenitor and mutated alleles, Jeffreys et al. have also found that the polymorphisms flanking the minisatellite repeat array have failed to reveal exchange of flanking markers [31]. This is true with the results obtained from this study that the 5' *HinfI* and the 3' *Fnu4HI* restriction site polymorphisms are concordant with the progenitor and the mutated alleles. This phenomenon suggests that the 5'- and 3'-flanking regions and at least four repeats on either end of the repeat array are not involved in the mutational process. These differences in the region of the mutation in the repeat array indicate possible different mutational mechanisms in different minisatellite loci.

The data we have reported highlights the importance of a number of questions. (1) Do repeats have functions and do they have an effect on genome structure? (2) Are the allelic clades the result of constraints on mutation? (3) Is selection acting at the D1S80 locus?, and (4) Why is the D1S80 locus conserved among primates (unpublished data) including humans?

Conflict of Interests

The authors declare no conflict of interests, financially or otherwise.

Acknowledgments

The authors would like to thank Mrs Yasotha Balasubramaniam, Toronto, ON, Canada for the artwork and the four anonymous reviewers for their comments and suggestions.

This study was partly supported by funding provided by the National Institute of Justice, Department of Justice, Washington, DC, USA, under project number 2009-D1-BX-0293 to the University of Southern Mississippi and by a startup grant for one of the authors (K. Balamurugan) from the University of Southern Mississippi, Hattiesburg, MS, USA. Points of view in the document are those of the authors and do not necessarily represent the official view of the U.S. Department of Justice.

References

- [1] B. Lewin, *Genes IV*, Oxford University Press, New York, NY, USA, 1990.
- [2] G. F. Richard, A. Kerrest, and B. Dujon, "Comparative genomics and molecular dynamics of DNA repeats in eukaryotes," *Microbiology and Molecular Biology Reviews*, vol. 72, no. 4, pp. 686–727, 2008.
- [3] D. L. Ellsworth, M. D. Shriver, and E. Boerwinkle, "Nucleotide sequence analysis of the apolipoprotein B 3' VNTR," *Human Molecular Genetics*, vol. 4, no. 5, pp. 937–944, 1995.
- [4] R. Saferstein, *Forensic Science Handbook*, Regents/Prentice Hall, Englewood Cliffs, NJ, USA, 1993.
- [5] J. A. L. Armour and A. J. Jeffreys, "Biology and applications of human minisatellite loci," *Current Opinion in Genetics and Development*, vol. 2, no. 6, pp. 850–856, 1992.
- [6] J. A. L. Armour, P. C. Harris, and A. J. Jeffreys, "Allelic diversity at minisatellite MS205 (D16S309): evidence for polarized variability," *Human Molecular Genetics*, vol. 2, no. 8, pp. 1137–1145, 1993.
- [7] A. J. Jeffreys, V. Wilson, and S. L. Thein, "Hypervariable 'minisatellite' regions in human DNA," *Nature*, vol. 314, no. 6006, pp. 67–73, 1985.
- [8] W. Stephan, "Tandem-repetitive noncoding DNA: forms and forces," *Molecular Biology and Evolution*, vol. 6, no. 2, pp. 198–212, 1989.
- [9] G. P. Smith, "Evolution of repeated DNA sequences by unequal crossover," *Science*, vol. 191, no. 4227, pp. 528–535, 1976.
- [10] P. R. J. Bois, "Hypermutable minisatellites, a human affair?" *Genomics*, vol. 81, no. 4, pp. 349–355, 2003.
- [11] C. B. Kunst and S. T. Warren, "Cryptic and polar variation of the fragile X repeat could result in predisposing normal alleles," *Cell*, vol. 77, no. 6, pp. 853–861, 1994.
- [12] A. J. Jeffreys, A. MacLeod, K. Tamaki, D. L. Neil, and D. Monckton, "Minisatellite repeat coding as a digital approach to DNA typing," *Nature*, vol. 354, no. 6350, pp. 204–209, 1991.
- [13] R. K. Wolff, R. Plaetke, A. J. Jeffreys, and R. White, "Unequal crossover between homologous chromosomes is not the major mechanism involved in the generation of new alleles at VNTR loci," *Genomics*, vol. 5, no. 2, pp. 382–384, 1989.
- [14] A. J. Jeffreys, M. J. Allen, J. A. L. Armour et al., "Mutation processes at human minisatellites," *Electrophoresis*, vol. 16, no. 9, pp. 1577–1585, 1995.
- [15] I. C. Gray and A. J. Jeffreys, "Evolutionary transience of hyper-variable minisatellites in man and the primates," *Proceedings of the Royal Society B*, vol. 243, no. 1308, pp. 241–253, 1991.
- [16] K. Kasai, Y. Nakamura, and R. White, "Amplification of a variable number of tandem repeats (VNTR) locus (pMCT118) by the polymerase chain reaction (PCR) and its application to forensic science," *Journal of Forensic Sciences*, vol. 35, no. 5, pp. 1196–1200, 1990.
- [17] Y. Nakamura, M. Carlson, K. Krapcho, and R. White, "Isolation and mapping of a polymorphic DNA sequence (pMCT118) on chromosome 1p [D1S80]," *Nucleic Acids Research*, vol. 16, no. 19, p. 9364, 1988.
- [18] W. James Kent, C. W. Sugnet, T. S. Furey et al., "The human genome browser at UCSC," *Genome Research*, vol. 12, no. 6, pp. 996–1006, 2002.
- [19] W. J. Kent, "BLAT—the BLAST-like alignment tool," *Genome Research*, vol. 12, no. 4, pp. 656–664, 2002.
- [20] B. Budowle, R. Chakraborty, A. M. Giusti, A. J. Eisenberg, and R. C. Allen, "Analysis of the VNTR locus D1S80 by the PCR followed by high-resolution PAGE," *American Journal of Human Genetics*, vol. 48, no. 1, pp. 137–144, 1991.
- [21] A. Sajantila, B. Budowle, M. Strom et al., "PCR amplification of alleles at the D1S80 locus: comparison of a Finnish and a North American Caucasian population sample, and forensic casework evaluation," *American Journal of Human Genetics*, vol. 50, no. 4, pp. 816–825, 1992.
- [22] R. Deka, S. DeCroo, L. Jin et al., "Population genetic characteristics of the D1S80 locus in seven human populations," *Human Genetics*, vol. 94, no. 3, pp. 252–258, 1994.
- [23] K. Balamurugan, R. Pomeroy, G. Duncan, and M. Tracey, "Investigating SNPs flanking the D1S80 locus in a Tamil population from India," *Human Biology*, vol. 82, no. 2, pp. 221–226, 2010.
- [24] K. Balamurugan, N. Prabakaran, G. Duncan, B. Budowle, M. Tahir, and M. Tracey, "Allele frequencies of 13 STR loci and the D1S80 locus in a Tamil population from Madras, India," *Journal of Forensic Sciences*, vol. 46, no. 6, pp. 1515–1517, 2001.
- [25] S. K. Shapira, C. McCaskill, H. Northrup et al., "Chromosome 1p36 deletions: the clinical phenotype and molecular characterization of a common newly delineated syndrome," *American Journal of Human Genetics*, vol. 61, no. 3, pp. 642–650, 1997.
- [26] G. T. Duncan, K. Balamurugan, B. Budowle, and M. L. Tracey, "Hinf I/Tsp509 I and BsoF I polymorphisms in the flanking regions of the human VNTR locus D1S80," *Genetic Analysis*, vol. 13, no. 5, pp. 119–121, 1996.
- [27] S. A. Limborska, A. V. Khrunin, O. V. Flegontova, V. A. Tasitz, and D. A. Verbenko, "Specificity of genetic diversity in D1S80 revealed by SNP-VNTR haplotyping," *Annals of Human Biology*, vol. 38, no. 5, pp. 564–569, 2011.
- [28] Annual Report Summary for Testing in 2008, Prepared by the Relationship Testing Program Unit, <http://www.aabb.org/sa/facilities/Documents/rtannrpt08.pdf>.
- [29] J. J. Wiens and M. R. Servedio, "Phylogenetic analysis and intraspecific variation: performance of parsimony, likelihood, and distance methods," *Systematic Biology*, vol. 47, no. 2, pp. 228–253, 1998.
- [30] E. Sober, "The contest between parsimony and likelihood," *Systematic Biology*, vol. 53, no. 4, pp. 644–653, 2004.
- [31] A. J. Jeffreys, K. Tamaki, A. MacLeod, D. G. Monckton, D. L. Neil, and J. A. L. Armour, "Complex gene conversion events in germline mutation at human minisatellites," *Nature Genetics*, vol. 6, no. 2, pp. 136–145, 1994.
- [32] K. Tamaki, C. H. Brenner, and A. J. Jeffreys, "Distinguishing minisatellite mutation from non-paternity by MVR-PCR," *Forensic Science International*, vol. 113, no. 1–3, pp. 55–62, 2000.
- [33] Y. E. Dubrova, G. Grant, A. A. Chumak, V. A. Stezhka, and A. N. Karakasian, "Elevated minisatellite mutation rate in the post-chernobyl families from Ukraine," *American Journal of Human Genetics*, vol. 71, no. 4, pp. 801–809, 2002.
- [34] A. J. Jeffreys, P. Bois, J. Buard et al., "Spontaneous and induced minisatellite instability," *Electrophoresis*, vol. 18, no. 9, pp. 1501–1511, 1997.

- [35] A. Sajantila, M. Lukka, and A. C. Syvänen, "Experimentally observed germline mutations at human micro- and minisatellite loci," *European Journal of Human Genetics*, vol. 7, no. 2, pp. 263–266, 1999.
- [36] M. D. Shriver, *Origins and evolution of VNTR loci: the apolipoprotein B 3' VNTR*, Ph.D. thesis, The University of Texas Health Science Center at Houston, Houston, Tex, USA , 1993.
- [37] M. D. Shriver, L. Jin, R. Chakraborty, and E. Boerwinkle, "VNTR allele frequency distributions under the stepwise mutation model: a computer simulation approach," *Genetics*, vol. 134, no. 3, pp. 983–993, 1993.
- [38] I. Berg, R. Neumann, H. Cederberg, U. Rannug, and A. J. Jeffreys, "Two modes of germline instability at human minisatellite MS1 (locus D1S7): complex rearrangements and paradoxical hyperdeletion," *American Journal of Human Genetics*, vol. 72, no. 6, pp. 1436–1447, 2003.

