

Research Article Strongly Lacunary Ward Continuity in 2-Normed Spaces

Hüseyin Çakalli¹ and Sibel Ersan²

¹ Department of Mathematics, Maltepe University, Marmara Eğitim Köyü, 34857 İstanbul, Turkey

² Faculty of Engineering and Natural Sciences, Maltepe University, Marmara Eğitim Köyü, 34857 İstanbul, Turkey

Correspondence should be addressed to Hüseyin Çakalli; huseyincakalli@maltepe.edu.tr

Received 19 April 2014; Accepted 6 June 2014; Published 23 June 2014

Academic Editor: S. A. Mohiuddine

Copyright © 2014 H. Çakalli and S. Ersan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A function f defined on a subset E of a 2-normed space X is strongly lacunary ward continuous if it preserves strongly lacunary quasi-Cauchy sequences of points in E; that is, $(f(x_k))$ is a strongly lacunary quasi-Cauchy sequence whenever (x_k) is strongly lacunary quasi-Cauchy. In this paper, not only strongly lacunary ward continuity, but also some other kinds of continuities are investigated in 2-normed spaces.

1. Introduction

Menger [1] introduced a notion called a generalized metric in 1928, and ten years later Vulich [2] defined a notion of a higher dimensional norm in linear spaces. Unfortunately, these studies had been neglected by many analysts for a long time. The concept of a 2-normed space was developed by Gähler in the middle of 1960s [3–5]. Since then, Mashadi [6], Gurdal [7], Mazaheri and Kazemi [8], and Sahiner [9] have studied this concept and obtained various results.

Using the main idea in the definition of sequential continuity, many kinds of continuities were introduced and investigated, and not all but some of them are in [10–13]. The concept of N_{θ} -convergence was introduced in [14] and further studied in [15]. Strongly lacunary ward continuity of a real function was introduced by Cakalli in [11] as named N_{θ} -ward continuity and further studied in [16].

The aim of this paper is to investigate strongly lacunary ward continuity in 2-normed spaces and prove interesting theorems.

2. Preliminaries

Throughout this paper, \mathbb{N} and \mathbb{R} will denote the set of all positive integers and the set of all real numbers, respectively. Now we recall the definition of a two-normed space. Let *X* be a real linear space with dim X > 1 and $\|\cdot, \cdot\| : X \times X \to \mathbb{R}$ a function. Then $(X, \|\cdot, \cdot\|)$ is called a linear 2-normed space if

(i) $||x, y|| = 0 \Leftrightarrow x$ and y are linearly dependent, (ii) ||x, y|| = ||y, x||, (iii) $||\alpha x, y|| = |\alpha| ||x, y||$, and (iv) $||x, y + z|| \le ||x, y|| + ||x, z||$ for $\alpha \in \mathbb{R}$ and $x, y, z \in X$. The function $||\cdot, \cdot||$ is called a 2-norm on X. Observe that in any 2-normed space $(X, ||\cdot, \cdot||)$ we have that $||\cdot, \cdot||$ is nonnegative, ||x-z, x-y|| = ||x-z, y-z||, and $||x, y + \alpha x|| = ||x, y||$ for all $x, y \in X$, $\alpha \in \mathbb{R}$. Throughout this paper by X we will mean a 2-normed space we may take $X = \mathbb{R}^2$ being equipped with the 2-norm

$$\|x_1, x_2\| = \operatorname{abs}\left(\begin{vmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{vmatrix}\right),$$
 (1)

where $x_1 = (x_{11}, x_{12}), x_2 = (x_{21}, x_{22}).$

A sequence (x_n) of points in X is said to converge to a point L of X in the 2-normed space X if $\lim_{n\to\infty} ||x_n-L,z|| = 0$ for every $z \in X$. This is denoted by $\lim_{n\to\infty} ||x_n,z|| = ||L,z||$. A sequence (x_n) of points in X is said to be a Cauchy sequence with respect to the 2-norm if $\lim_{n,m\to\infty} ||x_n - x_m, z|| = 0$ for every $z \in X$. A sequence of functions (f_n) is said to be uniformly convergent to a function f on a subset E of X if for each $\varepsilon > 0$, an integer N can be found such that $||f_n(x) - f(x), z|| < \varepsilon$ for $n \ge N$ and for all $x, z \in X$ [17]. A lacunary sequence $\theta = (k_r)$ is an increasing sequence of positive integers such that $k_0 = 0$ and $h_r = k_r - k_{r-1} \to \infty$ as $r \to \infty$. The intervals determined by θ will be denoted by $I_r = (k_{r-1}, k_r]$, and the ratio k_r/k_{r-1} will be abbreviated by q_r . A sequence (x_k) of points in *X* is called strongly lacunary convergent, or N_{θ} -convergent to an element *L* of *X* if

$$\lim_{r \to \infty} \frac{1}{h_r} \sum_{k \in I_r} \|x_k - L, z\| = 0,$$
(2)

for every $z \in X$ and it is denoted by $N_{\theta} - \lim_{k \to \infty} ||x_k, z|| = ||L, z||$ for every $z \in X$ [9]. Throughout the paper we will use the word " N_{θ} " instead of "strongly lacunary" and assume that $\liminf_r q_r > 1$.

3. Results

The set of seminorms $\{p_z\}$ defined by $p_z(x) = ||x, z||, \forall x$, for each $z \in X$, forms a locally convex topological vector space, and the topology formed by this family of seminorms gives the required topology on *X*. Since dim $X \ge 2$, for each $x \in X$ there exists a $y \in X$ such that *x* and *y* are linearly independent, and hence by $(i), p_y(x) = ||x, y|| \neq 0$. Thus the locally convex topological vector space induced by the set $\{p_z : z \in X\}$ of seminorms is Hausdorff so *X* is a Hausdorff space [18]. In this section, we investigate the concepts of a strongly lacunary quasi-Cauchy sequence and strongly lacunary ward continuity of a function on a 2-normed space.

Definition 1. A subset *E* of *X* is called N_{θ} -sequentially compact if any sequence of points in *E* has an N_{θ} -convergent sequence with an N_{θ} -limit in *E*.

We note that union of two N_{θ} -sequentially compact subsets of X is N_{θ} -sequentially compact, intersection of any N_{θ} -sequentially compact subsets is N_{θ} -sequentially compact, and any finite subset of X is N_{θ} -sequentially compact. Sum of N_{θ} sequentially compact subsets of X is N_{θ} -sequentially compact where sum of two subsets A and B is defined as $A+B = \{a+b : a \in A, b \in B\}$.

Definition 2. A function f defined on a subset E of X is said to be strongly lacunary sequentially continuous or N_{θ} sequentially continuous at a point x_0 of E if $(f(x_k))$ is an N_{θ} -convergent sequence to $f(x_0)$ whenever (x_k) is an N_{θ} convergent to x_0 sequence of points in E. If f is strongly lacunary sequentially continuous at every point of E, then it is said to be strongly lacunary sequentially continuous on E.

If a function f defined on a subset E of X is lacunary statistically sequentially continuous at a point x_0 , then $(f(x_k))$ is an N_{θ} -convergent sequence with $N_{\theta} - \lim ||f(x_k), z|| =$ $||f(x_0), z||$ for every $z \in X$ whenever (x_k) is an N_{θ} -convergent sequence with $N_{\theta} - \lim ||x_k, z|| = ||x_0, z||$ for every $z \in X$. We see that a function f defined on a subset E of X is strongly lacunary sequentially continuous if and only if it preserves strongly lacunary convergent sequences without stating limit of the sequence. We note that sum of two N_{θ} -sequentially continuous functions at a point x_0 of X is N_{θ} -sequentially continuous functions at a point x_0 of X is N_{θ} -sequentially continuous functions at a point x_0 of X is N_{θ} -sequentially continuous functions at a point x_0 of X is N_{θ} -sequentially continuous functions at a point x_0 of X is not sequentially continuous functions at a point x_0 of X is not sequentially continuous functions at a point x_0 of X is not sequentially continuous functions at a point x_0 of X is not sequentially continuous functions at a point x_0 of X is not sequentially continuous functions at a point x_0 of X is not sequentially continuous function is sequentially continuous; now we see that it is also true that not only uniform limit of sequentially continuous function is sequentially continuous, but also uniform limit of N_{θ} -sequentially continuous function is N_{θ} -sequentially continuous in 2-normed spaces. Now we give the latter in the following.

Theorem 3. Uniform limit of N_{θ} -sequentially continuous functions is N_{θ} -sequentially continuous.

Proof. Let (f_k) be a uniformly convergent sequence of each term defined on a subset *E* of *X* with uniform limit *f* and let (x_k) be any N_{θ} -convergent sequence of points in *E* with $N_{\theta} - \lim \|x_k, z\| = \|x, z\|$ for every $z \in X$. Take any $\varepsilon > 0$. By uniform convergence of (f_k) , there exists an $n_1 \in \mathbb{N}$ such that $\|f(x) - f_k(x), z\| < \varepsilon/3$ for $k \ge n_1$ and every $x \in E$ and $z \in X$. Hence,

$$\frac{1}{h_r}\sum_{k\in I_r} \left\| f\left(x\right) - f_k\left(x\right), z \right\| < \frac{1}{h_r} \left(k_r - k_{r-1}\right) \frac{\varepsilon}{3} = \frac{\varepsilon}{3}, \quad (3)$$

for $r \ge n_1$ and every $x \in E$ and $z \in X$. As f_{n_1} is N_{θ} -sequentially continuous on E, there exists an $n_2 \in \mathbb{N}$ such that, for $r \ge n_2$,

$$\frac{1}{h_r} \sum_{k \in I_r} \left\| f_{n_1}(x) - f_{n_1}(x_k), z \right\| < \frac{\varepsilon}{3},$$
(4)

for every $z \in X$. Now write $n_0 = \max\{n_1, n_2\}$. Thus for $r \ge n_0$ we have

$$\frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| f(x) - f(x_{k}), z \right\|
\leq \frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| v_{k}(x), z \right\| + \frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| f_{n_{1}}(x) - f_{n_{1}}(x_{k}), z \right\|
+ \frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| f_{n_{1}}(x_{k}) - f(x_{k}), z \right\|
< \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon,$$
(5)

where $v_k(x) = f(x) - f_{n_1}(x)$ for every $k \in \mathbb{N}$. Hence

$$\lim_{r \to \infty} \frac{1}{h_r} \sum_{k \in I_r} \| f(x) - f(x_k), z \| = 0.$$
 (6)

This completes the proof of the theorem.

Theorem 4. N_{θ} -sequentially continuous image of any N_{θ} -sequentially compact subset of X is N_{θ} -sequentially compact.

Proof. Assume that f is an N_{θ} -sequentially continuous function on a subset E of X and A is an N_{θ} -sequentially compact subset of E. Let $(f(x_n))$ be any sequence of points in f(A) where $x_n \in A$ for each positive integer n. N_{θ} -sequentially compactness of A implies that there is a subsequence $(\gamma_k) = (x_{n_k}) \text{ of } (x_n) \text{ with } N_{\theta} - \lim_{k \to \infty} ||\gamma_k, z|| = ||\ell, z|| \text{ for every } z \in E$. Write $(t_k) = (f(\gamma_k))$. As f is N_{θ} -sequentially continuous, $(f(\gamma_k))$ is N_{θ} -sequentially convergent which is a subsequence of the sequence $(f(x_n))$ with $N_{\theta} - \lim_{k \to \infty} ||t_k, z|| = ||\ell, z||$ for $\forall z \in E$. This completes the proof of the theorem.

The concept of a quasi-Cauchy sequence in a 2-normed space was studied in [19]. Now we give the following definition of an N_{θ} -quasi-Cauchy sequence.

Definition 5. A sequence (x_n) of points in a subset E of X is called N_{θ} -quasi-Cauchy if (Δx_k) is N_{θ} -convergent to 0, that is,

$$N_{\theta} - \lim_{k \to \infty} \left\| \Delta x_k, z \right\| = 0, \tag{7}$$

where $\Delta x_k = x_{k+1} - x_k$.

We note that any quasi-Cauchy sequence is N_{θ} -quasi-Cauchy, so any convergent sequence is N_{θ} -quasi-Cauchy in X. Any Cauchy sequence is N_{θ} -quasi-Cauchy, but the converse is not always true. However the converse is not always true, that is, there are N_{θ} -quasi-Cauchy sequences which are not convergent. Sum of two N_{θ} -quasi-Cauchy sequences is N_{θ} -quasi-Cauchy. Subsequence of an N_{θ} -quasi-Cauchy sequences needs not be N_{θ} -quasi-Cauchy. Now we give the definition of N_{θ} -ward compactness of a subset of X.

Definition 6. A subset E of X is called N_{θ} -ward compact if any sequence of points in E has an N_{θ} -quasi-Cauchy subsequence.

Union of two N_{θ} -ward compact subset of X is N_{θ} -ward compact, intersection of any N_{θ} -ward compact subsets is N_{θ} -ward compact, sum of two N_{θ} -ward compact subset of X is N_{θ} -ward compact, and any finite subset of X is N_{θ} -ward compact.

Definition 7. A function defined on a subset E of X is called N_{θ} -ward continuous if it preserves N_{θ} -quasi-Cauchy sequences, that is, $(f(x_k))$ is an N_{θ} -quasi-Cauchy sequence whenever (x_k) is.

We note that a composite of two N_{θ} -ward continuous functions is N_{θ} -ward continuous, and sum of two N_{θ} -ward continuous functions is N_{θ} -ward continuous.

Theorem 8. N_{θ} -ward continuous image of any N_{θ} -ward compact subset of X is N_{θ} -ward compact.

Proof. Assume that f is an N_{θ} -ward continuous function on a subset E of X and E is a N_{θ} -ward compact subset of A. Let $(f(x_n))$ be any sequence of points in f(E) where $x_n \in E$ for each positive integer n. N_{θ} -ward compactness of Eimplies that there is a subsequence $(\gamma_k) = (x_{n_k})$ of (x_n) with $N_{\theta} - \lim_{k \to \infty} ||\Delta \gamma_k, z|| = 0$ for every $z \in E$. Write $(t_k) = (f(\gamma_k))$. As f is N_{θ} -ward continuous, $(f(\gamma_k))$ is N_{θ} -quasi-Cauchy which is a subsequence of the sequence $(f(x_n))$ with $N_{\theta} - \lim_{k \to \infty} ||\Delta t_k, z|| = 0$ for $\forall z \in E$. This completes the proof of the theorem. \Box

Corollary 9. N_{θ} -ward continuous image of any compact subset of X is N_{θ} -ward compact.

Proof. The proof follows from the preceding theorem. \Box

A function f defined on a subset E of X is sequentially continuous at x_0 , if for any sequence (x_n) of points in E converging to x_0 , we have $(f(x_n))$ converges to $f(x_0)$. f is sequentially continuous on E if it is sequentially continuous at every point of E (see [19] for the infinite dimensional case

and [20] for the finite dimensional case). Concerning N_{θ} -quasi-Cauchy sequences, N_{θ} convergent sequences, and convergent sequences the problem arises to investigate the following types of continuity of functions on X. In the following N_{θ} , ΔN_{θ} , and c will denote the set of N_{θ} -convergent sequence, the set of all N_{θ} -quasi-Cauchy sequences, and the set of convergent sequences of points in X, respectively:

(1)
$$(x_k) \in \Delta N_\theta \Rightarrow (f(x_k)) \in \Delta N_\theta$$
,

(2)
$$(x_k) \in \Delta N_\theta \Rightarrow (f(x_k)) \in c$$

- (3) $(x_k) \in c \Rightarrow (f(x_k)) \in c$,
- (4) $(x_k) \in c \Rightarrow (f(x_k)) \in \Delta N_{\theta}$,
- (5) $(x_k) \in N_{\theta} \Rightarrow (f(x_k)) \in N_{\theta}$,
- (6) $(x_k) \in \Delta N_{\theta} \Rightarrow (f(x_n))$ is quasi-Cauchy,
- (7) $(x_k) \in \Delta N_\theta \Rightarrow (f(x_n)) \in N_\theta.$

We see that (1) is N_{θ} -ward continuity of f, (5) is N_{θ} -sequentially continuity of f, and (3) is the ordinary sequential continuity of f. It is easy to see that (2) implies (1), and (1) does not imply (2); and (1) implies (4), and (4) does not imply (1); (2) implies (3), and (3) does not imply (2). (6) implies (1), but (1) does not imply (6) and (7) implies (1), but (1) does not imply (7). Now we give that the implication (1) implies (5), that is, any N_{θ} -ward continuous function is N_{θ} -sequentially continuous.

Theorem 10. If f is N_{θ} -ward continuous function on a subset E of X, then it is N_{θ} -sequentially continuous on E.

Proof. Assume that f is an N_{θ} -ward continuous function on a subset E of X. Let (x_n) be any N_{θ} -convergent sequence in A with $N_{\theta} - \lim_{k \to \infty} ||x_k, z|| = ||x_0, z||$ for all $z \in X$. Then the sequence $x = (x_n)$ defined by

$$x_n = \begin{cases} x_k, & \text{if } n = 2k - 1 \text{ for a positive integer } k \\ x_0, & \text{if } n \text{ is even} \end{cases}$$
(8)

is also N_{θ} -convergent to x_0 . Hence it is N_{θ} -quasi-Cauchy sequence. As f is N_{θ} -ward continuous on E, the transformed sequence (y_n) obtained by

$$y_n = \begin{cases} f(x_k), & \text{if } n = 2k - 1 \text{ for a positive integer } k \\ f(x_0), & \text{if } n \text{ is even} \end{cases}$$
(9)

is also N_{θ} -quasi-Cauchy. Thus,

$$\lim_{r \to \infty} \frac{1}{h_r} \sum_{k \in I_r} \left\| f\left(x_k\right) - f\left(x_0\right), z \right\| = 0$$
(10)

for all $z \in X$. It follows from this that the sequence $(f(x_k))$ is N_{θ} convergent to $(f(x_0))$.

The converse of this theorem is not valid in general, a counterexample can be easily constructed via the function $f(x, y) = (x^2, y^2)$ on the 2-normed space \mathbb{R}^2 with the usual 2-norm.

Theorem 11. If (f_n) is a sequence of N_{θ} -ward continuous functions on a subset E of X and (f_n) is uniformly convergent to a function f, then f is N_{θ} -ward continuous on E.

Proof. Let (x_k) be any N_{θ} -quasi-Cauchy sequence of points in *E*, and let ε be any positive real number. By uniform convergence of (f_k) , there exists an $n_1 \in \mathbb{N}$ such that $||f(x) - f_k(x), z|| < \varepsilon/3$ for $k \ge n_1$ and every $x \in E$ and $z \in X$. Hence,

$$\frac{1}{h_r} \sum_{k \in I_r} \| f(x) - f_k(x), z \| < \frac{1}{h_r} \left(k_r - k_{r-1} \right) \frac{\varepsilon}{3} = \frac{\varepsilon}{3}, \quad (11)$$

for $r \ge n_1$ and every $x \in E$ and $z \in X$. As f_{n_1} is N_{θ} -ward continuous on E, there exists an $n_2 \in \mathbb{N}$ such that, for $r \ge n_2$,

$$\frac{1}{h_r} \sum_{k \in I_r} \left\| f_{n_1}(x_{k+1}) - f_{n_1}(x_k), z \right\| < \frac{\varepsilon}{3},$$
(12)

for every $z \in X$. Now write $n_0 = \max\{n_1, n_2\}$. Thus, for $r \ge n_0$, we have

$$\frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| f\left(x_{k+1}\right) - f\left(x_{k}\right), z \right\| \\
\leq \frac{1}{h_{r}} \sum_{k \in I_{r}} f \left\| (x_{k+1}) - f_{n_{1}}\left(x_{k+1}\right), z \right\| \\
+ \frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| f_{n_{1}}\left(x_{k+1}\right) - f_{n_{1}}\left(x_{k}\right), z \right\| \\
+ \frac{1}{h_{r}} \sum_{k \in I_{r}} \left\| f_{n_{1}}\left(x_{k}\right) - f\left(x_{k}\right), z \right\| \\
< \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon.$$
(13)

Hence,

$$\lim_{r \to \infty} \frac{1}{h_r} \sum_{k \in I_r} \| f(x_{k+1}) - f(x_k), z \| = 0.$$
(14)

Thus f preserves N_{θ} -quasi-Cauchy sequences. This completes the proof of the theorem.

4. Conclusion

In this paper, we investigate strongly lacunary continuity and some other kinds of continuities defined via a lacunary sequence and we prove interesting theorems related to these kinds of continuities. The results in this paper are extensively deeper than existing related results in the literature. We note that the notion of a strongly lacunary quasi-Cauchy sequence coincides with the notion of a strongly lacunary convergent sequence in a complete non-Archimedean 2-normed space, and so the set of strongly lacunary ward continuous functions coincides with the set of strongly lacunary sequentially continuous functions in a complete non-Archimedean 2-normed space (see [21] for the related concepts in an ultrametric field). For a further study, we suggest to investigate strongly lacunary quasi-Cauchy sequences of points for the fuzzy functions in a 2-normed fuzzy spaces. However, due to the change in settings, the definitions and methods of proofs will not always be analogous to those of the present work (see [22, 23] for the definitions and related concept in fuzzy setting). We note that the study in this paper can be generalized to *n*-normed spaces as another further study.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- K. Menger, "Untersuchungen über allgemeine Metrik," *Mathematische Annalen*, vol. 100, no. 1, pp. 75–163, 1928.
- [2] B. Vulich, "On a generalized notion of convergence in a Banach space," *Annals of Mathematics*, vol. 38, no. 1, pp. 156–174, 1937.
- [3] S. G. Gähler, "2-metrische Raume und ihre topologische Struktur," *Mathematische Nachrichten*, vol. 26, pp. 115–148, 1963.
- [4] S. G. Gähler, "Lineare 2-normietre Raume," Mathematische Nachrichten, vol. 28, pp. 1–43, 1965.
- [5] S. G. Gähler, "Uber der Uniformisierbarkeit 2-metrische Raume," Mathematische Nachrichten, vol. 28, pp. 235–244, 1965.
- [6] H. G. Mashadi, "On finite dimensional 2-normed spaces," Soochow Journal of Mathematics, vol. 27, no. 3, pp. 321–329, 2001.
- [7] M. Gurdal, "On ideal convergent sequences in 2-normed spaces," *Thai Journal of Mathematics*, vol. 4, no. 1, pp. 85–91, 2006.
- [8] H. Mazaheri and R. Kazemi, "Some results on 2-inner product spaces," *Novi Sad Journal of Mathematics*, vol. 37, no. 2, pp. 35– 40, 2007.
- [9] A. Sahiner, "On I-Lacunary strong convergence in 2-normed space," *International Journal of Contemporary Mathematical Sciences*, vol. 2, no. 20, pp. 991–998, 2007.
- [10] D. Burton and J. Coleman, "Quasi-cauchy sequences," American Mathematical Monthly, vol. 117, no. 4, pp. 328–333, 2010.
- H. Cakalli, "N_θ-ward continuity," Abstract and Applied Analysis, vol. 2012, Article ID 680456, 8 pages, 2012.
- [12] I. Canak and M. Dik, "New types of continuities," Abstract and Applied Analysis, vol. 2010, Article ID 258980, 6 pages, 2010.
- [13] R. W. Vallin, "Creating slowly oscillating sequences and slowly oscillating continuous functions," *Acta Mathematica Universitatis Comenianae*, vol. 80, no. 1, pp. 71–78, 2011.
- [14] A. R. Freedman, J. J. Sember, and M. Raphael, "Some Cesarotype summability spaces," *Proceedings of the London Mathematical Society*, vol. 3, no. 37, pp. 508–520, 1978.
- [15] I. H. Jebril, "A generalization of strongly Cesaro and strongly Lacunary summable spaces," *Acta Universitatis Apulensis*, vol. 23, pp. 49–61, 2010.
- [16] H. Cakalli and H. Kaplan, "A study on N_{θ} -quasi-Cauchy sequences," *Abstract and Applied Analysis*, vol. 2013, Article ID 836970, 4 pages, 2013.
- [17] R. Freese and Y. J. Cho, Geometry of Linear 2-Normed Spaces, Nova Science, Hauppauge, NY, USA, 2001.
- [18] N. Srivastava, S. Bhattacharya, and S. N. Lal, "2-Normed algebras-I," *Publications de l'Institut Mathematique*, vol. 88, no. 102, pp. 111–121, 2010.
- [19] H. Cakalli and S. Ersan, "On sequences in 2-normed spaces," http://arxiv.org/pdf/1306.2469.pdf.

- [20] S. Talebi and S. Sarabadan, "Statistical convergence and ideal convergence of sequences of functions in 2-normed spaces," *International Journal of Mathematics and Mathematical Sciences*, vol. 2011, Article ID 517841, 10 pages, 2011.
- [21] P. N. Natarajan, "An introduction to ultrametric summability theory," in *Springer Briefs in Mathematics*, 2014.
- [22] M. Mursaleen and S. A. Mohiuddine, "On lacunary statistical convergence with respect to the intuitionistic fuzzy normed space," *Journal of Computational and Applied Mathematics*, vol. 233, no. 2, pp. 142–149, 2009.
- [23] Lj. D. R. Kočinac, "Selection properties in fuzzy metric spaces," *Filomat*, vol. 26, no. 2, pp. 305–312, 2012.



The Scientific World Journal





Decision Sciences







Journal of Probability and Statistics



Hindawi Submit your manuscripts at http://www.hindawi.com



(0,1),

International Journal of Differential Equations





International Journal of Combinatorics





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society







Function Spaces



International Journal of Stochastic Analysis



Journal of Optimization